UNCLASSIFIED

ORNL 416
PHYSICS-GENERAL
SERIES B:
cy 7

MARTIN MARIETTA ENERGY SYSTEMS LIBRARIES
3 4456 0360537 4

ORNL SUMMER SHIELDING SESSION

GALE YOUNG, Leader

ON STRAIGHT-AHEAD γ-TRANSMISSION WITH A MINIMUM IN THE CROSS-SECTION

GALE YOUNG

CENTRAL RESEARCH LIBRARY
DOCUMENT COLLECTION
LIBRARY LOAN COPY
DO NOT TRANSFER TO ANOTHER PERSON
If you wish someone else to see this document, send in name with document and the library will arrange a loan.

OAK RIDGE NATIONAL LABORATORY

OPERATED BY
CARBIDE AND CARBON CHEMICALS CORPORATION
FOR THE
ATOMIC ENERGY COMMISSION
POST OFFICE BOX P
OAK RIDGE, TENNESSEE

UNCLASSIFIED

UNCLASSIFIED

ORNL-416
Series B

Contract No. W-7405, eng 26 *

Copy# 7

ORNL Summer Shielding Session

Gale Young, Leader

ON STRAIGHT-AHEAD γ-TRANSMISSION WITH A MINIMUM IN THE CROSS-SECTION

Gale Young

Date Issued: SEP 26 1950

OAK RIDGE NATIONAL LABORATORY
Operated By
Carbide and Carbon Chemicals Corporation
for the
Atomic Energy Commission
Post Office Box P
Oak Ridge, Tennessee

*Explanation on page 3.

UNCLASSIFIED

MARTIN MARIETTA ENERGY SYSTEMS LIBRARIES
3 4456 0360537 4

ORNL-416
Physics-General
Series B:

OAK RIDGE NATIONAL LABORATORY INTERNAL DISTRIBUTION:

1. G. T. Felbeck (C&CCC)
2. 706-A Library
3. 706-A Library
4. 706-B Library
5. Biology Library
6. Health Physics Library
7. Training School Library
8. Training School Library
9. Central Files
10. Central Files
11. Central Files
12. Central Files
13. J. A. Swartout
14. J. H. Gillette
15. A. S. Householder
16. J. H. Frye, Jr.
17. E. J. Murphy
18. M. D. Peterson
19. C. N. Rucker
20. W. D. Lavers
21. A. M. Weinberg
22. J. A. Lane
23. E. P. Blizard
24. R. N. Lyon
25. M. M. Mann
26. Gale Young
27. A. Hollaender

OAK RIDGE NATIONAL LABORATORY EXTERNAL DISTRIBUTION:

28-35. Argonne National Laboratory
36. Armed Forces Special Weapons Project
37-38. Atomic Energy Commission, Washington
39. Battelle Memorial Institute
40-47. Brookhaven National Laboratory
48. Bureau of Medicine and Surgery
49. Bureau of Ships
50. Bureau of Standards
51. Bureau of Aeronautics
52-55. Carbide & Carbon Chemicals Corp.(K-25)
56-59. Carbide & Carbon Chemicals Corp.(Y-12)
60. Chicago Operations Office
61. Cleveland Area Office
62. Columbia University (Dunning)
63. Columbia University (Failla)
64. Dow Chemical Company
65-70. General Electric Company, Richland
71. Hanford Operations Office
72. Idaho Operations Office
73-74. Iowa State College
75. Kansas City
76-79. Knolls Atomic Power Laboratory
80-82. Los Alamos
83. Mallinckrodt Chemical Works
84. Massachusetts Institute of Technology (Gaudin)
85. Massachusetts Institute of Technology (Kaufmann)
86. Massachusetts Institute of Technology
87-89. Mound Laboratory
90-91. National Advisory Committee for Aeronautics
92-93. National Bureau of Standards
94-95. Naval Radiological Defense Laboratory
96-97. NEPA Project
98. New Brunswick Laboratory
99-103. New York Operations Office
104. North American Aviation, Inc.
105-112. Nuclear Development Associates
113. Patent Advisor, Washington
114-116. Rand Corporation
117. Reactor Development Div. (AEC, Washington)
118. Sandia Base
119. Sylvania Electric Products, Inc.
120. U.S. Public Health Service
121. UCLA Medical Research Lab. (Warren)
122-126. University of California Radiation Laboratory
127-128. University of Rochester
129. University of Washington
130-131. Western Reserve University (Friedell)
132-135. Westinghouse
136. Los Alamos

This report is one of a series on shielding and represents part of the work of a summer group working principally at Oak Ridge National Laboratory. Participants are drawn from Nuclear Development Associates (Navy contract), NEPA, Bureau of Ships and Bureau of Aeronautics (Navy Department), AEC, RAND, General Electric, Argonne and Oak Ridge National Laboratory.

These reports are issued through ORNL on Contract No. W-7405, eng 26.

[redacted]

ON STRAIGHT-AHEAD γ- TRANSMISSION WITH A MINIMUM IN THE CROSS-SECTION

Gale Young

Under certain assumptions about the law of scattering and the variation of the cross-sections with energy, Greuling has obtained solutions of the attenuation problem (1, 2, 3). His results, as they stand, do not include instances where the cross-section can (as with γ rays) go through a minimum as the energy varies. The present note illustrates how different Greuling solutions can be pieced together to construct cases having a minimum.

Introduction

We give here only a brief sketch of the problem, referring to the above references for details. Let σ = scattering cross section, μ = total cross section. It is assumed that scattering distributes the energy of the emerging γ rays (we shall here write in terms of photons, though material particles such as neutrons can be discussed in exactly the same way) uniformly over lower energies, and that all the rays move straight ahead undeviated by scattering. This last assumption leads, of course, to an over-estimate of the penetration.

Let $F(x, E)$ be the number of photons per unit energy range at distance x. Then, with the above assumptions,

$$\frac{\delta F}{\delta x} + \mu F = \int_E^{E_o} F(x, t) \frac{\sigma(t)}{t} dt + S(x) \tag{1}$$

The source term $S(x)$ arises from scattering out of the primary or other beams above energy E_o.

A particularly simple case is when $\frac{d\mu}{dE} = -\frac{\sigma(E)}{E}$. Then (1) can be satisfied by a function of x alone, becoming

$$F' + \mu F = F \int_E^{E_o} \frac{\sigma(t)}{t}\, dt + S = (\mu - \mu_o) F + S$$

or

$$F' + \mu_o F = S \quad . \tag{2}$$

Thus all the properties of the medium disappear except μ_o, and it makes no difference how much of μ_o is contributed by scattering and how much by true absorption. The transmission (for secondaries) of the medium is just as if it had constant absorption equal to μ_o, and no scattering power at all.

In the case of a primary beam at E_o we have

$$S = \frac{\sigma_o}{E_o} e^{-\mu_o x} \tag{3}$$

and (2) has the familiar solution

$$F = x S = \frac{\sigma_o x}{E_o} e^{-\mu_o x} \quad . \tag{4}$$

The result does, of course, depend upon the strength of the scattering out of the primary beam.

The total number of photons penetrating a distance x is given by

$$Q(x) = e^{-\mu_o x} + \int_0^{E_o} F\, dE = (1 + \sigma_o x)\, e^{-\mu_o x} \quad , \tag{5}$$

so that the "buildup factor" is $1 + \sigma_o x$. As we have just seen, the quantity σ_o enters only from the primary beam and not at all from the secondaries.

First Example

Consider now the same situation, except that different cross-section values apply for the primary beam, as indicated in the following sketch.



The source term in (2) is now

$$S = \frac{\sigma_1}{E_o} e^{-\mu_1 x} , \tag{6}$$

and the solution is

$$F = \frac{\sigma_1}{E_o} \frac{e^{-\mu_o x} - e^{-\mu_1 x}}{u_1 - \mu_o} \tag{7}$$

with

$$Q = e^{-\mu_1 x} + \frac{\sigma_1}{\mu_1 - \mu_o} \left(e^{-\mu_o x} - e^{-u_1 x} \right) . \tag{8}$$

Considering $\mu_1 > \mu_o$, we write (8) as

$$Q(x) = \left[\frac{\sigma_1}{\mu_1 - \mu_o} + e^{-(\mu_1 - \mu_o)x} \left\{ 1 - \frac{\sigma_1}{\mu_1 - \mu_o} \right\} \right] e^{-\mu_o x} \tag{9}$$

where the expression in brackets is the build-up factor referred to the exponential taken at the minimum cross-section; it is unity at $x = 0$ and $\frac{\sigma_1}{\mu_1 - \mu_o}$ for x large. Thus, depending on the constants, it can be either a "build-up" or "build-down" factor. If $\sigma_1 = \mu_1 - \mu_o$, then $Q(x) = e^{-\mu_o x}$ everywhere.

Second Example

We now consider a slightly more complicated case in which the energy range is divided into two regions by $E_c < E_o$. In the top region we shall use a Greuling case with a rising cross-section, and in the bottom region a Greuling case with a falling cross-section.

One of the simplest assumptions for the top region is to take

$\frac{d\mu}{dE} = + \frac{\sigma(E)}{E}$. This is the Greuling case $\alpha = -1$ (see above references), and for a source term (3) on the right side of (1) the solution is

$$F(x, E) = \frac{\sigma_o}{E_o} \times e^{-\mu(E)x} , \tag{10}$$

as may be readily verified.

In the bottom region we shall take $\frac{d\mu}{dE} = -\sigma/E$, which is the case described in the introductory section. It may be noted that our assumptions regarding the top and bottom regions include the case with continuous $1/E$ scattering, absorption beginning at E_c and rising hyperbolically with energy, and total cross-section continuous (though with discontinuous derivative) at E_c; as shown in the adjacent sketch



The source fed into the bottom region below E_c is given by

$$\frac{\sigma_o}{E_o} e^{-\mu_o x} + \int_{E_c}^{E_o} F(x, E) \frac{\sigma(E)}{E} dE \tag{11}$$

which with (10) comes out to be

$$\frac{\sigma_o}{E_o} e^{-\mu_c x} . \tag{12}$$

Then, from (4), the solution for $E < E_c$ is

$$F = \frac{\sigma_o}{E_o} x e^{-\mu_c x} ; \tag{13}$$

provided, as assumed here, that μ is continuous across E_c.

The integral fluxes are

$$\left.\begin{aligned}
&\text{(primary)} \quad Qp = e^{-\mu_o x} \\
&(E_c < E < E_o) \quad Q_1 = \frac{\sigma_o x}{E_o} \int_{E_c}^{E_o} e^{-\mu(E)x} dE \\
&(E < E_c) \quad Q_2 = \frac{E_c}{E_o} \sigma_o x e^{-\mu_c x} .
\end{aligned}\right\} \tag{14}$$

The second of these may be written as

$$Q_1 = \frac{\sigma_o x}{E_o} \left(\frac{E}{\sigma}\right)_m \int_{E_c}^{E_o} e^{-\mu x} \frac{\sigma(E)}{E} dE$$

$$= \frac{\sigma_o}{E_o} \left(\frac{E}{\sigma}\right)_m \left(e^{-\mu_c x} - e^{-\mu_o x}\right) , \tag{15}$$

where $\left(\frac{E}{\sigma}\right)_m$ is evaluated at some mean energy E_m between E_c and E_o. If E/σ is an increasing function of E, then $Q_1 < \left(e^{-u_c x} - e^{-u_o x}\right)$. If μ is an increasing function of E, then for x large most of the value of the integral in (15) comes from near the lower limit, and thus asymptotically m tends to c. For x large the dominant term in $Q = Q_p + Q_1 + Q_2$ is Q_2 as given in (14).

Constant Total Cross-section

The above examples have a downward pointed cusp in μ at the minimum, and so presumably give too small a buildup factor. To approach the picture from the other side we consider the case where μ is flat.

A result due to Wigner (4; 3, page 11) can be generalized by considering the Greuling result as α becomes infinite (3, p.15 and 18); α is the coefficient in the Greuling condition $\alpha \frac{d\mu}{dE} = -\sigma/E$. Let μ be constant, and let σ be an arbitrary function of E. Then, as may be verified in a straight-forward manner by using the recurrence relations among the Bessel functions, the solution of (1) with a source $\frac{\sigma_o}{E_o} e^{-\mu x}$ is

$$F(x, E) = \frac{\sigma_o}{E_o} x e^{-\mu x} \frac{I_1\left(2\sqrt{x\varphi}\right)}{\sqrt{x\varphi}} \tag{16}$$

where

$$\left.\begin{aligned} \varphi(E) &= \int_E^{E_o} \frac{\sigma(t)}{t} dt \\ \varphi'(E) &= -\frac{\sigma(E)}{E} \end{aligned}\right\} \tag{17}$$

The integral secondary flux above energy E_2 is

$$\int_{E_2}^{E_o} F(x, E)\, dE = \frac{\sigma_o}{E_o}\, x\, e^{-\mu x} \left(\frac{E}{\sigma}\right)_m \int_{E_2}^{E_o} \frac{I_1(2\sqrt{x\varphi})}{\sqrt{x\varphi}}\, \frac{\sigma}{E}\, dE$$

$$= \frac{\sigma_o}{E_o} \left(\frac{E}{\sigma}\right)_m e^{-\mu x} \left[I_o\left(2\sqrt{x\varphi_2}\right) - 1 \right] . \qquad (18)$$

For x large most of the integral comes from near the bottom limit, and so asymptotically E_m approaches E_2.

In this case there is, for x large, a $e^{\sqrt{kx}}$ factor in the buildup.

Third Example

We now insert a flat μ central region between the upper and lower regions of the second example above, obtaining



a shape as sketched.

Working down from the initial energy E_o, and letting μ without a subscript denote the central value $\mu = \mu_1 = \mu_2$, we have

(primary) $$Q\,p = e^{-\mu_o x}$$

(region #1) $$S_1 = \frac{\sigma_o}{E_o}\, e^{-\mu_o x}$$

$$F_1 = \frac{\sigma_o}{E_o}\, x\, e^{-\mu(E)x} \qquad \text{from (10)}$$

$$Q_1 = \frac{\sigma_o}{E_o}\left(\frac{E}{\sigma}\right)_{m_1}\left(e^{-\mu x} - e^{-\mu_o x}\right) \qquad \text{from (15)}$$

(region #2) $$S_2 = \frac{\sigma_o}{E_o}\, e^{-\mu x} \qquad \text{from (12)}$$

$$F_2 = \frac{\sigma_o}{E_o}\, x\, e^{-\mu x}\, \frac{I_1\,(2\sqrt{x\varphi})}{\sqrt{x\varphi}} \qquad \text{from (16)}$$

$$Q_2 = \frac{\sigma_o}{E_o}\left(\frac{E}{\sigma}\right)_{m_2} e^{-\mu x}\left[I_o(2\sqrt{x\varphi_2}) - 1\right] \qquad \text{from (18)}$$

(region #3) Here we find

$$S_3 = S_2 + \int_{E_2}^{E_1} F_2(x,E)\,\frac{\sigma}{E}\,dE$$

$$= S_2 + \frac{\sigma_o}{E_o}\, e^{-\mu x}\left[I_o(2\sqrt{x\varphi_2}) - 1\right]$$

$$= \frac{\sigma_o}{E_o}\, e^{-\mu x}\, I_o\,(2\sqrt{x\varphi_2}) \qquad (19)$$

Then, with (2), we have $F' + \mu F = S_3$, of which the solution is

$$F_3 = \frac{\sigma_o}{E}\, x\, e^{-\mu x}\, \frac{I_1\,(2\sqrt{x\varphi_2})}{\sqrt{x\varphi_2}} \qquad (20)$$

$$Q_3 = \frac{E_2}{E_o}\,\sigma_o\, x\, e^{-\mu x}\, \frac{I_1\,(2\sqrt{x\varphi_2})}{\sqrt{x\varphi_2}} \qquad (21)$$

For x large, Q_3 is ultimately the dominant term in the total Q, namely

$$\frac{E_2}{E_o}\frac{\sigma_o}{\varphi_2}\frac{1}{2\sqrt{\pi}}\left(\varphi_2 x\right)^{\frac{1}{4}} e^{-\mu x + 2\sqrt{\varphi_2 x}} \tag{22}$$

- - - - -

References

1) E. Greuling in Mon P-172, p. 23-28

2) E. Greuling in Mon P-130, p. 7-9

3) G. Young, Mon P-293

4) E. P. Wigner, CH-137

GY;lg
8/12/49