ORNL-6086/V1
(First Volume
of ORNL-6086)

Physics Division

ATOMIC DATA FOR FUSION

VOLUME 1

COLLISIONS OF H, H_2, He and Li ATOMS and IONS with ATOMS and MOLECULES

C. F. Barnett
Oak Ridge National Laboratory

Published and Edited by
Controlled Fusion Atomic Data Center

H. T. Hunter and M. I. Kirkpatrick
Series Technical Editors

I. Alvarez, C. Cisneros, University of Mexico,
and R. A. Phaneuf
Contributing Editors

Date Published - July 1990

Prepared for the
Office of Fusion Energy
U.S. Department of Energy
Washington, DC 20545
Budget Activity No. AT 05 30 04 0

Prepared by the
OAK RIDGE NATIONAL LABORATORY
Oak Ridge, Tennessee 37831-6372
operated by
MARTIN MARIETTA ENERGY SYSTEMS, INC.
for the
U. S. DEPARTMENT OF ENERGY
under contract DE-AC05-84OR21400

MARTIN MARIETTA ENERGY SYSTEMS LIBRARIES

3 4456 0315830 2

Printed in the United States of America. Available from
National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road, Springfield, Virginia 22161
NTIS price codes—Printed Copy: A-99 Microfiche A01

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

ATOMIC DATA FOR FUSION

VOL. 1

COLLISIONS OF H, H_2, He, AND Li ATOMS AND IONS WITH ATOMS AND MOLECULES

Foreword xiii

Series Preface xv

Abstract xvii

Introduction I-1

A. Electron Capture Collisions

$H + H \rightarrow H^- + H^+$ A-2
$H + H_2 \rightarrow H^- + H_2^+$ A-8
$H + He \rightarrow H^- + He^+$ A-10
$H + Li \rightarrow H^- + Li^+$ A-16
$H + H^- \rightarrow H^- + H$ A-20

$H^+ + H \rightarrow H + H^+$ A-22
$H^+ + H_2 \rightarrow H$ A-28
$H^+ + H_2 \rightarrow H^-$ A-30
$H^+ + He \rightarrow H + He^+$ A-32
$H^+ + He \rightarrow H^- + He^{2+}$ A-38
$H^+ + Li \rightarrow H + Li^+$ A-40
$H^+ + H^- \rightarrow H^- + H^+$ A-44
$H^+ + H^- \rightarrow 2H$ A-46
$H^+ + He^+ \rightarrow \sigma(He^{2+}$ Total) A-48
$H^+ + He^+ \rightarrow H + He^{2+}$ A-54

$H_2^+ + H \rightarrow H_2 + H^+$ A-58
$H_2^+ + H_2 \rightarrow H_2$ A-58
$H_2^+ + He \rightarrow H_2 + He^+$ A-58
$H_2^+(\upsilon) + H_2 \rightarrow H_2$ A-60

$He^+ + H \rightarrow He + H^+$ A-62
$He^+ + H^- \rightarrow He + H$ A-86
$He^+ + H_2 \rightarrow He$ A-68
$He^+ + He \rightarrow He + He^+$ A-70
$He^+ + Li \rightarrow He + Li^+$ A-76
$He^+ + He^+ \rightarrow He + He^{2+}$ A-86

$He^{2+} + He^+ \rightarrow He^+ + He^{2+}$(Partial & Total) A-84
$He^{2+} + H^- \rightarrow He^+ + H$ A-86
$He^{2+} + H \rightarrow He^+ + H^+$ A-88
$He^{2+} + H_2 \rightarrow He^+$ A-94
$He^{2+} + H_2 \rightarrow He$ A-96
$He^{2+} + He \rightarrow He^+ + He^+$ A-98
$He^{2+} + He \rightarrow He + He^{2+}$ A-104
$He^{2+} + Li \rightarrow He^+ + Li^+$ A-110
$He^{2+} + Li \rightarrow He + Li^{2+}$ A-114

B. Electron Capture into Excited States

$H^+ + H \rightarrow H(2s) + H^+$ B-2
$H^+ + H \rightarrow H(2p) + H^+$ B-8
$H^+ + H_2 \rightarrow H(2s)$ B-14
$H^+ + H_2 \rightarrow H(2p)$ B-14
$H^+ + H_2 \rightarrow H(\text{Lyman-}\alpha\ \text{Total})$ B-16
$H^+ + H_2 \rightarrow H(\text{Balmer-}\alpha)$ B-18
$H^+ + H_2 \rightarrow H(\text{Balmer-}\beta)$ B-18
$H^+ + H_2 \rightarrow H(3s)$ B-20
$H^+ + H_2 \rightarrow H(3p)$ B-20
$H^+ + H_2 \rightarrow H(3d)$ B-20
$H^+ + He \rightarrow H(2s) + He^+$ B-22
$H^+ + He \rightarrow H(2p) + He^+$ B-28
$H^+ + He \rightarrow H(\text{Balmer-}\alpha) + He^+$ B-34
$H^+ + He \rightarrow H(3s) + He^+$ B-40
$H^+ + He \rightarrow H(3p) + He^+$ B-46
$H^+ + He \rightarrow H(3d) + He^+$ B-52
$H^+ + Li \rightarrow H(2s) + Li^+$ B-58
$H^+ + Li \rightarrow H(2p) + Li^+$ B-58
$H^+ + Li \rightarrow H(3s) + Li^+$ B-60
$H^+ + Li \rightarrow H(3p) + Li^+$ B-60
$H^+ + Li \rightarrow H(3d) + Li^+$ B-60

$He^+ + H_2 \rightarrow H(2^1S + 2^3S)$ B-62
$He^+ + H_2 \rightarrow He(2^1P)$ B-64
$He^+ + H_2 \rightarrow He(3^1P)$ B-64
$He^+ + H_2 \rightarrow He(\lambda = 447.1\ \text{nm})$ B-66
$He^+ + H_2 \rightarrow He(\lambda = 587.6\ \text{nm})$ B-66
$He^+ + H_2 \rightarrow He(\lambda = 667.8\ \text{nm})$ B-66
$He^+ + He \rightarrow He(2^1S + 2^3S) + He^+$ B-68
$He^+ + He \rightarrow He(2^1P) + He^+$ B-74
$He^+ + He \rightarrow He(3^1S) + He^+$ B-80
$He^+ + He \rightarrow He(3^1P) + He^+$ B-80
$He^+ + He \rightarrow He(3^1D) + He^+$ B-80
$He^+ + He \rightarrow He(3^3S) + He^+$ B-82
$He^+ + He \rightarrow He(3^3P) + He^+$ B-82
$He^+ + He \rightarrow He(3^3D) + He^+$ B-82
$He^+ + He \rightarrow He(4^1S) + He^+$ B-84
$He^+ + He \rightarrow He(4^1P) + He^+$ B-84
$He^+ + He \rightarrow He(4^1D) + He^+$ B-84
$He^+ + He \rightarrow He(4^3S) + He^+$ B-86
$He^+ + He \rightarrow He(4^3P) + He^+$ B-86
$He^+ + He \rightarrow He(4^3D) + He^+$ B-86
$He^+ + Li \rightarrow He(53.7\ \text{nm}) + Li^+$ B-88
$He^+ + Li \rightarrow He(58.4\ \text{nm}) + Li^+$ B-88

$He^{2+} + H \rightarrow He^+(2s) + H^+$ B-90
$He^{2+} + H \rightarrow He^+(2p) + H^+$ B-96
$He^{2+} + H \rightarrow He^+(\text{Lyman-}\alpha) + H^+$ B-102
$He^{2+} + H \rightarrow He^+(\text{Lyman-}\beta) + H^+$ B-102
$He^{2+} + H \rightarrow He^+(\text{Lyman-}\gamma) + H^+$ B-102
$He^{2+} + H \rightarrow He^+(\text{Balmer-}\alpha) + H^+$ B-102
$He^{2+} + H \rightarrow He^+(3p) + H^+$ B-104
$He^{2+} + H \rightarrow He^+(3s+3d) + H^+$ B-104
$He^{2+} + H \rightarrow He^+(4p) + H^+$ B-104
$He^{2+} + H_2 \rightarrow He^+(2s)$ B-106
$He^{2+} + H_2 \rightarrow He^+(2p)$ B-106
$He^{2+} + H_2 \rightarrow He^+(\text{Lyman-}\alpha)$ B-108
$He^{2+} + H_2 \rightarrow He^+(\text{Lyman-}\beta)$ B-108

B. Electron Capture into Excited States (Cont'd)

He^{2+} + H_2 -> He^+(Lyman-γ) B-108
He^{2+} + H_2 -> He^+(Balmer-α) B-108
He^{2+} + H_2 -> He^+(3p) B-110
He^{2+} + H_2 -> He^+(3s+3d) B-110
He^{2+} + H_2 -> He^+(4p) B-110
He^{2+} + He -> He^+(2s) + He^+ B-112
He^{2+} + Li -> He^+(2p) + Li^+ B-118
He^{2+} + Li -> He^+(3p) + Li^+ B-118
He^{2+} + Li -> He^+(4p) + Li^+ B-118
He^{2+} + Li -> He^+(3s+3d) + Li^+ B-118
He^{2+} + Li -> He^+(4s+4d) + Li^+ B-118
He^{2+} + Li -> He^+(Lyman-α) + Li^+ B-120
He^{2+} + Li -> He^+(Lyman-β) + Li^+ B-120
He^{2+} + Li -> He^+(Lyman-γ) + Li^+ B-120
He^{2+} + Li -> He^+(Balmer-α) + Li^+ B-122
He^{2+} + Li -> He^+(Balmer-β) + Li^+ B-122
He^{2+} + Li -> He^+(Balmer-γ) + Li^+ B-122
He^{2+} + Li -> He^+(Balmer-σ) + Li^+ B-122

C. Excitation and Spectral Line Emission

H + H -> H(2s) C-2
-> H(2p) C-2
-> H(3s) C-4
-> H(3p) C-4
-> H(3d) C-4
-> H(Balmer-α) C-4

H + H_2 -> H(2s) [projectile] C-6
-> H(2p) [projectile] C-6
-> (2s) [target] C-6
-> (2p) [target] C-6
-> (Lyman-α) C-8
-> H(3s) C-10
-> H(3p+3d) C-10
-> H(Balmer-α) C-12
-> H(4s) C-14
-> H(4p+4d) C-14
-> H(Balmer-β) C-14

H(2s) + H_2 -> H(Lyman-α) C-16
-> H(3s) C-18
-> H(Balmer-α) C-18

H(2s) + He -> H(Lyman-α) C-20
-> H(3s) C-22
-> H(Balmer-α) C-22

H + He -> H(Lyman-α) C-24
-> H(2s) C-30
-> H(2p) C-30
-> H(3s) C-32
-> H(3p) C-32
-> H(3d) C-32
-> H(3p+3d) C-32
-> H(Balmer-α) C-34
-> H(Balmer-β) C-34
-> H(4s) C-36
-> H(4p+4d) C-36
-> He(3^1P) C-38

C. Excitation and Spectral Line Emission (Cont'd)

$H + He \rightarrow He(3^3P)$ C-38
$\rightarrow He(3^3D)$ C-38
$\rightarrow He(4^1S)$ C-40
$\rightarrow He(4^1P)$ C-40
$\rightarrow He(4^1D)$ C-40
$\rightarrow He(4^3S)$ C-42
$\rightarrow He(4^3P)$ C-42
$\rightarrow He(4^3D)$ C-42

$H^+ + H \rightarrow H(2s)$ C-44
$\rightarrow H(2p)$ C-44
$\rightarrow H(n=2)$ C-48
$\rightarrow H(n=3)$ C-48
$\rightarrow H(n=4)$ C-48

$H^+ + H_2 \rightarrow H^+ + H(2s) + H$ C-50
$\rightarrow H^+ + H(2p) + H$ C-50
$\rightarrow H^+ + H(3p) + H$ C-52
$\rightarrow H(Lyman\text{-}\alpha)$ C-54
$\rightarrow H(Balmer\text{-}\alpha)$ C-56
$\rightarrow H(Balmer\text{-}\beta)$ C-56
$\rightarrow H(Balmer\text{-}\gamma)$ C-56
$\rightarrow H_2(Lyman\text{-}Band)$ C-58
$\rightarrow H_2(Lyman\text{-}Band)$ C-58

$H^+ + He \rightarrow He(2^1S)$ C-60
$\rightarrow He(2^1P)$ C-60
$\rightarrow He(2^1S + 2^1P)$ C-60
$\rightarrow He(3^1S)$ C-66
$\rightarrow He(3^1P)$ C-66
$\rightarrow He(3^1D)$ C-66
$\rightarrow He(4^1S)$ C-68
$\rightarrow He(4^1P)$ C-68
$\rightarrow He(4^1D)$ C-68
$\rightarrow$ v.u.v.(20-140 nm) C-70

$He^+ + H \rightarrow H(n=2)$ C-72
$\rightarrow H(2s)$ C-74
$\rightarrow H(2p)$ C-74

$He^+ + H_2 \rightarrow H(Lyman\text{-}\alpha)$ C-78
$\rightarrow H(Lyman\text{-}\beta)$ C-78
$\rightarrow H(Balmer\text{-}\alpha)$ C-80
$\rightarrow H(Balmer\text{-}\beta)$ C-80
$\rightarrow H(2s)$ C-82
$\rightarrow H(2p)$ C-82
$\rightarrow H(3s)$ C-84
$\rightarrow H(3p)$ C-84
$\rightarrow He^+(2s)$ C-86

$He^+ + He \rightarrow He(2^1S)$ [target] C-88
$\rightarrow He(2^1P)$ [target] C-88
$\rightarrow He(2^3S)$ [target] C-90
$\rightarrow He(2^3P)$ [target] C-90
$\rightarrow He(3^1S)$ [target] C-92
$\rightarrow He(3^1P)$ [target] C-92
$\rightarrow He(3^1D)$ [target] C-92
$\rightarrow He(3^3S)$ [target] C-94
$\rightarrow He(3^3P)$ [target] C-94
$\rightarrow He(3^3D)$ [target] C-94

C. Excitation and Spectral Line Emission (Cont'd)

He^+ + He -> $He(4^1S)$ [target] C-96
-> $He(4^1P)$ [target] C-96
-> $He(4^1D)$ [target] C-96
-> $He(4^3S)$ [target] C-98
-> $He(4^3P)$ [target] C-98
-> $He(4^3D)$ [target] C-98
-> $He^+(n=2)$ [projectile] C-100
-> $He^+(n=3)$ [projectile] C-102

He + H_2 -> H(Balmer-α) C-104
-> H(Balmer-β) C-104

He + He -> $He(2^1S)$ C-106
-> $He(2^1P)$ C-106
-> $He(3^1P)$ C-108
-> $He(3^1D)$ C-108
-> $He(3^3P)$ C-108
-> $He(3^3D)$ C-108
-> $He(4^1S)$ C-110
-> $He(4^1P)$ C-110
-> $He(4^3S)$ C-112
-> $He(4^3P)$ C-112
-> $He(2s^2\ {}^1S)$ C-114
-> $He(2s2p^3P)$ C-114
-> $He(2p^2\ {}^1D)$ C-114
-> $He(2s2p^1P)$ C-114
-> $He^-(1s2s^2\ {}^2S)$ C-114

He^{2+} + H -> H(2s) C-116
-> H(2p) C-116

$He(2^3S)$ + $He(1^1S)$ -> $He(3^1P)$ + He C-118
-> $He(3^3P)$ + He C-120
-> $He(4^3S)$ + He C-122
-> $He(4^3D)$ + He C-122
-> He + $He(4^3S)$ C-124
-> He + $He(4^3D)$ C-124

H^+ + Li(2s) -> Li(2p) C-126
-> Li(670.8 nm) C-126

He^+ + Li(2s) -> Li(670.8 nm) C-128

He^{2+} + Li(2s) -> Li(2p) C-130
-> Li(670.8 nm) C-130

D. Ionization Collisions and Charge Production

H + H_2 -> Total Slow Positive Ion Production D-2
-> Total Slow Electron Production D-2
-> H + H_2^+ + e D-4

H^+ + H -> H^+ + H^+ + e D-6

H^+ + H^- -> H_2^+ + e D-12

H^+ + H_2 -> Total Slow Electron Production D-14
-> H^+ + H_2 + e D-16

D. Ionization Collisions and Charge Production (Cont'd)

$H^+ + H_2$ -> Total Production of H_2^+ D-16
-> Total Production of H^+ D-16

$H^+ + He$ -> Electrons and H^- D-18
-> $H^+ + He^+ + e$ D-24
-> $(H + He^+)$ or $(H^+ + He^+ + e)$ D-24
-> $H^+ + He^{2+} + 2e$ D-34
-> $(H^- + He^{2+})$ or $(H^+ + He^{2+} + 2e)$ D-34

$He(2^1S) + H$ -> $He + H^+ + e$ D-42
-> $HeH^+ + e$ D-42
-> $(He + H^+ + e)$ or $(HeH^+ + e)$ D-42

$He(2^3S) + H$ -> $He + H^+ + e$ D-44
-> $HeH^+ + e$ D-44
-> $(He + H^+ + e)$ or $(HeH^+ + e)$ D-44

$He(2^3S) + H_2$ -> Total Ionization D-46
-> Rearrangement Ionization D-46

$He(2^1S) + H_2$ -> Total Ionization D-48
-> Rearrangement Ionization D-48

$H^+ + He^+$ -> $(H^+ + He^{2+} + e)$ D-50
-> $(H^+ + He^{2+} + e)$ or $(H^o + He^{2+})$ D-50

$He^+ + He^+$ -> $(He^+ + He^{2+} + e)$ D-58
-> $(He + He^{2+})$ or $(He^+ + He^{2+} + e)$ D-58

$He + H_2$ -> Total Slow Positive Ion Production D-64
-> Total Slow Electron Production D-64
-> $He + H_2^+ + e$ D-66
-> Total H^+ Production D-66

$He + He$ -> Total Slow Positive Ion Production D-68
-> Total Slow Electron Production D-68
-> $He + He^+ + e$ D-70
-> $He + He^{2+} + 2e$ D-70

$He^+ + H$ -> $He^+ + H^+ + e$ D-72

$He^+ + H_2$ -> Total Slow Positive Ion Production D-76
-> Total Slow Electron Production D-76

$He^+ + H_2$ -> $(He + H_2^+)$ or $(He^+ + H_2^+ + e)$ D-78
-> Total H^+ Production D-78
-> $He^+ + H^+ + H^+ + 2e$ D-78

$He^+ + He$ -> Total Slow Positive Ion Production D-80
-> Total Slow Electron Production D-80
-> $He^+ + He^+ + e$ D-90
-> $He^+ + He^{2+} + 2e$ D-90

$He^{2+} + H$ -> $He^{2+} + H^+ + e$ D-96

$He^{2+} + H_2$ -> Total Slow Positive Ion Production D-100
-> Total Slow Electron Production D-100
-> Total H^+ Production D-102
-> H^+ Production by Ionization Collisions D-102
-> Total H_2^+ Production D-102
-> H_2^+ Production by Single-Ionization D-102

D. Ionization Collisions and Charge Production (Cont'd)

He^{2+} + He -> Total Slow Positive Ion Production D-104
-> Total Slow Electron Production D-104
-> He^{2+} + He^{+} + e D-114
-> He^{2+} + He^{2+} + 2e D-114

H^{+} + Li -> H^{+} + Li^{+} + e D-122

He^{+} + Li -> He^{+} + Li^{+} + e D-126

He^{2+} + Li -> He^{2+} + Li^{+} + e D-128

E. Electron Loss or Stripping Collisions

H + H -> H^{+} + H + e E-2
H(2s) + H(1s) -> H^{+} + H + e E-4
H + H_2 -> H^{+} + H_2 + e E-6
H(2s) + H_2 -> H^{+} + H_2 + e E-8
H + He -> H^{+} + He + e E-10
H(2s) + He -> H^{+} + He + e E-14

He + H -> He^{+} + H + e E-16
He + H_2 -> He^{+} + H_2 + e E-18
He + H_2 -> He^{2+} + H_2 + 2e E-18
He* + H_2 -> He or He^{+} E-20
He* + H_2 -> He^{+} + H_2 + e E-20
He* + H_2 -> He + H_2 . . . . . E-20
He + He -> He^{+} + He + e E-22
He + He -> He^{2+} + He + 2e E-22

He^{+} + H -> He^{2+} + H + e E-28
He^{+} + H_2 -> He^{2+} + H_2 + e E-30
He^{+} + He -> He^{2+} + He + e E-32

F. Electron Detachment Collisions

H^{-} + H -> H + H + e F-2
-> (H + H + e) or (H + H^{-}) F-2
-> H^{+} + H + 2e F-4
H^{-} + H^{+} -> H + H^{+} + 2e F-6
H^{-} + H_2 -> H + H_2 + e F-8
-> H^{+} + H_2 + 2e F-10
H^{-} + He -> H + He + e F-12
-> H^{+} + He + 2e F-14

He^{-} + H -> He + H + e F-16
He^{-} + H_2 -> He + H_2 + e F-18
-> He^{+} + H_2 + 2e F-18
He^{-} + He -> He + He + e F-20
-> He^{+} + He + 2e F-20

Li^{-} + H_2 -> Li + H_2 + e F-22
-> Li^{+} + H_2 + 2e F-22
Li^{-} + He -> Li + He + e F-24
-> Li^{+} + He + 2e F-24

G. Dissociative Collisions

$H + H_2 \rightarrow H^- + (H^+ + H^+) + e$ G-2
$\rightarrow H + (H^+ + H^+) + 2e$ G-2
$\rightarrow H^+ + (H^+ + H^+) + 3e$ G-2
$\rightarrow H + (H + H^+) + e$ G-4
$\rightarrow H^+ + (H + H^+) + 2e$ G-6
$\rightarrow H^- + (H^+ + H)$ G-8

$H^+ + H_2 \rightarrow H^+ + H^+ + H^+ + 2e$ G-10

$He^+ + H_2 \rightarrow He^+ + H^+ + H^+ + 2e$ G-12

$He^{2+} + H_2 \rightarrow He^{2+} + (H^+ + H) + e$ (via $1s\sigma_g$) G-14
$\rightarrow He^{2+} + (H^+ + H) + e$ (via $2p\sigma_u$) G-16
$\rightarrow He^+ + (H^+ + H)$ G-18
$\rightarrow He^+ + (H^+ + H^+) + e$ G-20
$\rightarrow He^{2+} + (H^+ + H^+) + 2e$ G-22
$\rightarrow He + (H^+ + H^+)$ G-24

$H_2^+ + H_2 \rightarrow H^+_{Proj.}$ G-26

$H_2^+ + H \rightarrow H^+_{Proj.}$ G-28

$H_2^+ + H_2 \rightarrow H_{Proj.}$ G-30
$\rightarrow (H^+ + H) + H_2$ G-32

$H_2^+ + He \rightarrow$ Total H^+ G-34
$\rightarrow (H^+ + H) + He$ G-36

$H_2 + H_2 \rightarrow H^+$(Fast,Total) G-38
$\rightarrow$ H(Fast,Total) G-40
$\rightarrow$ Total Destruction of Fast H_2 Proj G-42

$H_3^+ + H_2 \rightarrow$ Total Destruction of H_3^+ G-44

$H_2^+ + H_2 \rightarrow$ Total Destruction of H_2^+ G-46

$H_3^+ + H_2 \rightarrow H^+$(Total,Projectile) G-48
$\rightarrow H_2^+$(Total,Projectile) G-50
$\rightarrow$ H(Total,Projectile) G-52
$\rightarrow H_2$(Total,Projectile) G-54

$H_3^+ + He \rightarrow H^+$(Total) G-56
$\rightarrow H_2^+$ G-58

$HeH^+ + H_2 \rightarrow$ Total HeH^+ Dissociation Prod. G-60

$HeH^+ + He \rightarrow$ Total HeH^+ Dissociation Prod. G-62

H. Particle Interchange Reactions

$H^+ + D_2 \rightarrow D^+ + HD$ H-2
$\rightarrow HD^+ + D$ H-2

$HD^+ + D_2 \rightarrow HD_2^+ + D$ H-4
$\rightarrow D_3^+ + H$ H-4

H. Particle Interchange Reactions (Cont'd)

$HD^+(\upsilon=0) + He \rightarrow HeH^+ + D$ H-6
$HD^+(\upsilon=1) + He \rightarrow HeH^+ + D$ H-6
$HD^+(\upsilon=2) + He \rightarrow HeH^+ + D$ H-6
$HD^+(\upsilon=3) + He \rightarrow HeH^+ + D$ H-6
$HD^+(\upsilon=4) + He \rightarrow HeH^+ + D$ H-6

$H_2^+ + He \rightarrow HeH^+ + H$ H-8

$H_2^+ + H_2 \rightarrow H_3^+ + H$ H-10

$D_2^+ + D_2 \rightarrow D_3^+ + D$ H-12

$D_2^+(\upsilon=0) + H_2 \rightarrow HD_2^+ + H$ H-14
$D_2^+(\upsilon=1) + H_2 \rightarrow HD_2^+ + H$ H-14
$D_2^+(\upsilon=2) + H_2 \rightarrow HD_2^+ + H$ H-14
$D_2^+(\upsilon=3) + H_2 \rightarrow HD_2^+ + H$ H-14

$H_2^+(\upsilon=0) + D_2 \rightarrow HD_2^+ + D$ H-16
$H_2^+(\upsilon=1) + D_2 \rightarrow HD_2^+ + D$ H-16
$H_2^+(\upsilon=2) + D_2 \rightarrow HD_2^+ + D$ H-16
$H_2^+(\upsilon=3) + D_2 \rightarrow HD_2^+ + D$ H-16
$H_2^+(\upsilon=4) + D_2 \rightarrow HD_2^+ + D$ H-16

References R-1

Appendix 1: Calculation of Cross Sections and Rate Coefficients from Chebyshev Fitting Parameters 1-1

Appendix 2: ALADDIN Database System and Data Files 2-1

Appendix 3: Atomic Physics (Handy Conversion Units and formulas) 3-1

FOREWORD

The Controlled Fusion Atomic Data Center (CFADC) was founded informally at Oak Ridge National Laboratory by Clarence F. Barnett in 1959, and was officially established six years later. The continuing mission of the CFADC is to identify, compile, evaluate and recommend data on atomic and molecular collision processes which are important in fusion energy research. "Barney" dedicated much of his professional career to the promotion of effective scientific communication between the atomic and fusion research communities, and was still actively engaged in this mission at the time of his death in 1989. This "Redbook" volume represents a major part of his continuing professional efforts during the years following his formal retirement from ORNL in 1985. Barnett's insights, guidance and dedication to this mission will be sorely missed by the CFADC and by the fusion research community.

Ronald A. Phaneuf
Hamilton T. Hunter
M. Imogene Kirkpatrick
David H. Crandall (DOE)

This photo is from his "official" retirement in 1985. His spirit will continue to encourage us in the future.



Series Preface

The primary objective of the Controlled Fusion Atomic Data Center at Oak Ridge National Laboratory is to publish and distribute handbooks containing numerical and graphical cross-section and other physical data relevant to fusion energy research. In 1977, a two-volume compilation was published as ORNL reports ORNL-5206 and ORNL-5207. Since that time, a large volume of pertinent data has become available, necessitating an update and expansion of the previous compilation. The specific volumes in this series, entitled "Atomic Data for Fusion," are listed below.

Vol. 1, "Collisions of H, H_2, He, and Li Atoms and Ions with Atoms and Molecules," C. F. Barnett, ORNL, ORNL-6086 (June 1990).

Vol. 2, "Collisions of Electrons with Atoms and Molecules," J. W. Gallagher, National Institute of Standards and Technology; and D. C. Gregory, ORNL (in preparation).

Vol. 3, "Particle Interactions with Surfaces," E. W. Thomas, Georgia Institute of Technology, ORNL-6088 (January 1985).

Vol. 4, "Spectroscopic Data for Iron," W. L. Wiese, National Bureau of Standards, ORNL-6089 (March 1985).

Vol. 5, "Collisions of Carbon and Oxygen Ions with Electrons, H, H_2, and He," R. A. Phaneuf, ORNL; R. K. Janev, Institute of Physics, Yugoslavia; and M. S. Pindzola, Auburn University, ORNL-6090 (January 1987).

Vol. 6, "Spectroscopic Data for Titanium, Chromium, and Nickel," W. L. Wiese and A. Musgrove, National Institute for Standards and Technology, ORNL-6551 (September 1989).

C. F. Barnett
H. T. Hunter
M. I. Kirkpatrick
R. A. Phaneuf

June 1990

ABSTRACT

This report provides a handbook of recommended cross-section and rate-coefficient data for inelastic collisions between hydrogen, helium and lithium atoms, molecules and ions, and encompasses more than 400 different reactions of primary interest in fusion research. Published experimental and theoretical data have been collected and evaluated, and the recommended data are presented in tabular, graphical and parametrized form. Processes include excitation and spectral line emission, charge exchange, ionization, stripping, dissociation and particle interchange reactions. The range of collision energies is appropriate to applications in fusion-energy research.

INTRODUCTION

This volume contains recommended cross-section and rate-coefficient data for inelastic collisions between hydrogen, helium and lithium atoms, molecules and ions, and encompasses 383 different reactions of primary interest in fusion research. Published experimental and theoretical data have been collected and evaluated, and the recommended data are presented in tabular, graphical and parametrized form. Inelastic processes include excitation and spectral-line emission, charge exchange, ionization, stripping, dissociation and particle interchange reactions. Emphasis has been given to the range of collision energies appropriate to diagnostic and modelling applications in fusion-energy research.

The data which form the basis for this volume of recommended data are those which were available as of September 1988. The bibliographic files of the ORNL Controlled Fusion Atomic Data Center provided the lists of references for this volume. In the vast majority of cases, the numerical data were obtained from the original publications, either from tables where available, or from figures using a precision graphical digitizer system. The rms uncertainty associated with the digitization system was determined to be less than 2%.

The recommended data in this compilation are based almost exclusively on experimental data. Theoretical data have been considered when it was desirable to extend the measurements to higher or lower collision energies, and also for a few important reactions where experimental data were unavailable. Specific references to the literature and notes are presented for each reaction, along with an estimated uncertainty in the recommended cross section. The latter was determined from uncertainties quoted in the original data sources, and also from the consistency of the data when more than one source was available. To determine the recommended cross sections in the latter case, relative weightings were made on the basis of the reliability of the different experimental methods used. In some cases, possible effects on the measurements due to other competing processes were also considered in determining the recommended cross section and its estimated uncertainty. For some reactions, it was not possible to estimate an uncertainty. In all cases, the estimated accuracies of the recommended data are believed to have been conservatively estimated, and are intended to represent upper limits.

Reaction rate coefficients have been calculated from the recommended cross-section data where appropriate for fusion applications, and where the energy range of the available data permits. These are based on Maxwellian velocity distributions for each reactant, where each may be characterized by a different temperature, and also for the case of a monoenergetic

(neutral) beam interacting with a Maxwellian distribution. The temperature ranges for the calculated rate coefficients are constrained by the energy range over which cross-section data are available. The tabulated values for the rate coefficients have been flagged in those temperature ranges where the limited energy range of the data appreciably influences their accuracy.

For convenience, least-squares Chebyshev polynomial fits to the recommended cross-sections and rate coefficients are presented for each reaction. These polynomial fits are indicated graphically, and a statement is made concerning the accuracy of each fit to the recommended data. Practical details on the least-squares fits and sample computer programs for rapidly generating the recommended cross sections and rate coefficients from the fitting coefficients are presented in Appendix 1.

For further ease of retrieval and application of the recommended cross-section data, a PC-DOS formatted diskette has been prepared and is available for distribution. Included on this diskette are data files with the Chebyshev fitting coefficients for each reaction encoded in *ALADDIN* format,[1,2] along with a copy of the *ALADDIN* program (written in FORTRAN-77). The *ALADDIN* program and data files may also be conveniently transmitted by electronic mail. Further details about *ALADDIN* are presented in Appendix 2, along with samples of the data entries from each chapter. Information about how to obtain the program and data files for this and other compilations may also be found in Appendix 2.

---

1. R. A. Hulse, "The ALADDIN Atomic Physics Database System," pp. 63-72 in Atomic Processes in Plasmas, AIP Conference Proceedings 206, ed. Y. K. Kim and R. C. Elton, AIP, New York, 1990.

2. ALADDIN Manual, A System for Storage, Exchange and Management of Atomic and Molecular Data for Fusion, IAEA-NDS-AM-17, Atomic and Molecular Data Unit, Nuclear Data Section, International Atomic Energy Agency, Vienna, June 1989.

A. Total Electron Capture Cross Sections

(* denotes rate coefficient data also)

	Page
Electron Capture by Neutral H	**A-2**
$H + H \rightarrow H^- + H^+$	A-2*
$H + H_2 \rightarrow H^- + H_2^+$	A-8
$H + He \rightarrow H^- + He^+$	A-10*
$H + Li \rightarrow H^- + Li^+$	A-16*
$H + H^- \rightarrow H^- + H$	A-20
Electron Capture by H^+	**A-22**
$H^+ + H \rightarrow H + H^+$	A-22*
$H^+ + H_2 \rightarrow H$	A-28
$H^+ + H_2 \rightarrow H^-$	A-30
$H^+ + He \rightarrow H + He^+$	A-32*
$H^+ + He \rightarrow H^- + He^{2+}$	A-38
$H^+ + Li \rightarrow H + Li^+$	A-40*
$H^+ + H^- \rightarrow H^- + H^+$	A-44
$H^+ + H^- \rightarrow 2H$	A-46
$H^+ + He^+ \rightarrow \sigma(He^{2+}$ Total)	A-48*
$H^+ + He^+ \rightarrow H + He^{2+}$	A-54*
Electron Capture by H_2^+	**A-58**
$H_2^+ + H \rightarrow H_2 + H^+$	A-58
$H_2^+ + H_2 \rightarrow H_2$	A-58
$H_2^+ + He \rightarrow H_2 + He^+$	A-58
$H_2^+(\nu) + H_2 \rightarrow H_2$	A-60
Electron Capture by He^+	**A-62**
$He^+ + H \rightarrow He + H^+$	A-62*
$He^+ + H^- \rightarrow He + H$	A-86
$He^+ + H_2 \rightarrow He$	A-68
$He^+ + He \rightarrow He + He^+$	A-70*
$He^+ + Li \rightarrow He + Li^+$	A-76*
$He^+ + He^+ \rightarrow He + He^{2+}$	A-86*
Electron Capture by He^{2+}	**A-84**
$He^{2+} + He^+ \rightarrow He^+ + He^{2+}$(Partial & Total)	A-84
$He^{2+} + H^- \rightarrow He^+ + H$	A-86
$He^{2+} + H \rightarrow He^+ + H^+$	A-88*
$He^{2+} + H_2 \rightarrow He^+$	A-94
$He^{2+} + H_2 \rightarrow He$	A-96
$He^{2+} + He \rightarrow He^+ + He^+$	A-98*
$He^{2+} + He \rightarrow He + He^{2+}$	A-104*
$He^{2+} + Li \rightarrow He^+ + Li^+$	A-110*
$He^{2+} + Li \rightarrow He + Li^{2+}$	A-114*

Electron Capture Cross Sections for
$H + H \rightarrow H^- + H^+$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.0E+03	6.21E+07	1.56E-18
4.0E+03	8.79E+07	3.64E-18
7.0E+03	1.16E+08	7.63E-18
1.0E+04	1.39E+08	1.18E-17
1.6E+04	1.76E+08	1.63E-17
2.0E+04	1.96E+08	1.55E-17
3.1E+04	2.45E+08	9.98E-18
4.0E+04	2.78E+08	5.58E-18
7.0E+04	3.68E+08	1.48E-18
7.3E+04	3.75E+08	1.37E-18

References: 33, 34, 396

Accuracy: $E < 8x10^3$ eV/amu - 50%; $E > 8x10^3$ eV/amu - 20%

Note: For energies less than $8x10^3$ eV/amu large discrepancies exist between the data.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 2.0E+03 eV/amu, E_{max} = 7.3E+04 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-79.9637	.00368622	-1.08628	-.302134	.0745111	.148356	-.0426883	.0535533	-.0289283

The fit represents the above cross sections with an rms deviation of 1.0%.
The maximum deviation is 2.0% at 2.0E+04 eV/amu.
See appendix for Chebyshev fit details.

$$H + H \rightarrow H^- + H^+$$



Electron Capture Rate Coefficients for
$H + H \rightarrow H^- + H^+$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H Temp. (eV)	Equal Temp.	H Temp. (eV) 1000.	2000.	5000.	10000.	12000.	15000.	20000.
1.0E+02	1.96E-14	7.56E-11	2.82E-10	1.01E-09	1.62E-09	1.70E-09	1.72E-09	1.65E-09
2.0E+02	2.57E-12	9.22E-11	3.06E-10	1.03E-09	1.63E-09	1.70E-09	1.72E-09	1.65E-09
4.0E+02	3.36E-11	1.29E-10	3.56E-10	1.07E-09	1.64E-09	1.71E-09	1.72E-09	1.64E-09
7.0E+02	1.29E-10	1.90E-10	4.32E-10	1.13E-09	1.65E-09	1.71E-09	1.72E-09	1.63E-09
1.0E+03	2.58E-10	2.58E-10	5.10E-10	1.18E-09	1.66E-09	1.72E-09	1.72E-09	1.63E-09
2.0E+03	7.64E-10	5.10E-10	7.64E-10	1.34E-09	1.70E-09	1.72E-09	1.71E-09	1.60E-09
4.0E+03	1.46E-09	9.91E-10	1.18E-09	1.55E-09	1.72E-09	1.72E-09	1.67E-09	1.55E-09
7.0E+03	1.72E-09	1.46E-09	1.55E-09	1.70E-09	1.71E-09	1.67E-09	1.60E-09	1.47E-09
1.0E+04	1.65E-09	1.66E-09	1.70E-09	1.72E-09	1.65E-09	1.60E-09	1.52E-09	1.38E-09
2.0E+04	1.14E-09*	1.63E-09	1.60E-09	1.52E-09	1.38E-09	1.33E-09	1.26E-09*	1.14E-09*

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Note: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+02 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

H Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
1000.	-4.34136E+01	1.77058E+00	-2.19086E-02	-2.54539E-01	-5.12704E-02	1.82644E-02	9.12815E-03
2000.	-4.22141E+01	1.02952E+00	3.42057E-02	-1.66830E-01	-5.60021E-02	5.86271E-03	8.15346E-03
5000.	-4.09368E+01	2.82198E-01	-5.96305E-03	-7.32604E-02	-3.54191E-02	-5.05727E-03	2.39134E-03
10000.	-4.04860E+01	-3.08846E-02	-5.68396E-02	-4.34280E-02	-1.90470E-02	-4.61028E-03	-1.80695E-05
12000.	-4.04639E+01	-7.77066E-02	-6.68905E-02	-3.99404E-02	-1.61322E-02	-3.99466E-03	-2.04042E-04
15000.	-4.04949E+01	-1.18325E-01	-7.60484E-02	-3.69771E-02	-1.33535E-02	-3.24675E-03	-2.69603E-04
20000.	-4.06294E+01	-1.47825E-01	-8.23614E-02	-3.42640E-02	-1.08621E-02	-2.44826E-03	-2.60694E-04
Equal Temp.	-4.70326E+01	4.89621E+00	-2.27782E+00	4.67042E-01	-2.66936E-01	1.22082E-01	-1.65425E-02

See appendix for Chebyshev fit details.

$$H + H \rightarrow H^- + H^+$$

Maxwellian – Maxwellian



Electron Capture Rate Coefficients for
$H + H \rightarrow H^- + H^+$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H Temp. (eV)	H Energy (eV/amu) 10000.	20000.	40000.	45000.	50000.	60000.	70000.
1.0E+00	1.63E-09**	3.03E-09**	1.55E-09**	1.24E-09**	1.02E-09**	7.26E-10**	5.45E-10**
2.0E+00	1.63E-09*	3.03E-09*	1.55E-09**	1.24E-09**	1.02E-09**	7.26E-10**	5.45E-10**
4.0E+00	1.63E-09*	3.02E-09*	1.55E-09**	1.24E-09**	1.02E-09**	7.26E-10**	5.45E-10**
7.0E+00	1.63E-09*	3.02E-09*	1.55E-09**	1.24E-09**	1.02E-09**	7.26E-10**	5.45E-10**
1.0E+01	1.63E-09*	3.01E-09*	1.55E-09**	1.24E-09**	1.02E-09**	7.26E-10**	5.43E-10**
2.0E+01	1.63E-09*	3.00E-09*	1.54E-09**	1.24E-09**	1.02E-09**	7.26E-10**	5.27E-10**
4.0E+01	1.63E-09*	2.98E-09*	1.54E-09**	1.25E-09**	1.02E-09**	7.27E-10**	4.94E-10**
7.0E+01	1.63E-09*	2.96E-09*	1.54E-09**	1.25E-09**	1.02E-09**	7.28E-10**	4.62E-10**
1.0E+02	1.64E-09*	2.94E-09*	1.54E-09**	1.25E-09**	1.03E-09**	7.29E-10**	4.42E-10**
2.0E+02	1.66E-09*	2.89E-09*	1.54E-09**	1.25E-09**	1.03E-09**	7.29E-10**	4.10E-10**
4.0E+02	1.71E-09*	2.81E-09*	1.55E-09*	1.26E-09**	1.04E-09**	7.19E-10**	3.91E-10**
7.0E+02	1.76E-09*	2.71E-09*	1.55E-09*	1.27E-09**	1.05E-09**	7.03E-10**	3.86E-10**
1.0E+03	1.80E-09*	2.62E-09*	1.54E-09*	1.28E-09*	1.05E-09**	6.93E-10**	3.88E-10**
2.0E+03	1.88E-09*	2.40E-09*	1.52E-09*	1.27E-09*	1.05E-09**	6.86E-10**	4.08E-10**
4.0E+03	1.90E-09	2.12E-09*	1.42E-09*	1.21E-09*	1.02E-09**	6.97E-10**	4.51E-10**
7.0E+03	1.86E-09*	1.85E-09*	1.28E-09*	1.12E-09*	9.66E-10*	7.01E-10**	4.92E-10**
1.0E+04	1.78E-09*	1.67E-09*	1.16E-09*	1.03E-09*	9.10E-10*	6.90E-10**	5.11E-10**
2.0E+04	1.48E-09*	1.29E-09*	9.21E-10*	8.39E-10*	7.61E-10**	6.22E-10**	5.03E-10**

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Note: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

H Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-4.04105E+01	2.91340E-02	-2.63688E-02	-6.02938E-02	-4.80367E-02	-1.89721E-02	-1.81756E-03
20000.	-3.96573E+01	-3.51288E-01	-1.95694E-01	-7.65178E-02	-1.76176E-02	6.76024E-05	8.20202E-04
40000.	-4.07630E+01	-1.72032E-01	-1.30927E-01	-8.12602E-02	-3.70555E-02	-8.68416E-03	3.26863E-03
45000.	-4.11299E+01	-1.16178E-01	-1.03253E-01	-7.22628E-02	-3.62733E-02	-1.02839E-02	1.83555E-03
50000.	-4.14810E+01	-7.82977E-02	-7.66264E-02	-5.87240E-02	-3.17702E-02	-1.03177E-02	-4.83902E-04
60000.	-4.21440E+01	-5.01687E-02	-2.65441E-02	-1.11314E-02	-8.49686E-03	-1.30422E-02	-1.43358E-02
70000.	-4.28827E+01	-7.90160E-02	1.09848E-01	8.45395E-02	-3.13642E-02	-4.50109E-02	-6.86268E-03

See appendix for Chebyshev fit details.

$$H + H \rightarrow H^- + H^+$$

Beam – Maxwellian



Electron Capture Cross Sections for
$H + H_2 \rightarrow H^-$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
4.0E+01	8.79E+06	1.05E-18
7.0E+01	1.16E+07	1.78E-18
1.0E+02	1.39E+07	2.52E-18
1.4E+02	1.64E+07	3.08E-18
2.0E+02	1.96E+07	2.74E-18
4.0E+02	2.78E+07	1.70E-18
7.0E+02	3.68E+07	1.24E-18
1.0E+03	4.39E+07	1.31E-18
2.0E+03	6.21E+07	5.28E-18
4.0E+03	8.79E+07	1.46E-17
7.0E+03	1.16E+08	2.18E-17
1.0E+04	1.39E+08	2.32E-17
2.0E+04	1.96E+08	1.91E-17
4.0E+04	2.78E+08	9.93E-18
7.0E+04	3.68E+08	4.07E-18
1.0E+05	4.39E+08	1.68E-18
2.0E+05	6.21E+08	1.72E-19
4.0E+05	8.78E+08	6.12E-21
4.6E+05	9.42E+08	2.97E-21

References: 26, 27, 28, 29, 30, 31, 32, 33, 35, 396

Accuracy: 25%

Note: The quoted results are believed to be accurate to within 25%, although the data presented in some of the references may deviate by 50-60%.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 4.0E+01 eV/amu, E_{max} = 4.6E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-82.8333	-1.65457	-2.60408	-1.53870	-.368649	.413620	-.165702	-.127116	.256041

The fit represents the above cross sections with an rms deviation of 9.5%.
The maximum deviation is 26.2% at 1.0E+03 eV/amu.
See appendix for Chebyshev fit details.

$H^- + H_2 \rightarrow H^-$



Electron Capture Cross Sections for
$H + He \rightarrow H^- + He^+$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
7.7E+02	3.85E+07	1.47E-19
1.0E+03	4.39E+07	2.15E-19
2.0E+03	6.21E+07	5.80E-19
3.0E+03	7.61E+07	1.04E-18
4.0E+03	8.79E+07	1.61E-18
5.0E+03	9.82E+07	2.23E-18
6.0E+03	1.08E+08	2.88E-18
7.0E+03	1.16E+08	3.46E-18
9.0E+03	1.32E+08	4.34E-18
1.0E+04	1.39E+08	4.83E-18
2.0E+04	1.96E+08	6.51E-18
3.0E+04	2.41E+08	6.06E-18
4.0E+04	2.78E+08	5.28E-18
7.0E+04	3.68E+08	2.92E-18
1.0E+05	4.39E+08	1.46E-18
2.0E+05	6.21E+08	2.33E-19
4.0E+05	8.78E+08	2.47E-20
7.0E+05	1.16E+09	1.87E-21
7.6E+05	1.21E+09	1.16E-21

References: 26, 27, 29, 35, 268, 269, 270

Accuracy: 30%

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 7.7E+02 eV/amu, E_{max} = 7.6E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-85.5215	-1.87685	-3.06302	-.492323	.117220	-.0269941	-.0479364	-.0315786	-.0439288

The fit represents the above cross sections with an rms deviation of 1.8%.
The maximum deviation is 2.9% at 7.0E+05 eV/amu.
See appendix for Chebyshev fit details.

$$H + He \rightarrow H^- + He^+$$



Electron Capture Rate Coefficients for
$H + He \rightarrow H^- + He^+$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H Temp. (eV)	Equal Temp.	He Temp. (eV) 1.	100.	500.	1000.	5000.	10000.	20000.
7.0E+01	3.79E-15	4.61E-16	7.47E-15	4.58E-13	2.43E-12	5.54E-11	1.74E-10	4.26E-10
1.0E+02	5.02E-14	1.13E-14	5.02E-14	7.92E-13	3.11E-12	5.78E-11	1.77E-10	4.28E-10
2.0E+02	1.14E-12	5.09E-13	7.92E-13	2.54E-12	5.86E-12	6.60E-11	1.87E-10	4.37E-10
4.0E+02	7.50E-12	4.40E-12	5.11E-12	8.38E-12	1.34E-11	8.33E-11	2.08E-10	4.55E-10
7.0E+02	2.48E-11	1.57E-11	1.69E-11	2.20E-11	2.93E-11	1.11E-10	2.40E-10	4.82E-10
1.0E+03	5.00E-11	3.24E-11	3.40E-11	4.08E-11	5.00E-11	1.41E-10	2.71E-10	5.08E-10
2.0E+03	1.67E-10	1.16E-10	1.19E-10	1.29E-10	1.41E-10	2.45E-10	3.72E-10	5.86E-10
4.0E+03	4.19E-10	3.22E-10	3.25E-10	3.35E-10	3.47E-10	4.42E-10	5.48E-10	7.17E-10
7.0E+03	7.02E-10	5.86E-10	5.88E-10	5.96E-10	6.05E-10	6.72E-10	7.44E-10	8.55E-10
1.0E+04	8.73E-10	7.70E-10	7.71E-10	7.76E-10	7.82E-10	8.26E-10	8.73E-10	9.43E-10
2.0E+04	1.04E-09	1.02E-09	1.02E-09	1.02E-09	1.02E-09	1.02E-09	1.03E-09	1.04E-09

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 7.0E+01 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
1.	-5.10941E+01	6.54972E+00	-2.07764E+00	6.21336E-01	-3.71355E-01	1.34135E-01	-1.51404E-02
100.	-5.00589E+01	5.62889E+00	-1.42974E+00	2.34011E-01	-1.74762E-01	5.17776E-02	1.40144E-02
500.	-4.81708E+01	4.05303E+00	-4.72178E-01	-2.11436E-01	-1.16459E-02	1.30366E-02	9.51225E-03
1000.	-4.71671E+01	3.28859E+00	-1.32390E-01	-2.90109E-01	-1.07340E-02	2.14696E-02	2.38292E-03
5000.	-4.45966E+01	1.61569E+00	1.93393E-01	-1.64393E-01	-5.91438E-02	7.61287E-03	6.54831E-03
10000.	-4.34391E+01	9.76929E-01	1.85422E-01	-8.18919E-02	-5.25533E-02	-5.26730E-03	3.99780E-03
20000.	-4.24393E+01	4.84530E-01	1.19557E-01	-3.07242E-02	-3.26702E-02	-9.20869E-03	1.64244E-04
Equal Temp.	-4.97998E+01	5.73848E+00	-1.74839E+00	4.04712E-01	-2.79431E-01	1.17900E-01	-2.07435E-02

See appendix for Chebyshev fit details.

$$H + He \rightarrow H^- + He^+$$

Maxwellian — Maxwellian



Electron Capture Rate Coefficients for
He + H -> H^- + He^+

Beam - Maxwellian Rate Coefficients (cm^3/s)

H Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	6.69E-10	1.28E-09	1.46E-09	1.07E-09	6.41E-10	1.45E-10	8.66E-12**
2.0E+00	6.68E-10	1.27E-09	1.46E-09	1.07E-09	6.40E-10	1.45E-10	8.67E-12**
4.0E+00	6.67E-10	1.27E-09	1.46E-09	1.07E-09	6.40E-10	1.45E-10	8.67E-12**
7.0E+00	6.66E-10	1.27E-09	1.46E-09	1.07E-09	6.39E-10	1.44E-10	8.67E-12**
1.0E+01	6.65E-10	1.27E-09	1.46E-09	1.07E-09	6.39E-10	1.44E-10	8.67E-12**
2.0E+01	6.64E-10	1.27E-09	1.46E-09	1.06E-09	6.38E-10	1.44E-10	8.67E-12**
4.0E+01	6.63E-10	1.26E-09	1.45E-09	1.06E-09	6.37E-10	1.44E-10	8.68E-12**
7.0E+01	6.63E-10	1.26E-09	1.45E-09	1.06E-09	6.36E-10	1.44E-10	8.68E-12**
1.0E+02	6.64E-10	1.25E-09	1.44E-09	1.05E-09	6.35E-10	1.44E-10	8.69E-12**
2.0E+02	6.69E-10	1.24E-09	1.43E-09	1.05E-09	6.33E-10	1.44E-10	8.72E-12**
4.0E+02	6.82E-10	1.24E-09	1.42E-09	1.04E-09	6.31E-10	1.44E-10	8.77E-12*
7.0E+02	7.04E-10	1.23E-09	1.40E-09	1.03E-09	6.30E-10	1.44E-10	8.85E-12*
1.0E+03	7.25E-10	1.22E-09	1.39E-09	1.02E-09	6.30E-10	1.45E-10	8.93E-12*
2.0E+03	7.87E-10	1.21E-09	1.34E-09	9.96E-10	6.29E-10	1.47E-10	9.20E-12*
4.0E+03	8.79E-10	1.18E-09	1.27E-09	9.60E-10	6.26E-10	1.52E-10	9.70E-12*
7.0E+03	9.65E-10	1.15E-09	1.18E-09	9.11E-10	6.18E-10	1.60E-10	1.04E-11*
1.0E+04	1.01E-09	1.12E-09	1.11E-09	8.67E-10	6.07E-10	1.68E-10	1.10E-11*
2.0E+04	1.05E-09	1.04E-09	9.50E-10	7.54E-10	5.63E-10	1.90E-10	1.34E-11*

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-4.19935E+01	2.19891E-01	1.24242E-01	2.58739E-02	-1.96928E-02	-2.10662E-02	-9.41303E-03
20000.	-4.10635E+01	-7.94718E-02	-3.94031E-02	-1.87660E-02	-9.93076E-03	-4.65478E-03	-1.66689E-03
40000.	-4.08823E+01	-1.64742E-01	-9.93840E-02	-4.75948E-02	-1.79398E-02	-4.36181E-03	4.08769E-04
70000.	-4.14695E+01	-1.30558E-01	-7.61140E-02	-3.80809E-02	-1.75717E-02	-6.97521E-03	-1.71873E-03
100000.	-4.23878E+01	-4.00395E-02	-2.39094E-02	-1.67316E-02	-1.22134E-02	-7.11472E-03	-2.89733E-03
200000.	-4.52157E+01	9.17644E-02	7.16076E-02	4.20628E-02	1.78905E-02	4.62301E-03	-6.38872E-04
500000.	-5.07701E+01	1.52640E-01	1.03844E-01	5.55224E-02	2.32810E-02	7.67676E-03	2.39728E-03

See appendix for Chebyshev fit details.

$He + H \rightarrow H^- + He^+$

Beam – Maxwellian



Electron Capture Cross Sections for
$H + Li \rightarrow H^- + Li^+$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
3.0E+05	7.61E+08	1.71E-18
4.0E+05	8.78E+08	8.97E-19
5.0E+05	9.82E+08	6.99E-19
6.0E+05	1.08E+09	6.13E-19
7.0E+05	1.16E+09	5.48E-19
8.0E+05	1.24E+09	4.93E-19
9.0E+05	1.32E+09	4.49E-19
1.0E+06	1.39E+09	4.15E-19

References: 268

Accuracy: 30%

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 3.0E+05 eV/amu, E_{max} = 1.0E+06 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-83.5167	-.651028	.130590	-.0639887	.0130205	6.98064E-03	-3.26857E-03	5.16789E-05	

The fit represents the above cross sections with an rms deviation of 0.0%.
The maximum deviation is 0.0%.
See appendix for Chebyshev fit details.

$$H + Li \rightarrow H^- + Li^+$$



Electron Capture Rate Coefficients for
$Li + H \rightarrow H^- + Li^+$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H Temp. (eV)	Li Energy (eV/amu) 300000.	350000.	400000.	450000.	500000.	550000.	600000.
1.0E+00	6.49E-10**	9.94E-10**	7.89E-10**	7.33E-10**	6.87E-10**	6.72E-10**	6.59E-10**
2.0E+00	6.48E-10**	9.94E-10**	7.89E-10**	7.33E-10**	6.87E-10**	6.72E-10**	6.59E-10**
4.0E+00	6.47E-10**	9.94E-10**	7.89E-10**	7.33E-10**	6.87E-10**	6.72E-10**	6.59E-10**
7.0E+00	6.46E-10**	9.94E-10**	7.90E-10**	7.33E-10**	6.87E-10**	6.72E-10**	6.59E-10**
1.0E+01	6.45E-10**	9.94E-10**	7.90E-10**	7.33E-10**	6.87E-10**	6.72E-10**	6.59E-10**
2.0E+01	6.43E-10**	9.94E-10**	7.91E-10**	7.33E-10**	6.87E-10**	6.72E-10**	6.59E-10**
4.0E+01	6.40E-10**	9.95E-10**	7.93E-10**	7.33E-10**	6.88E-10**	6.72E-10**	6.59E-10**
7.0E+01	6.37E-10**	9.95E-10**	7.95E-10**	7.33E-10**	6.88E-10**	6.72E-10**	6.59E-10**
1.0E+02	6.34E-10**	9.95E-10**	7.96E-10**	7.33E-10**	6.89E-10**	6.72E-10**	6.59E-10**
2.0E+02	6.28E-10**	9.95E-10**	7.99E-10**	7.33E-10**	6.89E-10**	6.72E-10**	6.59E-10**
4.0E+02	6.19E-10**	9.96E-10**	8.05E-10**	7.33E-10**	6.91E-10**	6.72E-10**	6.59E-10**
7.0E+02	6.09E-10**	9.87E-10**	8.10E-10**	7.33E-10**	6.92E-10**	6.72E-10**	6.59E-10**
1.0E+03	6.02E-10**	9.70E-10**	8.15E-10**	7.34E-10**	6.93E-10**	6.72E-10**	6.59E-10**
2.0E+03	5.84E-10**	9.04E-10**	8.24E-10**	7.39E-10**	6.96E-10**	6.74E-10**	6.59E-10**
4.0E+03	5.62E-10**	8.10E-10**	8.13E-10**	7.49E-10**	7.02E-10**	6.76E-10**	6.60E-10**
7.0E+03	5.43E-10**	7.33E-10**	7.78E-10**	7.49E-10**	7.09E-10**	6.81E-10**	6.61E-10**
1.0E+04	5.31E-10**	6.89E-10**	7.46E-10**	7.39E-10**	7.11E-10**	6.84E-10**	6.63E-10**
2.0E+04	5.11E-10**	6.17E-10**	6.75E-10**	6.96E-10**	6.92E-10**	6.77E-10**	6.55E-10**

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Note: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

Li Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
300000.	-4.24630E+01	-1.12797E-01	-4.73823E-02	-9.52319E-03	2.18237E-03	2.88098E-03	1.39108E-03
350000.	-4.16874E+01	-2.00135E-01	-1.31634E-01	-5.41437E-02	-1.93565E-03	1.56830E-02	1.05923E-02
400000.	-4.19480E+01	-3.48038E-02	-4.56039E-02	-3.81117E-02	-2.13775E-02	-6.73960E-03	1.93438E-03
450000.	-4.20718E+01	-4.96888E-03	-8.42094E-03	-1.19219E-02	-1.19411E-02	-8.70975E-03	-4.48543E-03
500000.	-4.21803E+01	1.24481E-02	3.32025E-03	-3.09575E-03	-5.02533E-03	-4.35643E-03	-3.20289E-03
550000.	-4.22344E+01	5.60707E-03	3.46072E-03	4.86332E-04	-1.43623E-03	-1.96514E-03	-1.82432E-03
600000.	-4.22797E+01	-3.01170E-04	-3.84669E-04	-9.55080E-04	-1.27882E-03	-1.24680E-03	-1.16149E-03

See appendix for Chebyshev fit details.

$$Li + H \rightarrow H^- + Li^+$$



Electron Capture Cross Sections for

$H + H^- \rightarrow H^- + H$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.0E+00	1.96E+06	1.33E-14
4.0E+00	2.78E+06	1.24E-14
7.0E+00	3.68E+06	1.15E-14
1.0E+01	4.39E+06	1.07E-14
2.0E+01	6.21E+06	8.97E-15
4.0E+01	8.79E+06	7.24E-15
7.0E+01	1.16E+07	5.99E-15
9.9E+01	1.38E+07	5.27E-15
2.0E+02	1.96E+07	4.20E-15
4.0E+02	2.78E+07	3.19E-15
7.0E+02	3.68E+07	2.43E-15
1.0E+03	4.39E+07	1.90E-15
2.0E+03	6.21E+07	9.49E-16
4.0E+03	8.79E+07	3.29E-16
7.1E+03	1.17E+08	1.03E-16

References: 162, 271, 272, 281

Accuracy: 30%

Note: For energies less than 40 eV the data were taken from the theoretical papers of Ref. 271 and 272.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 2.0E+00 eV/amu, E_{max} = 7.1E+03 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-67.0414	-2.13773	-.748888	-.288550	-.136764	-.0116298	.0247820	.00770226	2.25210E-04

The fit represents the above cross sections with an rms deviation of 0.3%.
The maximum deviation is 0.7% at 4.0E+02 eV/amu.
See appendix for Chebyshev fit details.

$H + H^- \rightarrow H^- + H$

Cross Section vs. Energy



Electron Capture Cross Sections for
$H^+ + H \rightarrow H + H^+$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.2E-01	4.79E+05	4.96E-15
2.0E-01	6.21E+05	4.70E-15
4.0E-01	8.79E+05	4.45E-15
7.0E-01	1.16E+06	4.24E-15
1.0E+00	1.39E+06	4.10E-15
2.0E+00	1.96E+06	3.83E-15
4.0E+00	2.78E+06	3.58E-15
7.0E+00	3.68E+06	3.33E-15
1.0E+01	4.39E+06	3.17E-15
2.0E+01	6.21E+06	2.93E-15
4.0E+01	8.79E+06	2.70E-15
7.0E+01	1.16E+07	2.54E-15
1.0E+02	1.39E+07	2.44E-15
2.0E+02	1.96E+07	2.22E-15
4.0E+02	2.78E+07	2.03E-15
7.0E+02	3.68E+07	1.88E-15
1.0E+03	4.39E+07	1.71E-15
2.0E+03	6.21E+07	1.44E-15
4.0E+03	8.79E+07	1.20E-15
7.0E+03	1.16E+08	9.42E-16
1.0E+04	1.39E+08	7.75E-16
2.0E+04	1.96E+08	4.45E-16
4.0E+04	2.78E+08	1.67E-16
7.0E+04	3.68E+08	3.77E-17
1.0E+05	4.39E+08	1.01E-17
2.0E+05	6.21E+08	6.09E-19
4.0E+05	8.78E+08	1.76E-20
6.3E+05	1.10E+09	1.35E-21

References: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 273, 395

Accuracy: H^+ Energy: < 10 eV/amu, 10%; 10 eV/amu - $1x10^3$ eV/amu, 15%; $1x10^3$ eV/amu - $1x10^5$ eV/amu, 5%; $1x10^5$ eV/amu - $6.25x10^5$ eV/amu, 20%

Notes: (1) This reaction has been measured by three different techniques: 1. Static Gas Target, 2. Cross Beams, 3. Merged Beams.
(2) The maximum deviation of the recommended data from the data points in the references cited is less than 45%.
(3) Newman in reference 9 found that when D atoms were the target the cross section decreased 16% from that using H atoms at an energy of .12 eV. At high energies the cross sections were the same.
(4) The recent ref. 273 extends the measurements up to an energy of $7.5x10^6$ eV/amu. The cross sections are (a) $2.0x10^6$ eV/amu - $5.25x10^{-24}$ cm^2; (b) $3.5x10^6$ eV/amu - $2.14x10^{-25}$ cm^2; (c) $5.0x10^6$ eV/amu - $3.1x10^{-26}$ cm^2; (d) $7.5x10^6$ eV/amu - $4.10x10^{-27}$ cm^2. Quoted errors are as great as 43%.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 1.2E-01 eV/amu, E_{max} = 6.3E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-72.6656	-5.49142	-3.42948	-1.98377	-.878009	-.198932	.0837431	.121252	.0827182

The fit represents the above cross sections with an rms deviation of 4.5%.
The maximum deviation is 13.5% at 1.0E+05 eV/amu.
See appendix for Chebyshev fit details.

$H^+ + H \rightarrow H + H^+$



Electron Capture Rate Coefficients for
$H^+ + H \rightarrow H + H^+$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	Equal Temp.	H Temp. (eV) 1.	10.	100.	1000.	5000.	10000.	20000.
1.0E+00	8.10E-09	8.10E-09	1.56E-08	3.63E-08	7.65E-08	9.74E-08	8.95E-08	6.84E-08
2.0E+00	1.06E-08	9.50E-09	1.61E-08	3.64E-08	7.65E-08	9.74E-08	8.95E-08	6.84E-08
4.0E+00	1.38E-08	1.16E-08	1.71E-08	3.67E-08	7.66E-08	9.74E-08	8.95E-08	6.84E-08
7.0E+00	1.71E-08	1.38E-08	1.85E-08	3.71E-08	7.66E-08	9.74E-08	8.95E-08	6.84E-08
1.0E+01	1.96E-08	1.56E-08	1.96E-08	3.75E-08	7.67E-08	9.74E-08	8.95E-08	6.84E-08
2.0E+01	2.56E-08	2.00E-08	2.30E-08	3.87E-08	7.69E-08	9.74E-08	8.95E-08	6.84E-08
4.0E+01	3.33E-08	2.59E-08	2.79E-08	4.10E-08	7.73E-08	9.74E-08	8.94E-08	6.84E-08
7.0E+01	4.10E-08	3.18E-08	3.33E-08	4.40E-08	7.78E-08	9.74E-08	8.94E-08	6.83E-08
1.0E+02	4.66E-08	3.63E-08	3.75E-08	4.66E-08	7.84E-08	9.74E-08	8.93E-08	6.83E-08
2.0E+02	5.88E-08	4.67E-08	4.74E-08	5.36E-08	8.00E-08	9.73E-08	8.91E-08	6.81E-08
4.0E+02	7.21E-08	5.89E-08	5.93E-08	6.30E-08	8.30E-08	9.72E-08	8.87E-08	6.77E-08
7.0E+02	8.30E-08	6.95E-08	6.98E-08	7.21E-08	8.65E-08	9.70E-08	8.80E-08	6.72E-08
1.0E+03	8.92E-08	7.65E-08	7.67E-08	7.84E-08	8.92E-08	9.67E-08	8.73E-08	6.66E-08
2.0E+03	9.70E-08	8.92E-08	8.93E-08	9.00E-08	9.48E-08	9.54E-08	8.51E-08	6.49E-08
4.0E+03	9.36E-08	9.70E-08	9.70E-08	9.71E-08	9.74E-08	9.17E-08	8.06E-08	6.16E-08
7.0E+03	8.06E-08	9.54E-08	9.54E-08	9.52E-08	9.36E-08	8.51E-08	7.43E-08	5.70E-08
1.0E+04	6.84E-08	8.95E-08	8.95E-08	8.93E-08	8.73E-08	7.84E-08	6.84E-08	5.30E-08
2.0E+04	4.23E-08	6.84E-08	6.84E-08	6.83E-08	6.66E-08	6.00E-08	5.30E-08	4.23E-08

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

H Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
1.	-3.44869E+01	1.27901E+00	-2.75515E-01	-1.94518E-01	-4.14913E-02	-1.79417E-02	-4.17985E-03
10.	-3.41227E+01	9.80499E-01	-1.17702E-01	-2.36330E-01	-4.90489E-02	-5.73981E-03	-8.76848E-03
100.	-3.34478E+01	5.02257E-01	-2.99001E-03	-1.75672E-01	-9.20126E-02	-1.06753E-02	3.21804E-03
1000.	-3.26740E+01	4.69533E-02	-4.61309E-02	-8.81367E-02	-6.79803E-02	-2.93738E-02	-4.42210E-03
5000.	-3.24495E+01	-1.48073E-01	-1.15160E-01	-7.59145E-02	-4.13891E-02	-1.79143E-02	-5.55664E-03
10000.	-3.26521E+01	-1.74458E-01	-1.25225E-01	-7.36305E-02	-3.55149E-02	-1.38261E-02	-4.10388E-03
20000.	-3.31787E+01	-1.63538E-01	-1.15089E-01	-6.54004E-02	-3.01053E-02	-1.10769E-02	-3.10752E-03
Equal Temp.	-3.44313E+01	1.06711E+00	-5.19907E-01	-2.29247E-01	-6.98896E-02	-1.36578E-02	4.42580E-04

See appendix for Chebyshev fit details.

$H^+ + H \rightarrow H + H^+$

Maxwellian – Maxwellian



Electron Capture Rate Coefficients for
$H + H^+ \rightarrow H + H^+$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	H Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	1.08E-07	8.72E-08	4.62E-08	1.38E-08	4.43E-09	3.78E-10	4.79E-12**
2.0E+00	1.07E-07	8.71E-08	4.62E-08	1.38E-08	4.43E-09	3.78E-10	4.79E-12**
4.0E+00	1.07E-07	8.70E-08	4.61E-08	1.38E-08	4.43E-09	3.77E-10	4.79E-12**
7.0E+00	1.07E-07	8.69E-08	4.60E-08	1.38E-08	4.43E-09	3.77E-10	4.79E-12**
1.0E+01	1.07E-07	8.68E-08	4.59E-08	1.38E-08	4.43E-09	3.77E-10	4.79E-12**
2.0E+01	1.07E-07	8.65E-08	4.57E-08	1.37E-08	4.43E-09	3.77E-10	4.79E-12**
4.0E+01	1.07E-07	8.61E-08	4.55E-08	1.37E-08	4.44E-09	3.76E-10	4.80E-12**
7.0E+01	1.06E-07	8.57E-08	4.52E-08	1.37E-08	4.44E-09	3.76E-10	4.80E-12**
1.0E+02	1.06E-07	8.53E-08	4.50E-08	1.37E-08	4.45E-09	3.76E-10	4.81E-12**
2.0E+02	1.05E-07	8.45E-08	4.45E-08	1.37E-08	4.48E-09	3.77E-10	4.83E-12**
4.0E+02	1.04E-07	8.34E-08	4.39E-08	1.37E-08	4.55E-09	3.79E-10	4.88E-12**
7.0E+02	1.03E-07	8.21E-08	4.34E-08	1.38E-08	4.67E-09	3.84E-10	4.96E-12**
1.0E+03	1.01E-07	8.10E-08	4.31E-08	1.40E-08	4.79E-09	3.89E-10	5.03E-12**
2.0E+03	9.70E-08	7.79E-08	4.23E-08	1.46E-08	5.17E-09	4.11E-10	5.29E-12**
4.0E+03	9.00E-08	7.24E-08	4.11E-08	1.56E-08	5.91E-09	4.64E-10	5.82E-12*
7.0E+03	8.16E-08	6.57E-08	3.92E-08	1.66E-08	6.90E-09	5.69E-10	6.71E-12*
1.0E+04	7.46E-08	6.03E-08	3.74E-08	1.72E-08	7.74E-09	7.01E-10	7.80E-12*
2.0E+04	5.70E-08	4.72E-08	3.20E-08	1.75E-08	9.46E-09	1.28E-09	1.36E-11*

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

H Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-3.23588E+01	-2.28428E-01	-1.47539E-01	-7.73418E-02	-3.29002E-02	-1.09141E-02	-2.93158E-03
20000.	-3.27829E+01	-2.26775E-01	-1.38750E-01	-7.04432E-02	-3.03096E-02	-1.00013E-02	-1.47228E-03
40000.	-3.39547E+01	-1.36550E-01	-7.15786E-02	-3.65682E-02	-2.10115E-02	-1.11355E-02	-4.14062E-03
70000.	-3.60736E+01	1.09084E-01	7.35296E-02	2.31264E-02	-7.74992E-03	-1.38165E-02	-8.49906E-03
100000.	-3.81082E+01	3.09107E-01	1.95996E-01	8.14408E-02	1.35787E-02	-9.21346E-03	-1.02081E-02
200000.	-4.29608E+01	3.96726E-01	2.99000E-01	1.81207E-01	8.74182E-02	3.26701E-02	7.33350E-03
500000.	-5.17564E+01	3.37283E-01	2.43298E-01	1.48098E-01	7.72749E-02	3.54105E-02	1.56884E-02

See appendix for Chebyshev fit details.

$$H + H^+ \rightarrow H + H^+$$

Beam – Maxwellian



Electron Capture Cross Sections for
$H^+ + H_2 \rightarrow H$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.6E+00	2.24E+06	1.81E-17
4.0E+00	2.78E+06	4.96E-17
4.7E+00	3.01E+06	5.68E-17
7.0E+00	3.68E+06	2.76E-17
1.0E+01	4.39E+06	1.30E-17
1.4E+01	5.20E+06	1.01E-17
2.0E+01	6.21E+06	1.11E-17
4.0E+01	8.79E+06	1.68E-17
7.0E+01	1.16E+07	2.61E-17
1.0E+02	1.39E+07	3.65E-17
2.0E+02	1.96E+07	6.96E-17
4.0E+02	2.78E+07	1.56E-16
7.0E+02	3.68E+07	2.93E-16
1.0E+03	4.39E+07	4.26E-16
2.0E+03	6.21E+07	6.88E-16
4.0E+03	8.79E+07	9.33E-16
7.0E+03	1.16E+08	9.57E-16
1.0E+04	1.39E+08	8.66E-16
2.0E+04	1.96E+08	5.79E-16
4.0E+04	2.78E+08	2.50E-16
7.0E+04	3.68E+08	7.78E-17
1.0E+05	4.39E+08	2.91E-17
2.0E+05	6.21E+08	1.76E-18
4.0E+05	8.78E+08	6.09E-20
7.0E+05	1.16E+09	3.64E-21
1.0E+06	1.39E+09	4.14E-22
2.0E+06	1.96E+09	1.68E-23
4.0E+06	2.77E+09	1.03E-24

References: 7, 27, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 47, 48, 62, 273, 395

Accuracy: E < 100 eV/amu - Unknown; E > 100 eV/amu - 10%

Notes: (1) The peak in the cross section at 4 eV comes from the data of reference 41. The origin of this maximum is unknown.
(2) Recent data of reference 273 extend the energy range to 7.5×10^6 eV/amu. The cross sections reported are: (a) 2.0×10^6 eV/amu - 7.00×10^{-24} cm^2; (b) 3.5×10^6 eV/amu - 4.85×10^{-25} cm^2; (c) 5.0×10^6 eV/amu - 6.47×10^{-26} cm^2; (d) 7.5×10^6 eV/amu - 6.30×10^{-27} cm^2. Errors of 10-15% are reported.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 2.6E+00 eV/amu, E_{max} = 4.0E+06 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-82.5164	-6.70755	-6.10977	-2.62810	.709759	.639033	.102980	.261240	-.263817

The fit represents the above cross sections with an rms deviation of 15.5%.
The maximum deviation is 41.3% at 1.0E+01 eV/amu.
See appendix for Chebyshev fit details.

$H^+ + H_2 \rightarrow H$



Double Electron Capture Cross Sections for
$H^+ + H_2 \rightarrow H^- + 2H^+$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.0E+02	1.96E+07	1.17E-20
4.0E+02	2.78E+07	2.22E-20
7.0E+02	3.68E+07	4.82E-20
1.0E+03	4.39E+07	8.94E-20
2.0E+03	6.21E+07	2.80E-19
4.0E+03	8.79E+07	8.64E-19
7.0E+03	1.16E+08	2.09E-18
1.0E+04	1.39E+08	3.63E-18
2.0E+04	1.96E+08	8.89E-18
4.0E+04	2.78E+08	2.21E-18
7.0E+04	3.68E+08	2.42E-19
1.0E+05	4.39E+08	3.15E-20
2.0E+05	6.21E+08	1.84E-22
4.0E+05	8.78E+08	7.22E-25
7.0E+05	1.16E+09	6.64E-27
1.0E+06	1.39E+09	2.70E-28

References: 35, 44, 50, 51, 52, 53, 54

Accuracy: $E < 1x10^4$ eV/amu - 25%; $1x10^4$ eV/amu $< E < 1x10^5$ eV/amu - 15%; $E > 1x10^5$ eV/amu - 40%

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 2.0E+02 eV/amu, E_{max} = 1.0E+06 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-95.8165	-7.17049	-7.48288	-1.93034	.761153	.556689	-.0542859	-.270184	-.0147551

The fit represents the above cross sections with an rms deviation of 9.4%.
The maximum deviation is 15.0% at 1.0E+04 eV/amu.
See appendix for Chebyshev fit details.

$$H^+ + H_2 \rightarrow H^- + 2H^+$$



Electron Capture Cross Sections for
$H^+ + He \rightarrow H + He^+$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
9.9E+01	1.38E+07	4.28E-22
2.0E+02	1.96E+07	2.46E-21
4.0E+02	2.78E+07	1.77E-20
7.0E+02	3.68E+07	1.12E-19
1.0E+03	4.39E+07	4.73E-19
2.0E+03	6.21E+07	4.25E-18
4.0E+03	8.79E+07	1.98E-17
7.0E+03	1.16E+08	5.18E-17
1.0E+04	1.39E+08	8.52E-17
2.0E+04	1.96E+08	1.75E-16
2.4E+04	2.15E+08	1.83E-16
4.0E+04	2.78E+08	1.41E-16
7.0E+04	3.68E+08	6.98E-17
1.0E+05	4.39E+08	3.38E-17
2.0E+05	6.21E+08	3.95E-18
4.0E+05	8.78E+08	3.22E-19
7.0E+05	1.16E+09	3.07E-20
1.0E+06	1.39E+09	5.42E-21
2.0E+06	1.96E+09	1.42E-22
4.0E+06	2.77E+09	3.29E-24
7.0E+06	3.65E+09	1.62E-25
1.0E+07	4.36E+09	2.09E-26
1.1E+07	4.57E+09	1.02E-26

References: 27, 35, 37, 40, 47, 49, 51, 82, 83, 84, 85, 86, 87, 88, 89, 272

Accuracy: $E < 5x10^3$ eV/amu - 40%; $E > 5x10^3$ eV/amu - 20%

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 9.9E+01 eV/amu, E_{max} = 1.1E+07 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-92.7819	-4.69073	-8.99666	-.490484	.935276	-.0504853	-.0410750	-.0730136	.0189320

The fit represents the above cross sections with an rms deviation of 10.0%.
The maximum deviation is 18.7% at 7.0E+02 eV/amu.
See appendix for Chebyshev fit details.

$H^+ + He \rightarrow H + He^+$



Electron Capture Rate Coefficients for
$H^+ + He \rightarrow H + He^+$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	Equal Temp.	He Temp. (eV) 50.	100.	500.	1000.	5000.	10000.	20000.
7.0E+01	7.21E-14	5.81E-14	9.77E-14	1.60E-12	1.02E-11	6.38E-10	2.68E-09	8.50E-09
1.0E+02	2.79E-13	1.85E-13	2.79E-13	2.78E-12	1.39E-11	6.73E-10	2.74E-09	8.57E-09
2.0E+02	4.14E-12	2.23E-12	2.78E-12	1.08E-11	3.22E-11	7.95E-10	2.95E-09	8.81E-09
4.0E+02	4.51E-11	2.43E-11	2.68E-11	5.24E-11	9.92E-11	1.07E-09	3.37E-09	9.28E-09
7.0E+02	2.27E-10	1.29E-10	1.36E-10	1.94E-10	2.82E-10	1.53E-09	4.03E-09	9.99E-09
1.0E+03	5.60E-10	3.32E-10	3.42E-10	4.33E-10	5.60E-10	2.06E-09	4.73E-09	1.07E-08
2.0E+03	2.54E-09	1.64E-09	1.66E-09	1.83E-09	2.06E-09	4.15E-09	7.12E-09	1.29E-08
4.0E+03	8.33E-09	5.94E-09	5.97E-09	6.21E-09	6.52E-09	8.93E-09	1.18E-08	1.66E-08
7.0E+03	1.62E-08	1.29E-08	1.29E-08	1.31E-08	1.34E-08	1.53E-08	1.74E-08	2.07E-08
1.0E+04	2.12E-08	1.82E-08	1.82E-08	1.83E-08	1.85E-08	1.98E-08	2.12E-08	2.33E-08
2.0E+04	2.60E-08	2.55E-08	2.55E-08	2.55E-08	2.55E-08	2.57E-08	2.59E-08	2.60E-08

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 7.0E+01 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
50.	-4.54767E+01	6.67186E+00	-1.30388E+00	-1.33476E-01	4.39632E-02	-4.52889E-02	1.85831E-02
100.	-4.51974E+01	6.44612E+00	-1.18362E+00	-1.72127E-01	4.66619E-02	-4.04263E-02	1.55388E-02
500.	-4.35812E+01	5.17359E+00	-5.65656E-01	-3.28985E-01	3.26661E-02	-7.75934E-03	1.09010E-03
1000.	-4.23753E+01	4.27018E+00	-2.07470E-01	-3.69408E-01	-9.50854E-04	1.17863E-02	-8.41076E-04
5000.	-3.90038E+01	2.05191E+00	2.49124E-01	-2.12101E-01	-8.19217E-02	7.92950E-03	1.06018E-02
10000.	-3.75454E+01	1.24999E+00	2.32693E-01	-1.10685E-01	-6.99375E-02	-5.62572E-03	6.74424E-03
20000.	-3.62570E+01	6.12670E-01	1.45612E-01	-4.25717E-02	-4.13562E-02	-1.03361E-02	9.23441E-04
Equal Temp.	-4.47894E+01	6.48647E+00	-1.50363E+00	-5.56465E-02	1.00897E-02	-3.20031E-02	2.61002E-02

See appendix for Chebyshev fit details.

$H^+ + He \rightarrow H + He^+$

Maxwellian – Maxwellian



Electron Capture Rate Coefficients for
$He + H^+ \rightarrow H + He^+$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	1.18E-08	3.43E-08	3.91E-08	2.56E-08	1.48E-08	2.45E-09	1.24E-10
2.0E+00	1.18E-08	3.42E-08	3.91E-08	2.56E-08	1.48E-08	2.45E-09	1.24E-10
4.0E+00	1.18E-08	3.42E-08	3.90E-08	2.56E-08	1.48E-08	2.45E-09	1.24E-10
7.0E+00	1.18E-08	3.41E-08	3.90E-08	2.55E-08	1.48E-08	2.45E-09	1.24E-10
1.0E+01	1.18E-08	3.41E-08	3.89E-08	2.55E-08	1.48E-08	2.45E-09	1.24E-10
2.0E+01	1.18E-08	3.40E-08	3.88E-08	2.55E-08	1.47E-08	2.45E-09	1.24E-10
4.0E+01	1.19E-08	3.38E-08	3.87E-08	2.54E-08	1.47E-08	2.45E-09	1.24E-10
7.0E+01	1.19E-08	3.37E-08	3.85E-08	2.53E-08	1.46E-08	2.45E-09	1.24E-10
1.0E+02	1.20E-08	3.36E-08	3.84E-08	2.53E-08	1.46E-08	2.45E-09	1.24E-10
2.0E+02	1.24E-08	3.33E-08	3.80E-08	2.51E-08	1.45E-08	2.45E-09	1.24E-10
4.0E+02	1.31E-08	3.29E-08	3.76E-08	2.49E-08	1.45E-08	2.45E-09	1.25E-10
7.0E+02	1.41E-08	3.25E-08	3.70E-08	2.47E-08	1.44E-08	2.47E-09	1.26E-10
1.0E+03	1.51E-08	3.21E-08	3.66E-08	2.46E-08	1.44E-08	2.48E-09	1.27E-10
2.0E+03	1.77E-08	3.14E-08	3.53E-08	2.42E-08	1.44E-08	2.55E-09	1.30E-10
4.0E+03	2.11E-08	3.05E-08	3.30E-08	2.34E-08	1.44E-08	2.70E-09	1.36E-10
7.0E+03	2.39E-08	2.95E-08	3.03E-08	2.23E-08	1.43E-08	2.94E-09	1.45E-10
1.0E+04	2.54E-08	2.87E-08	2.82E-08	2.12E-08	1.41E-08	3.18E-09	1.55E-10
2.0E+04	2.64E-08	2.61E-08	2.37E-08	1.83E-08	1.32E-08	3.84E-09	1.94E-10

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-3.59874E+01	4.17430E-01	1.93060E-01	1.41316E-02	-4.62466E-02	-3.20698E-02	-6.14466E-03
20000.	-3.45210E+01	-1.10051E-01	-5.31364E-02	-2.09508E-02	-8.28916E-03	-4.29131E-03	-2.80760E-03
40000.	-3.43497E+01	-1.93363E-01	-1.15051E-01	-5.41354E-02	-1.98052E-02	-4.14043E-03	1.32023E-03
70000.	-3.51059E+01	-1.20106E-01	-7.36757E-02	-4.02144E-02	-2.03546E-02	-8.65732E-03	-2.24607E-03
100000.	-3.61082E+01	-3.88036E-02	-1.73976E-02	-1.16387E-02	-1.05935E-02	-7.51826E-03	-3.79933E-03
200000.	-3.94814E+01	1.55478E-01	1.14512E-01	6.41579E-02	2.61502E-02	6.20889E-03	-1.41106E-03
500000.	-4.54567E+01	1.49642E-01	1.06212E-01	6.21478E-02	3.03614E-02	1.26308E-02	4.94057E-03

See appendix for Chebyshev fit details.

$He + H^+ \rightarrow H + He^+$

Beam – Maxwellian



Double Electron Capture Cross Sections for
$H^+ + He \rightarrow H^- + He^{2+}$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
4.7E+02	3.01E+07	2.70E-22
7.0E+02	3.68E+07	5.75E-22
1.0E+03	4.37E+07	1.12E-21
2.0E+03	6.21E+07	4.20E-21
4.0E+03	8.89E+07	1.78E-20
7.0E+03	1.16E+08	5.08E-20
1.0E+04	1.38E+08	1.06E-19
2.0E+04	1.96E+08	3.75E-19
3.2E+04	2.48E+08	7.08E-19
4.0E+04	2.78E+08	6.42E-19
7.0E+04	3.65E+08	1.73E-19
1.0E+05	4.37E+08	3.76E-20
2.0E+05	6.21E+08	9.67E-22
4.0E+05	8.78E+08	7.64E-24
7.0E+05	1.15E+09	9.68E-26
1.0E+06	1.38E+09	3.76E-27

References: 35, 50, 52, 53, 54

Accuracy: $E < 1x10^4$ eV/amu - factor 2 or more; $E > 1x10^4$ eV/amu - 40%

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 4.7E+02 eV/amu, E_{max} = 9.9E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-98.3453	-4.13574	-6.43185	-1.70697	.435158	.417370	-.125015	-.191020	.0356727

The fit represents the above cross sections with an rms deviation of 10.0%.
The maximum deviation is 20.7% at 2.0E+04 eV/amu.
See appendix for Chebyshev fit details.

$H^+ + He \rightarrow H^- + He^{2+}$

Cross Section vs. Energy



Electron Capture Cross Sections for
$H^+ + Li \rightarrow H + Li^+$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.5E+02	2.20E+07	1.75E-16
3.0E+02	2.41E+07	2.38E-16
4.0E+02	2.78E+07	3.79E-16
5.0E+02	3.11E+07	5.40E-16
7.0E+02	3.68E+07	8.95E-16
1.0E+03	4.39E+07	1.40E-15
2.0E+03	6.21E+07	3.28E-15
3.0E+03	7.61E+07	4.26E-15
3.6E+03	8.33E+07	4.35E-15
5.0E+03	9.82E+07	4.07E-15
8.0E+03	1.24E+08	2.68E-15
1.0E+04	1.39E+08	1.80E-15
1.5E+04	1.70E+08	7.52E-16
3.0E+04	2.41E+08	7.98E-17
5.0E+04	3.11E+08	2.84E-17
8.0E+04	3.93E+08	1.77E-17
1.5E+05	5.38E+08	8.73E-18
2.0E+05	6.21E+08	5.21E-18
3.0E+05	7.61E+08	1.88E-18
4.0E+05	8.78E+08	7.17E-19
5.0E+05	9.82E+08	2.93E-19
5.6E+05	1.04E+09	2.02E-19

References: 230, 231, 232, 233, 234, 235, 236, 237, 238, 239

Accuracy: 30%

Notes: (1) Gruebler et al. (Ref. 235) found up to a factor of 2 greater cross section when H^+ was the projectile compared to D^+ incident at equivalent velocities.
(2) For data on electron capture cross sections of $Li^+ + H$ see references 247, 248, 252, 253.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 2.5E+02 eV/amu, E_{max} = 5.6E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-74.2905	-3.64125	-2.48131	.598612	.0178325	-.507845	-.166745	.177271	.132733

The fit represents the above cross sections with an rms deviation of 6.4%.
The maximum deviation is 17.7% at 3.0E+04 eV/amu.
See appendix for Chebyshev fit details.

$H^+ + Li \rightarrow H + Li^+$



Electron Capture Rate Coefficients for
$Li + H^+ \rightarrow H + Li^+$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	Li Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	2.50E-07	5.82E-08	1.24E-08	7.44E-09	6.05E-09*	3.23E-09*	2.88E-10**
2.0E+00	2.49E-07	5.83E-08	1.24E-08	7.44E-09	6.05E-09*	3.23E-09*	2.88E-10**
4.0E+00	2.49E-07	5.83E-08	1.24E-08	7.44E-09	6.05E-09*	3.23E-09*	2.88E-10**
7.0E+00	2.49E-07	5.84E-08	1.24E-08	7.44E-09	6.05E-09*	3.23E-09*	2.88E-10**
1.0E+01	2.49E-07	5.84E-08	1.24E-08	7.44E-09	6.05E-09*	3.23E-09*	2.88E-10**
2.0E+01	2.48E-07	5.86E-08	1.24E-08	7.44E-09	6.05E-09*	3.22E-09*	2.89E-10**
4.0E+01	2.48E-07	5.89E-08	1.24E-08	7.44E-09	6.05E-09	3.22E-09*	2.89E-10**
7.0E+01	2.48E-07	5.95E-08	1.24E-08	7.44E-09	6.05E-09	3.21E-09*	2.89E-10**
1.0E+02	2.47E-07	6.00E-08	1.24E-08	7.44E-09	6.05E-09	3.21E-09*	2.90E-10**
2.0E+02	2.44E-07	6.17E-08	1.25E-08	7.44E-09	6.05E-09	3.20E-09*	2.91E-10**
4.0E+02	2.39E-07	6.46E-08	1.27E-08	7.44E-09	6.05E-09	3.19E-09*	2.93E-10**
7.0E+02	2.30E-07	6.81E-08	1.30E-08	7.44E-09	6.05E-09	3.18E-09*	2.93E-10**
1.0E+03	2.21E-07	7.10E-08	1.35E-08	7.45E-09	6.05E-09	3.17E-09*	2.91E-10**
2.0E+03	1.96E-07	7.73E-08	1.56E-08	7.53E-09	6.04E-09	3.16E-09*	2.86E-10**
4.0E+03	1.64E-07	8.02E-08	2.04E-08	7.94E-09	6.04E-09	3.14E-09*	2.85E-10**
7.0E+03	1.34E-07	7.67E-08	2.59E-08	9.12E-09	6.14E-09	3.11E-09	2.94E-10**
1.0E+04	1.14E-07	7.14E-08	2.90E-08	1.06E-08	6.44E-09	3.08E-09	3.08E-10**
2.0E+04	7.50E-08	5.52E-08	3.08E-08	1.44E-08	8.16E-09	3.06E-09	3.61E-10**

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

Li Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-3.09481E+01	-4.64901E-01	-2.95108E-01	-1.33838E-01	-3.59323E-02	-1.29218E-03	8.76916E-04
20000.	-3.31581E+01	9.29024E-02	-3.27874E-02	-9.34862E-02	-7.42714E-02	-2.98862E-02	-6.36610E-04
40000.	-3.59040E+01	4.30313E-01	2.62445E-01	8.26693E-02	-2.56118E-02	-5.35625E-02	-3.78019E-02
70000.	-3.72045E+01	2.09566E-01	1.66816E-01	1.08374E-01	5.50349E-02	1.81051E-02	-2.94289E-03
100000.	-3.77748E+01	7.11403E-02	6.19800E-02	5.19417E-02	3.83027E-02	2.39693E-02	1.32084E-02
200000.	-3.91351E+01	-2.54412E-02	-9.95070E-03	-2.34861E-03	-2.74660E-04	3.66029E-04	9.29558E-04
500000.	-4.38736E+01	5.64031E-02	4.45512E-02	3.82771E-02	3.08448E-02	1.91863E-02	5.93228E-03

See appendix for Chebyshev fit details.

$$Li + H^+ \rightarrow H + Li^+$$

Beam – Maxwellian



Double Electron Capture Cross Sections for
$H^+ + H^- \rightarrow H^- + H^+$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
5.9E+01	1.07E+07	9.98E-17
6.4E+01	1.11E+07	1.02E-16
6.9E+01	1.15E+07	8.45E-17
7.4E+01	1.19E+07	5.63E-17
8.0E+01	1.24E+07	3.96E-17
8.1E+01	1.25E+07	3.95E-17
8.4E+01	1.27E+07	4.63E-17
8.9E+01	1.31E+07	6.11E-17
9.4E+01	1.35E+07	7.54E-17
9.9E+01	1.38E+07	8.72E-17
1.0E+02	1.39E+07	8.82E-17
1.1E+02	1.46E+07	6.84E-17
1.2E+02	1.52E+07	4.70E-17
1.3E+02	1.58E+07	4.23E-17
1.4E+02	1.64E+07	4.84E-17
1.5E+02	1.70E+07	6.33E-17
1.6E+02	1.76E+07	7.20E-17
1.7E+02	1.81E+07	6.12E-17
1.8E+02	1.86E+07	4.27E-17
1.9E+02	1.91E+07	4.10E-17
2.0E+02	1.96E+07	4.62E-17
2.5E+02	2.20E+07	8.33E-17
3.0E+02	2.41E+07	1.08E-16
3.5E+02	2.60E+07	1.27E-16
4.0E+02	2.78E+07	1.44E-16
4.5E+02	2.95E+07	1.50E-16
4.7E+02	3.01E+07	1.51E-16
5.0E+02	3.11E+07	1.50E-16
6.0E+02	3.40E+07	1.40E-16
8.0E+02	3.93E+07	1.14E-16
8.4E+02	4.03E+07	1.07E-16
1.0E+03	4.39E+07	8.79E-17
1.1E+03	4.61E+07	7.48E-17

References: 69, 274, 275, 276

Accuracy: 30%

Notes: (1) The oscillatory behaviour results from the formation of a quasi-H_2 molecule.
(2) The cross section as plotted is an average of that measured by Peart (Ref. 274) and that of Brouillard et al. (Ref. 275).
(3) Plotted are the H^+ energies with the H^- at rest.
(4) The interaction energy is given by $E_{int}=\frac{E_{H^+}}{2}$ for equal masses.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 5.9E+01 eV/amu, E_{max} = 1.1E+03 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-74.0098	.227379	.0232305	-.388332	-.00604588	.0351500	.101175	-.132693	-.0302977

The fit represents the above cross sections with an rms deviation of 2[illegible].9%.
The maximum deviation is 46.4% at 8.1E+01 eV/amu.
See appendix for Chebyshev fit details.

$H^+ + H^- \rightarrow H^- + H^+$



Mutual Neutralization Cross Sections for
$H^+ + H^- \rightarrow H + H$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.0E-01	6.21E+05	1.69E-13
4.0E-01	8.79E+05	9.09E-14
7.0E-01	1.16E+06	5.74E-14
1.0E+00	1.39E+06	4.27E-14
2.0E+00	1.96E+06	2.40E-14
4.0E+00	2.78E+06	1.40E-14
7.0E+00	3.68E+06	9.21E-15
1.0E+01	4.39E+06	7.36E-15
2.0E+01	6.21E+06	6.53E-15
4.0E+01	8.79E+06	7.11E-15
7.0E+01	1.16E+07	8.20E-15
1.0E+02	1.39E+07	8.80E-15
2.0E+02	1.96E+07	1.06E-14
4.0E+02	2.78E+07	1.30E-14
7.0E+02	3.68E+07	1.39E-14
1.0E+03	4.39E+07	1.38E-14
2.0E+03	6.21E+07	1.17E-14
4.0E+03	8.79E+07	7.54E-15
4.8E+03	9.62E+07	6.36E-15

References: 63, 64, 65, 66, 67, 68, 69, 280

Accuracy: E > 20 eV/amu - 20%; E < 20 eV/amu - factor 2 (see notes)

Notes: (1) Measurements made in the 1970's showed structure in the cross section curve in the energy range of 40 - 700 eV/amu. More recent measurements (Ref. 67 and 68) show no evidence of structure and the cross sections are approximately a factor of 2 less than those reported in Refs. 63-66 for energies less than 400 eV.
(2) Mosley (Ref. 63) found a sharp rise in the cross section for energies less than 30 eV/amu. In the region of overlap, with the most recent papers, Mosely was a factor of two greater. Thus, the data of Mosely have been reduced by approximately a factor of two which more nearly coincides with the theoretical data of Ref. 280.
(3) The data below 10 eV/amu should be used with caution.
(4) Plotted is the H^+ energy with H^- at rest.
(5) The interaction energy is given by $E_{int}=E_{H^+}/2$.
(6) See Dolder's review (Ref. 69) for a discussion of this reaction.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 2.0E-01 eV/amu, E_{max} = 4.8E+03 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-63.1439	-1.23277	.753504	-.408901	-.276190	.0381613	.0408940	-.0423914	.00269073

The fit represents the above cross sections with an rms deviation of 2.4%.
The maximum deviation is 5.8% at 1.0E+01 eV/amu.
See appendix for Chebyshev fit details.

$H^+ + H^- \rightarrow H + H$



Sum of the Electron Capture and Ionization Cross Sections for
$H^+ + He^+ \rightarrow [(H + He^{2+}) + (H^+ + He^{2+} + e^-)]$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
3.3E+03	7.98E+07	7.43E-20
4.0E+03	8.79E+07	1.04E-19
7.0E+03	1.16E+08	3.62E-19
1.0E+04	1.39E+08	9.89E-19
2.0E+04	1.96E+08	1.00E-17
4.0E+04	2.78E+08	2.31E-17
7.0E+04	3.68E+08	2.55E-17
1.0E+05	4.39E+08	2.19E-17
2.0E+05	6.21E+08	1.38E-17
4.0E+05	8.78E+08	6.84E-18
5.2E+05	1.00E+09	4.82E-18

References: 69, 73, 75, 76, 77, 81, 277, 278, 279

Accuracy: 20%

Notes: (1) Plotted are H^+ energies with the He^+ particles at rest.
(2) The interaction energy is given by $E_{int} = \frac{E_{H^+}}{1.25}$.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 3.3E+03 eV/amu, E_{max} = 5.2E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-80.8329	2.22908	-1.82091	-.0490894	.341964	-.160988	-.0383566	.0746090	-.0322523

The fit represents the above cross sections with an rms deviation of 4.3%.
The maximum deviation is 9.2% at 1.0E+04 eV/amu.
See appendix for Chebyshev fit details.

$$H^+ + He^+ \rightarrow ((H + He^{2+}) + (H^+ + He^{2+} + e^-))$$



Sum of the Electron Capture and Ionization Rate Coefficients for
$H^+ + He^+ \rightarrow [(H + He^{2+}) + (H^+ + He^{2+} + e^-)]$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	Equal Temp.	He^+ Temp. (eV) 500.	750.	1000.	2500.	5000.	10000.	20000.
2.0E+02	5.59E-17	1.11E-15	5.53E-15	1.78E-14	5.49E-13	4.80E-12	4.15E-11	3.10E-10
4.0E+02	3.71E-14	5.08E-14	1.00E-13	1.74E-13	1.35E-12	7.55E-12	5.28E-11	3.42E-10
7.0E+02	7.08E-13	5.49E-13	7.52E-13	9.96E-13	3.49E-12	1.35E-11	7.31E-11	3.93E-10
1.0E+03	2.83E-12	1.92E-12	2.35E-12	2.83E-12	7.15E-12	2.22E-11	9.75E-11	4.46E-10
2.0E+03	3.19E-11	1.82E-11	2.01E-11	2.22E-11	3.77E-11	7.69E-11	2.08E-10	6.41E-10
4.0E+03	2.79E-10	1.62E-10	1.69E-10	1.76E-10	2.25E-10	3.18E-10	5.41E-10	1.08E-09
7.0E+03	1.02E-09	6.66E-10	6.79E-10	6.92E-10	7.72E-10	9.10E-10	1.20E-09	1.78E-09
1.0E+04	1.90E-09	1.34E-09	1.36E-09	1.37E-09	1.46E-09	1.61E-09	1.90E-09	2.47E-09
2.0E+04	4.31E-09	3.51E-09	3.52E-09	3.53E-09	3.60E-09	3.71E-09	3.92E-09	4.31E-09

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 2.0E+02 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He^+ Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
500.	-5.13885E+01	7.38050E+00	-1.17079E+00	6.70967E-02	-1.42287E-01	3.79377E-02	5.35545E-02
750.	-5.05826E+01	6.73738E+00	-8.10583E-01	-8.73094E-02	-9.21944E-02	3.17877E-02	4.71527E-02
1000.	-4.99515E+01	6.24899E+00	-5.58489E-01	-1.82760E-01	-6.61605E-02	3.20490E-02	4.02355E-02
2500.	-4.77567E+01	4.68556E+00	6.56666E-02	-3.34420E-01	-3.90804E-02	4.27066E-02	1.48636E-02
5000.	-4.59364E+01	3.57181E+00	2.68644E-01	-2.79208E-01	-4.43108E-02	3.19685E-02	6.39440E-03
10000.	-4.38282E+01	2.41498E+00	3.24673E-01	-1.50289E-01	-4.79414E-02	9.61266E-03	5.76632E-03
20000.	-4.16749E+01	1.36451E+00	2.92509E-01	-4.41725E-02	-3.60762E-02	-3.52436E-03	2.65593E-03
Equal Temp.	-5.18052E+01	8.42203E+00	-2.10191E+00	5.09552E-01	-3.49726E-01	1.50596E-01	1.11481E-02

See appendix for Chebyshev fit details.

$$H^+ + He^+ \rightarrow ((H + He^{2+}) + (H^+ + He^{2+} + e^-))$$

Maxwellian – Maxwellian



Total Electron Capture Rate Coefficients for
$He^+ + H^+ \rightarrow ((H + He^{2+}) + (H^+ + He^{2+} + e^-))$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	He^+ Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	1.42E-10	1.95E-09	6.39E-09	9.31E-09	9.56E-09	8.57E-09	4.98E-09
2.0E+00	1.43E-10	1.94E-09	6.39E-09	9.31E-09	9.56E-09	8.57E-09	4.98E-09
4.0E+00	1.43E-10	1.94E-09	6.38E-09	9.31E-09	9.55E-09	8.56E-09	4.98E-09
7.0E+00	1.44E-10	1.93E-09	6.37E-09	9.31E-09	9.55E-09	8.56E-09	4.98E-09
1.0E+01	1.45E-10	1.92E-09	6.36E-09	9.30E-09	9.55E-09	8.56E-09	4.98E-09
2.0E+01	1.47E-10	1.91E-09	6.34E-09	9.29E-09	9.55E-09	8.55E-09	4.98E-09
4.0E+01	1.53E-10	1.90E-09	6.31E-09	9.27E-09	9.54E-09	8.55E-09	4.98E-09
7.0E+01	1.60E-10	1.89E-09	6.29E-09	9.25E-09	9.53E-09	8.54E-09	4.94E-09
1.0E+02	1.68E-10	1.89E-09	6.27E-09	9.23E-09	9.53E-09	8.53E-09	4.87E-09
2.0E+02	1.94E-10	1.90E-09	6.22E-09	9.19E-09	9.51E-09	8.52E-09	4.59E-09
4.0E+02	2.53E-10	1.95E-09	6.18E-09	9.13E-09	9.49E-09	8.49E-09	4.20E-09*
7.0E+02	3.50E-10	2.06E-09	6.15E-09	9.07E-09	9.46E-09	8.47E-09	3.88E-09*
1.0E+03	4.49E-10	2.17E-09	6.16E-09	9.02E-09	9.44E-09	8.45E-09	3.69E-09*
2.0E+03	7.69E-10	2.52E-09	6.19E-09	8.90E-09	9.36E-09	8.39E-09	3.40E-09*
4.0E+03	1.37E-09	3.11E-09	6.29E-09	8.70E-09	9.23E-09	8.32E-09	3.18E-09*
7.0E+03	2.18E-09	3.79E-09	6.44E-09	8.49E-09	9.05E-09	8.23E-09	3.05E-09*
1.0E+04	2.88E-09	4.33E-09	6.57E-09	8.34E-09	8.89E-09	8.16E-09	2.99E-09*
2.0E+04	4.61E-09	5.54E-09	6.93E-09	8.06E-09	8.48E-09	7.90E-09	2.90E-09*

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He^+ Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-43.1609	1.70961	.748336	.0779426	-.109478	-.0550219	.00900977
20000.	-39.6126	.447496	.287036	.0983789	-.00499034	-.0233174	-.00975898
40000.	-37.7344	.0172991	.0327260	.0232302	.00718786	6.985132E-05	-.00119392
70000.	-37.0628	-.0628379	-.0317506	-.0103654	-.00206175	4.854519E-04	.00112569
100000.	-36.9842	-.0440273	-.0277168	-.0140155	-.00593012	-.00184524	-1.575483E-04
200000.	-37.1894	-.0312141	-.0170352	-.00770730	-.00339568	-.00158336	-7.883493E-04
500000.	-38.6322	-.298865	-.0952539	.0317736	.0355560	-.00369627	-.0132072

See appendix for Chebyshev fit details

$$He^{+} + H^{+} \rightarrow ((H + He^{2+}) + (H^{+} + He^{2+} + e^{-}))$$

Beam – Maxwellian



Electron Capture Cross Sections for
$H^+ + He^+ \rightarrow H + He^{2+}$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.8E+03	5.89E+07	5.84E-22
2.0E+03	6.21E+07	1.55E-21
4.0E+03	8.79E+07	8.84E-20
7.0E+03	1.16E+08	4.91E-19
1.0E+04	1.39E+08	1.40E-18
2.0E+04	1.96E+08	1.17E-17
4.0E+04	2.78E+08	2.46E-17
4.4E+04	2.91E+08	2.48E-17
5.6E+04	3.29E+08	2.35E-17
7.0E+04	3.68E+08	1.92E-17
1.0E+05	4.39E+08	1.20E-17
2.0E+05	6.21E+08	4.06E-18
2.4E+05	6.80E+08	3.06E-18

References: 69, 71, 73, 74, 75, 76, 77, 78, 79, 80

Accuracy: $E < 1x10^4$ eV/amu - factor 2; $1x10^4$ eV/amu $< E < 2.2x10^5$ eV/amu - 25%

Notes: (1) At energies less than $1x10^4$ eV/amu the assumption was made that the electron capture cross section (σ_c) is equal to the total cross section (σ_T). σ_c: $H^+ + He^+ \rightarrow H + He^{2+}$; σ_T: $H^+ + He^+ \rightarrow [(H + He^+) + (H^+ + He^{2+} + e^-)]$.
(2) Data below $3x10^3$ eV/amu rely on theory of references 71, 79, and 80.
(3) Plotted is the H^+ energy with He^+ at rest.
(4) The interaction energy is given by $E_{int} = \frac{E_{H^+}}{1.25}$.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 1.8E+03 eV/amu, E_{max} = 2.4E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-83.5391	3.94447	-2.70255	.0791317	-.0544615	.290337	-.133801	-.0294272	.0518279

The fit represents the above cross sections with an rms deviation of 2.4%.
The maximum deviation is 3.8% at 4.0E+04 eV/amu.
See appendix for Chebyshev fit details.

$$H^+ + He^+ \rightarrow H + He^{2+}$$



Electron Capture Rate Coefficients for
$H^+ + He^+ \rightarrow H + He^{2+}$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	Equal Temp.	He^+ Temp. (eV) 200.	750.	1000.	2500.	5000.	10000.	20000.
1.0E+02	1.84E-19	2.17E-18	1.20E-15	4.99E-15	3.22E-13	4.38E-12	4.62E-11	3.50E-10
2.0E+02	3.92E-16	3.92E-16	9.86E-15	2.52E-14	5.84E-13	5.73E-12	5.24E-11	3.68E-10
4.0E+02	4.67E-14	2.52E-14	1.13E-13	1.89E-13	1.48E-12	9.25E-12	6.64E-11	4.05E-10
7.0E+02	7.57E-13	3.78E-13	8.05E-13	1.08E-12	4.07E-12	1.69E-11	9.13E-11	4.62E-10
1.0E+03	3.26E-12	1.63E-12	2.67E-12	3.26E-12	8.74E-12	2.81E-11	1.21E-10	5.22E-10
2.0E+03	4.04E-11	2.02E-11	2.54E-11	2.81E-11	4.77E-11	9.59E-11	2.51E-10	7.37E-10
4.0E+03	3.33E-10	1.87E-10	2.06E-10	2.14E-10	2.71E-10	3.77E-10	6.27E-10	1.20E-09
7.0E+03	1.14E-09	7.48E-10	7.79E-10	7.93E-10	8.80E-10	1.03E-09	1.32E-09	1.91E-09
1.0E+04	2.02E-09	1.46E-09	1.49E-09	1.50E-09	1.59E-09	1.74E-09	2.02E-09	2.53E-09
2.0E+04	3.87E-09	3.34E-09	3.35E-09	3.36E-09	3.41E-09	3.49E-09	3.63E-09	3.87E-09

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+02 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He^+ Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
200.	-5.56691E+01	1.03931E+01	-2.17666E+00	2.00410E-01	-1.29725E-01	-1.82527E-02	4.71659E-02
750.	-5.24322E+01	7.77856E+00	-7.52983E-01	-3.38100E-01	-3.27433E-03	-1.92121E-02	3.80958E-02
1000.	-5.15600E+01	7.11555E+00	-4.59118E-01	-4.02692E-01	-1.26409E-02	-3.27140E-03	3.15657E-02
2500.	-4.86136E+01	5.03638E+00	2.33460E-01	-4.36889E-01	-7.67903E-02	3.44015E-02	1.99228E-02
5000.	-4.63267E+01	3.63467E+00	4.33629E-01	-3.18641E-01	-1.01089E-01	2.48311E-02	1.64483E-02
10000.	-4.39403E+01	2.34146E+00	4.31460E-01	-1.58490E-01	-8.81424E-02	-4.91624E-04	1.03754E-02
20000.	-4.16815E+01	1.25619E+00	3.21917E-01	-4.30200E-02	-5.44652E-02	-1.22828E-02	2.56368E-03
Equal Temp.	-5.56573E+01	1.11221E+01	-3.07961E+00	6.63079E-01	-3.65584E-01	9.91144E-02	1.99414E-02

See appendix for Chebyshev fit details.

$H^+ + He^+ \rightarrow H + He^{2+}$

Maxwellian – Maxwellian



Electron Capture Cross Sections for

$H_2^+ + H_2 \rightarrow H_2 + H_2^+$

Energy eV/amu	Cross Section cm^2
2.0E+00	1.22E-15
4.0E+00	1.12E-15
7.0E+00	1.06E-15
1.0E+01	1.01E-15
2.0E+01	9.21E-16
4.0E+01	7.83E-16
8.0E+01	6.06E-16
1.0E+02	5.64E-16
2.0E+02	4.95E-16
3.0E+02	4.89E-16
6.0E+02	4.96E-16
8.0E+02	5.28E-16
1.0E+03	5.39E-16
2.0E+03	5.49E-16
3.0E+03	5.26E-16
5.0E+03	4.81E-16
8.0E+03	3.86E-16
1.0E+04	3.39E-16
2.0E+04	2.04E-16
3.0E+04	1.26E-16
5.0E+04	4.58E-17
6.0E+04	2.72E-17
8.0E+04	1.15E-17
1.0E+05	6.43E-18

$H_2^+ + H \rightarrow H_2 + H^+$

Energy eV/amu	Cross Section cm^2
5.0E+01	8.00E-16
7.0E+01	8.36E-16
1.0E+02	8.75E-16
2.0E+02	9.54E-16
4.0E+02	1.02E-15
7.0E+02	1.07E-15
1.0E+03	1.09E-15
2.0E+03	1.02E-15
3.5E+03	8.96E-16
5.0E+03	7.23E-16

$H_2^+ + He \rightarrow H_2 + He^+$

Energy eV/amu	Cross Section cm^2
1.2E+03	1.39E-18
2.0E+03	1.50E-18
4.0E+03	1.64E-18
7.0E+03	1.73E-18
8.0E+03	1.82E-18
9.0E+03	2.18E-18
1.0E+04	2.68E-18
1.5E+04	5.45E-18
2.0E+04	7.26E-18
2.4E+04	7.49E-18

<u>References:</u> $H_2^+ + H_2$ - 39, 43, 82, 208, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220
$H_2^+ + H$ - 3, 228
$H_2^+ + He$ - 82

<u>Accuracy:</u> $H_2^+ + H_2$ - 30% E < $1x10^3$ eV/amu; 20% E > $1x10^3$ eV/amu
$H_2^+ + H$ - 30%
$H_2^+ + He$ - unknown

<u>Notes:</u> (1) At energies less than $1.5x10^3$ eV/amu the cross sections were determined by measuring the slow ion current produced in the collisions. The cross section was calculated assuming that H_2^+ ions formed in ionizing collisions were negligible. Latimer et al. (Ref. 213) found the ratio of H^+/H_2^+ produced to be .024 at $7.5x10^2$ eV/amu and increased monotonically to .2 at $25x10^3$ eV/amu.

(2) A proportional counter was used in many experiments at energies greater than $1x10^3$ eV/amu to measure H_2 formed in the collision.

(3) It is well known that the H_2^+ electron capture cross sections are dependent on the vibrational levels of the incident H_2^+ ion. See the following set of data for the magnitude of this effect which increases as the energy decreases.

(4) Karpas et al. (Ref. 228) found the rate constants at thermal energies to be $6.4x10^{-10}$ cm^3/s for $H_2^+ + H$ and $5x10^{-10}$ cm^3/s for $D_2^+ + D$.

(5) Electron capture data for H_2^+ in He are very scarce. The data presented here should be used with caution.

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

$H_2^+ + H_2$: E_{min} = 2.0E+00 eV/amu, E_{max} = 1.0E+05 eV/amu
$H_2^+ + H$: E_{min} = 5.0E+01 eV/amu, E_{max} = 5.0E+03 eV/amu
$H_2^+ + He$: E_{min} = 1.2E+03 eV/amu, E_{max} = 2.4E+04 eV/amu

<u>Chebyshev Fitting Parameters for Cross Sections</u>

	A0	A1	A2	A3	A4	A5	A6	A7	A8
$H_2^+ + H_2$:	-71.4572	-1.88878	-.906965	-.676593	-.388666	-.0528444	.0283239	-.0386419	.00767518
$H_2^+ + H$:	-69.2642	.0203046	-.153359	-.0596508	-.0175975	-.00448389	-.00355258	-.00672273	-.00592839
$H_2^+ + He$:	-81.0697	.916248	.310573	.0729101	-.120222	-.148642	.0679087		

The fit represents the $H_2^+ + H_2$ cross section with an rms deviation of 3.8%.
The maximum deviation is 7.1% at 8.1E+04 eV/amu.

The fit represents the $H_2^+ + H$ cross section with an rms deviation of 0.1%.
The maximum deviation is 0.2% at 7.0E+02 eV/amu.

The fit represents the $H_2^+ + He$ cross section with an rms deviation of 2.0%.
The maximum deviation is 4.0% at 8.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$H_2^+ + H, H_2, He \rightarrow H_2$

Cross Section vs. Energy



Electron Capture Cross Sections for

$H_2^+(\nu = 0,1,2,3) + H_2 \rightarrow H_2$

(ν = 0)		(ν = 1)		(ν = 2)		(ν = 3)	
Energy eV/amu	Cross Section cm^2	Energy eV/amu	Cross Section cm^2	Energy eV/amu	Cross Section cm^2	Energy eV/amu	Cross Section cm^2
4.0E+00	7.65E-16	4.0E+00	1.20E-15	4.0E+00	1.22E-15	4.0E+00	1.10E-15
5.0E+00	7.78E-16	6.0E+00	1.15E-15	6.0E+00	1.14E-15	7.5E+00	1.09E-15
6.0E+00	7.90E-16	8.0E+00	1.11E-15	7.5E+00	1.10E-15	1.0E+01	1.05E-15
7.7E+00	8.05E-16	1.5E+01	1.05E-15	1.5E+01	1.04E-15	1.5E+01	9.92E-16
1.0E+01	8.03E-16	2.5E+01	9.97E-16	2.5E+01	9.98E-16	3.0E+01	8.95E-16
1.5E+01	8.04E-16	4.0E+01	9.54E-16	4.0E+01	9.56E-16	6.0E+01	7.98E-16
2.0E+01	8.03E-16	6.0E+01	9.16E-16	7.0E+01	9.11E-16	7.5E+01	7.65E-16
4.0E+01	8.09E-16	9.0E+01	8.72E-16	7.5E+01	9.03E-16	1.5E+02	7.36E-16
6.0E+01	8.07E-16	1.5E+02	8.09E-16	1.0E+02	8.55E-16	2.0E+02	7.20E-16
7.5E+01	8.09E-16	2.0E+02	7.74E-16	2.0E+02	7.45E-16	4.0E+02	6.75E-16
9.0E+01	7.86E-16	3.0E+02	7.62E-16	5.0E+02	5.97E-16	5.0E+02	6.59E-16
1.5E+02	7.19E-16	5.0E+02	7.43E-16				
2.0E+02	6.83E-16						
3.0E+02	6.98E-16						
4.0E+02	7.11E-16						
5.0E+02	7.20E-16						

References: 221, 222, 223, 224, 225, 226, 227, 295

Accuracy: unknown

Note: (1) These experimental data are taken from Campbell et al. (Ref. 221). For further information see theoretical references 222 - 227 and experimental measurements of Liao et al. (Ref. 295).

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

(ν = 0) E_{min} = 4.0E+00 eV/amu, E_{max} = 5.0E+02 eV/amu
(ν = 1) E_{min} = 4.0E+00 eV/amu, E_{max} = 5.0E+02 eV/amu
(ν = 2) E_{min} = 4.0E+00 eV/amu, E_{max} = 5.0E+02 eV/amu
(ν = 3) E_{min} = 4.0E+00 eV/amu, E_{max} = 5.0E+02 eV/amu

Chebyshev Fitting Parameters for Cross Sections

	A0	A1	A2	A3	A4	A5	A6	A7	A8
(ν = 0)	-69.6245	-.0593713	-.0380137	.0269366	.0238283	.0156238	-.00842422	-.0146446	-.00321688
(ν = 1)	-69.2068	-.245756	.00730004	.0108746	.0127040	.00132443	-.00641668	-.00611834	-.00633415
(ν = 2)	-69.2536	-.288634	-.0220048	-.0199150	2.45200E-04	-.0241700	-.0316741	-.0245964	-.0170459
(ν = 3)	-69.3794	-.283633	.0144526	.0213835	-.0200482	-.00269142	-.00368054	.00893694	-5.08689E-04

The fit represents the ν=0 cross section with an rms deviation of 0.6%.
The maximum deviation is 0.9% at 2.0E+02 eV/amu.

The fit represents the ν=1 cross section with an rms deviation of 0.1%.
The maximum deviation is 0.2% at 2.0E+02 eV/amu.

The fit represents the ν=2 cross section with an rms deviation of 0.1%.
The maximum deviation is 0.2% at 4.0E+01 eV/amu.

The fit represents the ν=3 cross section with an rms deviation of 0.3%.
The maximum deviation is 0.7% at 7.5E+01 eV/amu.

See appendix for Chebyshev fit details.

$$H_2^+(\nu=0,1,2,3) + H_2 \rightarrow H_2$$



Electron Capture Cross Sections for
$He^+ + H \rightarrow He + H^+$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.5E+02	2.20E+07	1.54E-17
4.0E+02	2.78E+07	2.39E-17
7.0E+02	3.68E+07	3.99E-17
1.0E+03	4.39E+07	4.74E-17
2.0E+03	6.21E+07	5.57E-17
4.0E+03	8.79E+07	9.08E-17
7.0E+03	1.16E+08	1.71E-16
1.0E+04	1.39E+08	2.33E-16
2.0E+04	1.96E+08	2.42E-16
4.0E+04	2.78E+08	1.00E-16
7.0E+04	3.68E+08	2.97E-17
1.0E+05	4.39E+08	1.29E-17
2.0E+05	6.21E+08	1.01E-18
4.0E+05	8.78E+08	7.94E-20
7.0E+05	1.16E+09	2.46E-21

References: 7, 22, 59, 60, 61

Accuracy: E < 600 eV/amu - unknown; E > 600 eV - 30%

Note: The lowest experimental measurement was at an energy of 500 eV/amu. Data was extrapolated to 250 eV/amu by using the theoretical results of Ref. 61.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 2.5E+02 eV/amu, E_{max} = 7.0E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-79.2449	-3.43445	-3.38701	-.953151	.0800233	.153935	-.198503	-.126958	.0492936

The fit represents the above cross sections with an rms deviation of 6.7%.
The maximum deviation is 15.8% at 2.0E+05 eV/amu.
See appendix for Chebyshev fit details.

$He^+ + H \rightarrow He + H^+$



Electron Capture Rate Coefficients for
$He^+ + H \rightarrow He + H^+$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

He^+ Temp. (eV)	Equal Temp.	H Temp. (eV) 10.	100.	1000.	2500.	5000.	10000.	20000.
7.0E+01	7.00E-11	1.60E-13	1.51E-10	2.97E-09	9.16E-09	1.88E-08	2.75E-08	2.77E-08
1.0E+02	1.74E-10	1.05E-12	1.74E-10	3.00E-09	9.20E-09	1.88E-08	2.75E-08	2.77E-08
2.0E+02	5.93E-10	1.89E-11	2.54E-10	3.08E-09	9.31E-09	1.89E-08	2.75E-08	2.77E-08
4.0E+02	1.37E-09	1.29E-10	4.23E-10	3.25E-09	9.53E-09	1.91E-08	2.75E-08	2.77E-08
7.0E+02	2.50E-09	3.71E-10	6.76E-10	3.52E-09	9.86E-09	1.93E-08	2.76E-08	2.77E-08
1.0E+03	3.79E-09	6.26E-10	9.18E-10	3.79E-09	1.02E-08	1.95E-08	2.77E-08	2.77E-08
2.0E+03	9.08E-09	1.40E-09	1.67E-09	4.76E-09	1.13E-08	2.02E-08	2.78E-08	2.76E-08
4.0E+03	1.88E-08	2.95E-09	3.25E-09	6.87E-09	1.34E-08	2.15E-08	2.81E-08	2.74E-08
7.0E+03	2.63E-08	5.83E-09	6.21E-09	1.02E-08	1.63E-08	2.32E-08	2.84E-08	2.72E-08
1.0E+04	2.86E-08	9.13E-09	9.53E-09	1.34E-08	1.88E-08	2.46E-08	2.86E-08	2.69E-08
2.0E+04	2.59E-08	1.88E-08	1.91E-08	2.15E-08	2.46E-08	2.75E-08	2.88E-08	2.59E-08

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 7.0E+01 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

H Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
10.	-4.44012E+01	5.33316E+00	-1.31634E+00	5.53831E-01	-1.05230E-01	-4.75481E-02	-6.48771E-03
100.	-4.08648E+01	2.48428E+00	2.58050E-01	-3.12250E-02	9.54007E-04	-3.30158E-02	-2.12794E-02
1000.	-3.79160E+01	9.81592E-01	3.54607E-01	2.81517E-02	-3.71282E-02	-1.95845E-02	-3.45087E-03
2500.	-3.63706E+01	4.76956E-01	1.86293E-01	2.58640E-02	-1.33869E-02	-9.40073E-03	-2.40891E-03
5000.	-3.53362E+01	1.79870E-01	7.40800E-02	1.32222E-02	-3.81465E-03	-3.86405E-03	-1.31950E-03
10000.	-3.47840E+01	2.52895E-02	8.72489E-03	-3.06100E-04	-2.09262E-03	-1.25992E-03	-3.74090E-04
20000.	-3.48330E+01	-2.63999E-02	-1.50886E-02	-6.64467E-03	-2.39705E-03	-7.47129E-04	-1.92495E-04
Equal Temp.	-3.92825E+01	2.95989E+00	-6.50284E-01	-6.36867E-02	-1.59695E-01	6.25401E-02	2.55537E-02

See appendix for Chebyshev fit details.

$He^+ + H \rightarrow He + H^+$

Maxwellian — Maxwellian



Electron Capture Rate Coefficients for
$H + He^+ \rightarrow He + H^+$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^+ Temp. (eV)	H Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	3.19E-08	4.74E-08	2.77E-08	1.09E-08	5.66E-09	6.28E-10	1.95E-11**
2.0E+00	3.23E-08	4.74E-08	2.77E-08	1.09E-08	5.66E-09	6.27E-10	1.95E-11**
4.0E+00	3.22E-08	4.73E-08	2.77E-08	1.09E-08	5.65E-09	6.27E-10	1.95E-11**
7.0E+00	3.22E-08	4.72E-08	2.77E-08	1.09E-08	5.65E-09	6.27E-10	1.95E-11**
1.0E+01	3.21E-08	4.71E-08	2.77E-08	1.09E-08	5.65E-09	6.27E-10	1.95E-11**
2.0E+01	3.21E-08	4.70E-08	2.76E-08	1.09E-08	5.64E-09	6.27E-10	1.95E-11**
4.0E+01	3.19E-08	4.68E-08	2.76E-08	1.09E-08	5.63E-09	6.28E-10	1.95E-11**
7.0E+01	3.18E-08	4.65E-08	2.75E-08	1.09E-08	5.62E-09	6.28E-10	1.95E-11**
1.0E+02	3.17E-08	4.63E-08	2.74E-08	1.09E-08	5.61E-09	6.28E-10	1.95E-11**
2.0E+02	3.16E-08	4.58E-08	2.73E-08	1.09E-08	5.59E-09	6.29E-10	1.96E-11**
4.0E+02	3.14E-08	4.51E-08	2.72E-08	1.09E-08	5.56E-09	6.30E-10	1.96E-11**
7.0E+02	3.12E-08	4.44E-08	2.70E-08	1.09E-08	5.54E-09	6.32E-10	1.97E-11**
1.0E+03	3.12E-08	4.38E-08	2.68E-08	1.09E-08	5.52E-09	6.34E-10	1.98E-11*
2.0E+03	3.12E-08	4.24E-08	2.65E-08	1.10E-08	5.50E-09	6.41E-10	2.01E-11*
4.0E+03	3.15E-08	4.04E-08	2.62E-08	1.11E-08	5.51E-09	6.55E-10	2.07E-11*
7.0E+03	3.17E-08	3.82E-08	2.57E-08	1.13E-08	5.56E-09	6.77E-10	2.17E-11*
1.0E+04	3.18E-08	3.65E-08	2.53E-08	1.15E-08	5.63E-09	7.01E-10	2.26E-11*
2.0E+04	3.13E-08	3.26E-08	2.37E-08	1.18E-08	5.92E-09	7.89E-10	2.54E-11*

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

H Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-3.45301E+01	-1.16460E-02	2.95116E-03	6.60447E-03	-4.30159E-03	-3.33439E-03	-4.15571E-03
20000.	-3.39259E+01	-1.54811E-01	-7.89802E-02	-3.03904E-02	-9.78923E-03	-2.83585E-03	-6.35026E-04
40000.	-3.48770E+01	-5.98881E-02	-3.03031E-02	-1.35966E-02	-7.00807E-03	-4.08339E-03	-2.34824E-03
70000.	-3.66351E+01	2.84585E-02	2.06398E-02	1.05841E-02	3.32277E-03	-4.79714E-05	-9.38437E-04
100000.	-3.79874E+01	2.72710E-03	1.55802E-02	1.57294E-02	9.39045E-03	3.99108E-03	1.23431E-03
200000.	-4.22957E+01	7.45856E-02	5.41115E-02	3.23181E-02	1.66050E-02	6.94819E-03	2.78456E-03
500000.	-4.92176E+01	9.14498E-02	6.40861E-02	3.57259E-02	1.57066E-02	5.13868E-03	8.46885E-04

See appendix for Chebyshev fit details.

$$H + He^{+} \rightarrow He + H^{+}$$

Beam – Maxwellian



Electron Capture Cross Sections for
$He^+ + H_2^* \rightarrow He$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.4E-03	6.81E+04	1.65E-18
7.0E-03	1.16E+05	1.20E-18
1.0E-02	1.39E+05	1.07E-18
4.0E-02	2.78E+05	6.66E-19
1.0E-01	4.39E+05	4.35E-19
4.0E-01	8.79E+05	1.89E-19
7.0E-01	1.16E+06	9.93E-20
1.0E+00	1.39E+06	5.21E-20
1.3E+00	1.58E+06	4.40E-20
1.5E+00	1.70E+06	5.70E-20
3.0E+00	2.41E+06	2.15E-19
4.0E+00	2.78E+06	2.41E-19
7.0E+00	3.68E+06	2.65E-19
1.0E+01	4.39E+06	2.77E-19
4.0E+01	8.79E+06	5.03E-19
7.0E+01	1.16E+07	1.13E-18
1.0E+02	1.39E+07	1.88E-18
4.0E+02	2.78E+07	1.18E-17
7.0E+02	3.68E+07	2.46E-17
1.0E+03	4.39E+07	3.77E-17
4.0E+03	8.79E+07	1.42E-16
7.0E+03	1.16E+08	2.09E-16
1.0E+04	1.39E+08	2.42E-16
1.6E+04	1.76E+08	2.69E-16
4.0E+04	2.78E+08	1.54E-16
7.0E+04	3.70E+08	6.41E-17
1.0E+05	4.39E+08	2.84E-17
4.0E+05	8.78E+08	2.15E-19
7.0E+05	1.16E+09	1.35E-20

References: 7, 22, 82, 91, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107

Accuracy: $E < 2x10^3$ eV/amu - unknown; $E > 2x10^3$ eV/amu - 20%

Notes: (1) The data between 40 - 2000 eV/amu have been interpolated
(2) The data of Moran and Conrads (Ref. 107) are essentially independent of energy in the 250 - 800 eV/amu region and are approximately a factor of 10 above the values shown.
(3) The low energy data ($E < 2x10^3$ eV/amu) of Stedeford and Hasted (Ref. 82) are 1 to 2 orders of magnitude greater than the extrapolated data.
(4) The data below $2x10^3$ eV/amu have large uncertainties
(5) In the experiments cited the He^+ metastable state population is dependent on the ion source used and the operating parameters of the source. As the metastable population increases the total cross section increases.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 2.4E-03 eV/amu, E_{max} = 7.0E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-81.9486	.720256	-.856412	-3.13728	-1.68226	.0267213	.0474764	3.15823E-03	.164787

The fit represents the above cross sections with an rms deviation of 34.8%.
The maximum deviation is 132.4% at 1.3E+00 eV/amu.
See appendix for Chebyshev fit details.

$He^+ + H_2 \rightarrow He$



Electron Capture Cross Sections for
$He^+ + He \rightarrow He + He^+$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
4.0E-02	2.78E+05	3.32E-15
7.0E-02	3.68E+05	3.12E-15
1.0E-01	4.39E+05	2.97E-15
2.0E-01	6.21E+05	2.85E-15
4.0E-01	8.79E+05	2.64E-15
7.0E-01	1.16E+06	2.43E-15
1.0E+00	1.39E+06	2.31E-15
2.0E+00	1.96E+06	2.10E-15
4.0E+00	2.78E+06	1.96E-15
7.0E+00	3.68E+06	1.83E-15
1.0E+01	4.39E+06	1.75E-15
2.0E+01	6.21E+06	1.54E-15
4.0E+01	8.79E+06	1.39E-15
7.0E+01	1.16E+07	1.31E-15
1.0E+02	1.39E+07	1.26E-15
2.0E+02	1.96E+07	1.11E-15
4.0E+02	2.78E+07	9.87E-16
7.0E+02	3.68E+07	8.56E-16
1.0E+03	4.39E+07	7.92E-16
2.0E+03	6.21E+07	6.48E-16
4.0E+03	8.79E+07	5.26E-16
7.0E+03	1.16E+08	4.22E-16
1.0E+04	1.39E+08	3.60E-16
2.0E+04	1.96E+08	2.36E-16
4.0E+04	2.78E+08	1.28E-16
7.0E+04	3.68E+08	5.58E-17
1.0E+05	4.39E+08	2.68E-17
2.0E+05	6.21E+08	5.49E-18
4.0E+05	8.78E+08	4.12E-19
5.3E+05	1.01E+09	1.08E-19

References: 91, 96, 97, 98, 99, 103, 104, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 390, 397

Accuracy: 20% throughout the energy range

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 4.0E-02 eV/amu, E_{max} = 5.3E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-71.6915	-3.72702	-1.97026	-1.16040	-.605858	-.259050	-.0892322	-.0165873	9.75783E-03

The fit represents the above cross sections with an rms deviation of 2.4%.
The maximum deviation is 9.2% at 1.0E+05 eV/amu.
See appendix for Chebyshev fit details.

$He^+ + He \rightarrow He + He^+$



Electron Capture Rate Coefficients for
$He^+ + He \rightarrow He + He^+$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

He^+ Temp. (eV)	Equal Temp.	He Temp. (eV) 1.	100.	500.	1000.	5000.	10000.	20000.
1.0E+00	2.66E-09	2.66E-09	1.12E-08	1.97E-08	2.44E-08	3.68E-08	4.18E-08	4.51E-08
2.0E+00	3.45E-09	3.10E-09	1.13E-08	1.97E-08	2.44E-08	3.68E-08	4.18E-08	4.51E-08
4.0E+00	4.49E-09	3.75E-09	1.13E-08	1.97E-08	2.44E-08	3.68E-08	4.18E-08	4.51E-08
7.0E+00	5.54E-09	4.49E-09	1.15E-08	1.97E-08	2.44E-08	3.68E-08	4.18E-08	4.51E-08
1.0E+01	6.30E-09	5.06E-09	1.16E-08	1.98E-08	2.44E-08	3.68E-08	4.18E-08	4.51E-08
2.0E+01	8.06E-09	6.42E-09	1.19E-08	1.99E-08	2.45E-08	3.68E-08	4.18E-08	4.51E-08
4.0E+01	1.03E-08	8.13E-09	1.26E-08	2.01E-08	2.47E-08	3.68E-08	4.18E-08	4.51E-08
7.0E+01	1.26E-08	9.87E-09	1.35E-08	2.05E-08	2.49E-08	3.69E-08	4.18E-08	4.51E-08
1.0E+02	1.43E-08	1.12E-08	1.43E-08	2.08E-08	2.51E-08	3.69E-08	4.18E-08	4.51E-08
2.0E+02	1.82E-08	1.44E-08	1.65E-08	2.19E-08	2.57E-08	3.71E-08	4.19E-08	4.51E-08
4.0E+02	2.28E-08	1.83E-08	1.96E-08	2.36E-08	2.69E-08	3.74E-08	4.20E-08	4.51E-08
7.0E+02	2.69E-08	2.19E-08	2.28E-08	2.57E-08	2.83E-08	3.78E-08	4.22E-08	4.52E-08
1.0E+03	2.96E-08	2.44E-08	2.51E-08	2.74E-08	2.96E-08	3.82E-08	4.24E-08	4.52E-08
2.0E+03	3.50E-08	2.96E-08	2.99E-08	3.13E-08	3.27E-08	3.93E-08	4.29E-08	4.53E-08
4.0E+03	4.03E-08	3.50E-08	3.52E-08	3.59E-08	3.68E-08	4.11E-08	4.37E-08	4.54E-08
7.0E+03	4.37E-08	3.93E-08	3.94E-08	3.98E-08	4.03E-08	4.29E-08	4.46E-08	4.55E-08
1.0E+04	4.51E-08	4.18E-08	4.18E-08	4.21E-08	4.24E-08	4.40E-08	4.51E-08	4.54E-08
2.0E+04	4.47E-08	4.51E-08	4.51E-08	4.51E-08	4.52E-08	4.55E-08	4.54E-08	4.47E-08

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
1.	-3.65091E+01	1.50347E+00	-7.73477E-02	-8.34424E-02	-7.59297E-04	-3.17776E-03	2.46929E-03
100.	-3.55092E+01	7.61944E-01	1.84506E-01	-6.99606E-02	-4.58282E-02	5.01278E-03	6.02365E-03
500.	-3.49239E+01	4.30514E-01	1.66682E-01	-1.76952E-03	-3.51444E-02	-1.39113E-02	1.27578E-03
1000.	-3.46672E+01	3.07854E-01	1.38199E-01	1.53048E-02	-2.27026E-02	-1.52299E-02	-2.87864E-03
5000.	-3.41193E+01	9.72499E-02	5.39119E-02	1.55767E-02	-2.97046E-03	-6.16891E-03	-3.80972E-03
10000.	-3.39337E+01	3.94479E-02	2.21217E-02	6.22476E-03	-1.76048E-03	-3.26193E-03	-2.26998E-03
20000.	-3.38270E+01	1.35988E-03	-9.13724E-04	-2.77809E-03	-3.01978E-03	-2.17733E-03	-1.25142E-03
Equal Temp.	-3.62362E+01	1.48422E+00	-1.94700E-01	-6.90443E-02	-1.94257E-02	-1.35547E-03	-2.79202E-03

See appendix for Chebyshev fit details.

$He^+ + He \rightarrow He + He^+$

Maxwellian – Maxwellian



Electron Capture Rate Coefficients for
$He + He^+ \rightarrow He + He^+$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^+ Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	400000.
1.0E+00	5.00E-08	4.63E-08	3.55E-08	2.05E-08	1.18E-08	3.41E-09*	3.61E-10**
2.0E+00	5.00E-08	4.63E-08	3.55E-08	2.05E-08	1.18E-08	3.40E-09*	3.62E-10**
4.0E+00	5.00E-08	4.63E-08	3.55E-08	2.05E-08	1.18E-08	3.40E-09*	3.61E-10**
7.0E+00	4.99E-08	4.63E-08	3.55E-08	2.05E-08	1.18E-08	3.40E-09*	3.61E-10**
1.0E+01	4.99E-08	4.63E-08	3.55E-08	2.05E-08	1.18E-08	3.40E-09*	3.61E-10**
2.0E+01	4.99E-08	4.62E-08	3.54E-08	2.05E-08	1.18E-08	3.40E-09*	3.61E-10**
4.0E+01	4.99E-08	4.62E-08	3.54E-08	2.04E-08	1.18E-08	3.39E-09*	3.61E-10**
7.0E+01	4.98E-08	4.61E-08	3.53E-08	2.04E-08	1.18E-08	3.39E-09*	3.61E-10**
1.0E+02	4.98E-08	4.61E-08	3.53E-08	2.04E-08	1.18E-08	3.38E-09*	3.60E-10**
2.0E+02	4.97E-08	4.60E-08	3.51E-08	2.03E-08	1.17E-08	3.37E-09*	3.60E-10**
4.0E+02	4.95E-08	4.58E-08	3.50E-08	2.03E-08	1.17E-08	3.36E-09*	3.60E-10**
7.0E+02	4.94E-08	4.57E-08	3.48E-08	2.02E-08	1.17E-08	3.34E-09*	3.59E-10**
1.0E+03	4.93E-08	4.55E-08	3.47E-08	2.02E-08	1.18E-08	3.33E-09	3.59E-10**
2.0E+03	4.89E-08	4.52E-08	3.43E-08	2.01E-08	1.18E-08	3.31E-09	3.60E-10**
4.0E+03	4.83E-08	4.46E-08	3.38E-08	2.00E-08	1.18E-08	3.30E-09	3.62E-10**
7.0E+03	4.75E-08	4.39E-08	3.32E-08	1.99E-08	1.19E-08	3.29E-09	3.67E-10**
1.0E+04	4.68E-08	4.31E-08	3.28E-08	1.98E-08	1.20E-08	3.30E-09	3.72E-10**
2.0E+04	4.48E-08	4.09E-08	3.13E-08	1.96E-08	1.22E-08	3.37E-09	3.91E-10*

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-3.36725E+01	-4.07805E-02	-2.43723E-02	-1.21532E-02	-5.34881E-03	-2.01573E-03	-5.49929E-04
20000.	-3.38300E+01	-4.55632E-02	-2.69923E-02	-1.37596E-02	-6.60783E-03	-3.00454E-03	-1.25225E-03
40000.	-3.43690E+01	-5.00191E-02	-2.61460E-02	-1.10247E-02	-4.45099E-03	-1.92372E-03	-9.15345E-04
70000.	-3.54336E+01	-2.00865E-02	-7.82001E-03	-2.03461E-03	-7.58265E-04	-7.02792E-04	-6.07996E-04
100000.	-3.65032E+01	1.22086E-02	1.00798E-02	5.91914E-03	2.15183E-03	3.37849E-04	-2.72081E-04
200000.	-3.90194E+01	-1.48944E-02	8.12695E-04	5.76317E-03	5.00317E-03	2.19384E-03	1.07961E-03
400000.	-4.34614E+01	2.08264E-02	2.00251E-02	1.46660E-02	7.48524E-03	3.55071E-03	7.16357E-04

See appendix for Chebyshev fit details.

$He + He^+ \rightarrow He + He^+$

Beam – Maxwellian



Electron Capture Cross Sections for
$He^+ + Li \rightarrow He + Li^+$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
6.4E+01	1.11E+07	7.80E-16
1.0E+02	1.39E+07	1.53E-15
2.0E+02	1.96E+07	3.27E-15
5.0E+02	3.11E+07	5.25E-15
8.0E+02	3.93E+07	5.61E-15
1.0E+03	4.39E+07	5.60E-15
2.0E+03	6.21E+07	5.12E-15
4.0E+03	8.79E+07	4.17E-15
7.0E+03	1.16E+08	2.93E-15
1.0E+04	1.39E+08	1.76E-15
2.0E+04	1.96E+08	3.26E-16
2.5E+04	2.20E+08	1.33E-16
3.0E+04	2.41E+08	6.37E-17
4.0E+04	2.78E+08	4.17E-17
5.0E+04	3.11E+08	3.42E-17
7.0E+04	3.68E+08	2.60E-17
1.0E+05	4.39E+08	2.00E-17

References: 230, 231, 232, 233, 234, 242

Accuracy: 30%

Note: For electron capture cross sections of Li^+ and Li^{2+} in He see references 249, 250, 251, 254.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 6.4E+01 eV/amu, E_{max} = 1.0E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-69.9420	-2.17479	-1.90237	.153408	.309985	.316010	.0848497	-.118385	-.140948

The fit represents the above cross sections with an rms deviation of 8.8%.
The maximum deviation is 26.5% at 3.0E+04 eV/amu.
See appendix for Chebyshev fit details.

$He^+ + Li \rightarrow He + Li^+$

Cross Section vs. Energy



Electron Capture Rate Coefficients for
$Li + He^+ \rightarrow He + Li^+$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^+ Temp. (eV)	Li Energy (eV/amu) 10000.	20000.	30000.	40000.	60000.	80000.	100000.
1.0E+00	2.44E-07	6.38E-08*	1.54E-08**	1.16E-08**	1.00E-08**	9.26E-09**	4.39E-09**
2.0E+00	2.44E-07	6.38E-08*	1.54E-08**	1.16E-08**	1.00E-08**	9.26E-09**	4.39E-09**
4.0E+00	2.43E-07	6.37E-08*	1.55E-08**	1.16E-08**	1.00E-08**	9.26E-09**	4.39E-09**
7.0E+00	2.43E-07	6.35E-08*	1.55E-08**	1.16E-08**	1.00E-08**	9.26E-09**	4.39E-09**
1.0E+01	2.42E-07	6.35E-08*	1.55E-08**	1.16E-08**	1.00E-08**	9.26E-09**	4.39E-09**
2.0E+01	2.42E-07	6.32E-08*	1.56E-08**	1.16E-08**	1.00E-08**	9.26E-09**	4.38E-09**
4.0E+01	2.40E-07	6.30E-08*	1.58E-08**	1.16E-08**	1.00E-08**	9.26E-09**	4.38E-09**
7.0E+01	2.39E-07	6.27E-08*	1.59E-08**	1.17E-08**	1.00E-08**	9.26E-09**	4.38E-09**
1.0E+02	2.38E-07	6.25E-08*	1.60E-08**	1.17E-08**	1.00E-08**	9.26E-09**	4.37E-09**
2.0E+02	2.36E-07	6.22E-08*	1.64E-08**	1.17E-08**	1.00E-08**	9.26E-09**	4.36E-09**
4.0E+02	2.34E-07	6.19E-08*	1.70E-08**	1.18E-08**	1.00E-08**	9.26E-09**	4.35E-09**
7.0E+02	2.31E-07	6.19E-08*	1.77E-08*	1.19E-08**	1.00E-08**	9.26E-09**	4.34E-09**
1.0E+03	2.29E-07	6.22E-08*	1.83E-08*	1.19E-08**	1.00E-08**	9.24E-09**	4.33E-09**
2.0E+03	2.21E-07	6.38E-08*	2.03E-08*	1.22E-08**	1.00E-08**	9.09E-09**	4.30E-09**
4.0E+03	2.09E-07	6.77E-08*	2.36E-08*	1.31E-08**	1.01E-08**	8.59E-09**	4.26E-09**
7.0E+03	1.93E-07	7.22E-08	2.78E-08*	1.48E-08*	1.01E-08**	7.97E-09**	4.22E-09**
1.0E+04	1.80E-07	7.48E-08	3.14E-08*	1.66E-08*	1.01E-08**	7.54E-09**	4.18E-09**
2.0E+04	1.50E-07	7.63E-08	3.93E-08*	2.22E-08*	1.08E-08**	6.91E-09**	4.13E-09**

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

Li Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-3.06784E+01	-1.83367E-01	-1.08562E-01	-5.42037E-02	-2.28601E-02	-6.74194E-03	-2.90887E-04
20000.	-3.30630E+01	7.50860E-02	6.65545E-02	2.87114E-02	-2.24836E-03	-1.23958E-02	-1.01177E-02
30000.	-3.54800E+01	4.01173E-01	2.16479E-01	7.71491E-02	1.07563E-02	-8.06147E-03	-8.04622E-03
40000.	-3.62976E+01	2.18893E-01	1.57718E-01	9.39441E-02	4.47005E-02	1.42990E-02	-8.03805E-04
60000.	-3.68192E+01	1.63206E-02	1.40921E-02	1.21589E-02	9.29880E-03	6.22735E-03	4.24079E-03
80000.	-3.71186E+01	-1.09657E-01	-8.05414E-02	-4.26549E-02	-1.26465E-02	3.18261E-03	7.94923E-03
100000.	-3.85227E+01	-2.71037E-02	-1.31775E-02	-4.13455E-03	-5.05274E-04	3.90860E-04	4.43525E-04

See appendix for Chebyshev fit details.

$Li + He^+ \rightarrow He + Li^+$

Beam – Maxwellian



Electron Capture Cross Sections for
$He^+ + He^+ \rightarrow He + He^{2+}$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.5E+04	1.70E+08	1.95E-18
2.0E+04	1.96E+08	3.25E-18
3.0E+04	2.41E+08	6.28E-18
4.0E+04	2.78E+08	9.44E-18
5.0E+04	3.11E+08	1.22E-17
6.0E+04	3.40E+08	1.45E-17
7.0E+04	3.68E+08	1.64E-17
9.0E+04	4.17E+08	1.88E-17
1.0E+05	4.39E+08	1.90E-17
1.5E+05	5.38E+08	1.73E-17
2.0E+05	6.21E+08	1.42E-17
2.3E+05	6.66E+08	1.24E-17

References: 69, 277, 278, 283, 284

Accuracy: 20%

Note: (1) Good agreement of the experimental data exists with the theoretical data of Ref. 284.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 1.5E+04 eV/amu, E_{max} = 2.3E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-78.6559	.981266	-.533866	-.0667013	.0133417	.00898626	.00154609	.00128923	-.00693562

The fit represents the above cross sections with an rms deviation of 0.3%.
The maximum deviation is 0.6% at 1.0E+05 eV/amu.
See appendix for Chebyshev fit details.

$He^{+} + He^{+} \rightarrow He + He^{2+}$



Electron Capture Rate Coefficients for
$He^+ + He^+ \rightarrow He + He^{2+}$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

He^+ Temp. (eV)	Equal Temp.	He^+ Temp. (eV) 10000.	12000.	14000.	16000.	18000.	20000.	25000.
4.0E+03	8.29E-13	2.06E-11	3.60E-11	5.61E-11	8.08E-11	1.10E-10	1.43E-10	2.40E-10
6.0E+03	9.94E-12	3.60E-11	5.61E-11	8.08E-11	1.10E-10	1.43E-10	1.79E-10	2.85E-10
8.0E+03	3.60E-11	5.61E-11	8.08E-11	1.10E-10	1.43E-10	1.79E-10	2.19E-10	3.32E-10
1.0E+04	8.08E-11	8.08E-11	1.10E-10	1.43E-10	1.79E-10	2.19E-10	2.62E-10	3.81E-10
1.2E+04	1.43E-10	1.10E-10	1.43E-10	1.79E-10	2.19E-10	2.62E-10	3.08E-10	4.33E-10
1.4E+04	2.19E-10	1.43E-10	1.79E-10	2.19E-10	2.62E-10	3.08E-10	3.56E-10	4.87E-10
1.6E+04	3.08E-10	1.79E-10	2.19E-10	2.62E-10	3.08E-10	3.56E-10	4.07E-10	5.42E-10
1.8E+04	4.07E-10	2.19E-10	2.62E-10	3.08E-10	3.56E-10	4.07E-10	4.60E-10	5.99E-10
2.0E+04	5.14E-10	2.62E-10	3.08E-10	3.56E-10	4.07E-10	4.60E-10	5.14E-10	6.58E-10
2.2E+04	6.28E-10	3.08E-10	3.56E-10	4.07E-10	4.60E-10	5.14E-10	5.71E-10	7.18E-10
2.4E+04	7.49E-10	3.56E-10	4.07E-10	4.60E-10	5.14E-10	5.71E-10	6.28E-10	7.80E-10

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 4.0E+03 eV, E_{max} = 2.4E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He^+ Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
10000.	-4.64541E+01	1.43962E+00	4.63030E-02	-1.49060E-02	-2.30149E-04	2.44855E-04	-4.28539E-05
12000.	-4.57963E+01	1.22412E+00	6.31757E-02	-1.13559E-02	-6.21329E-04	1.52187E-04	1.97060E-04
14000.	-4.52461E+01	1.05998E+00	7.17125E-02	-8.34828E-03	-9.77820E-04	2.30920E-04	-9.51457E-06
16000.	-4.47759E+01	9.31470E-01	7.53759E-02	-5.93773E-03	-1.18832E-03	1.95011E-04	-1.89337E-04
18000.	-4.43672E+01	8.28548E-01	7.61027E-02	-4.06263E-03	-1.22968E-03	-1.52699E-05	-1.37761E-04
20000.	-4.40069E+01	7.44390E-01	7.51176E-02	-2.63629E-03	-1.17745E-03	-2.47636E-04	7.54863E-05
25000.	-4.32601E+01	5.88552E-01	7.03846E-02	-1.20370E-04	-1.13108E-03	1.84062E-04	-1.45794E-04
Equal Temp.	-4.76973E+01	3.30679E+00	-5.53811E-01	9.53225E-02	-1.30960E-02	6.76109E-04	1.60674E-04

See appendix for Chebyshev fit details.

$He^+ + He^+ \rightarrow He + He^{2+}$

Maxwellian – Maxwellian



Partial and Total Electron Capture Cross Sections for

$He^{2+} + He^{+} \rightarrow He^{+} + He^{2+}$

(Partial)

He^{+} Energy eV/amu	Cross Section cm^2
7.0E+00	3.04E-17
1.0E+01	4.02E-17
1.5E+01	7.57E-17
2.0E+01	2.00E-16
2.5E+01	2.98E-16
3.0E+01	3.48E-16
4.0E+01	4.04E-16
6.0E+01	4.45E-16
7.0E+01	4.55E-16
1.0E+02	4.79E-16
2.0E+02	4.84E-16
4.0E+02	4.64E-16
7.0E+02	4.38E-16
1.0E+03	4.19E-16
2.0E+03	3.76E-16
4.0E+03	3.28E-16
7.0E+03	2.83E-16
1.0E+04	2.52E-16
1.6E+04	2.07E-16

(Total)

He^{+} Energy eV/amu	Cross Section cm^2
8.5E+00	3.39E-16
1.0E+01	4.09E-16
1.5E+01	6.01E-16
2.0E+01	6.99E-16
4.0E+01	7.77E-16
7.0E+01	7.80E-16
1.0E+02	7.66E-16
2.0E+02	7.16E-16
4.0E+02	6.64E-16
7.0E+02	6.13E-16
1.0E+03	5.75E-16
2.0E+03	5.01E-16
4.0E+03	4.30E-16
7.0E+03	3.76E-16
1.0E+04	3.23E-16
2.0E+04	2.03E-16
4.0E+04	1.01E-16
7.0E+04	4.50E-17
1.0E+05	2.65E-17
1.5E+05	1.33E-17
2.0E+05	6.34E-18
2.5E+05	2.83E-18
3.0E+05	1.27E-18

References: 69, 288, 391, 392, 393

Accuracy: unknown

Notes: (1) In the measurements a problem arises from the mutual repulsion between the He^{+} and He^{2+}. The repulsion results in scattering with some of the ions not reaching the detector which has a finite acceptance angle. These measurements are termed partial cross sections.
(2) The total cross section data are from theoretical values in Ref. 288 and 391. The theoretical data has been joined with a smooth curve between 6×10^3 and 10^5 eV/amu.
(3) For this simple one electron, resonant system, the theoretical data should be accurate to within 20%.

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

(Partial) E_{min} = 7.1E+00 eV/amu, E_{max} = 1.6E+04 eV/amu
(Total) E_{min} = 8.5E+00 eV/amu, E_{max} = 3.0E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

	A0	A1	A2	A3	A4	A5	A6	A7	A8
(Partial)	-72.2134	.725003	-1.00558	.400148	-.150634	-.0556029	.119500	-.118086	.0830868
(Total)	-72.8308	-2.29484	-1.54555	-.441776	-.328931	-.00327441	-.0709033	-.0333345	-.0595299

The fit represents the partial cross section with an rms deviation of 7.4%.
The maximum deviation is 25.2% at 1.5E+01 eV/amu.

The fit represents the total cross section with an rms deviation of 2.8%.
The maximum deviation is 5.6% at 7.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$He^{2+} + He^{+} \rightarrow He^{+} + He^{2+}$



Single Electron Capture Cross Sections for

$He^{2+} + H^- \rightarrow He^+ + H$
$He^+ + H^- \rightarrow He + H$

$He^{2+} + H^-$ Energy eV/amu	Cross Section cm^2
4.7E-01	1.82E-13
5.0E-01	1.71E-13
6.0E-01	1.52E-13
8.0E-01	1.30E-13
1.0E+00	1.17E-13
2.0E+00	8.90E-14
3.0E+00	7.78E-14
4.0E+00	7.13E-14
6.0E+00	6.44E-14
8.0E+00	6.01E-14
1.0E+01	5.79E-14
2.0E+01	5.31E-14
3.0E+01	5.08E-14
5.0E+01	4.77E-14
8.0E+01	4.55E-14
1.0E+02	4.43E-14
2.0E+02	4.12E-14
4.0E+02	3.73E-14
7.0E+02	3.44E-14
1.0E+03	3.22E-14
2.0E+03	2.79E-14
3.0E+03	2.50E-14

$He^+ + H^-$ Energy eV/amu	Cross Section cm^2
4.2E+01	3.03E-14
5.0E+01	2.18E-14
6.0E+01	1.83E-14
7.0E+01	1.66E-14
8.0E+01	1.56E-14
1.0E+02	1.46E-14
1.5E+02	1.40E-14
2.0E+02	1.37E-14
3.0E+02	1.32E-14
4.0E+02	1.30E-14
5.0E+02	1.28E-14
7.0E+02	1.24E-14
1.0E+03	1.19E-14
2.0E+03	9.78E-15
3.0E+03	7.89E-15
4.0E+03	6.06E-15
5.0E+03	4.40E-15
7.0E+03	2.43E-15
7.7E+03	2.01E-15

References: ($He^+ + H^-$) 68, 292, 293, 394 ; ($He^{2+} + H^-$) 69, 290, 291, 294

Accuracy: ($He^+ + H^-$) - 20%; ($He^{2+} + H^-$) - 40%

Notes: (1) In the energy region of overlap, the $He^{2+} + H^-$ cross sections of Terao et al (ref. 290, 291) are consistently greater than those of the Dolder and Peart group (ref. 69, 294).
(2) The measurements indicate some structure in the cross sections for both the He^+ and He^{2+} reactions. This oscillation has been smoothed out in the data presented here.

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

($He^{2+} + H^-$) E_{min} = 4.7E-01 eV/amu, E_{max} = 3.0E+03 eV/amu
($He^+ + H^-$) E_{min} = 4.2E+01 eV/amu, E_{max} = 7.7E+03 eV/amu

Chebyshev Fitting Parameters for Cross Sections

	A0	A1	A2	A3	A4	A5	A6	A7	A8
($He^{2+} + H^-$)	-61.0067	-.855510	.155087	-.126012	.0124474	-6.60133E-04	-4.30771E-04	-.00636246	.00615254
($He^+ + H^-$)	-64.4915	-.959186	-.269411	-.331423	-.00745857	-.0611579	.0257220	-.00396428	.0113568

The fit represents the $He^{2+} + H^-$ cross section with an rms deviation of 0.4%.
The maximum deviation is 0.8% at 5.0E-01 eV/amu.

The fit represents the $He^+ + H^-$ cross section with an rms deviation of 0.6%.
The maximum deviation is 1.2% at 1.0E+02 eV/amu.

See appendix for Chebyshev fit details.

$He^{q+} + H^- \rightarrow He^{(q-1)+} + H \ (q=1,2)$

Cross Section vs. Energy



Electron Capture Cross Sections for
$He^{2+} + H \rightarrow He^{+} + H^{+}$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
7.0E+01	1.16E+07	1.25E-20
1.0E+02	1.39E+07	2.20E-19
2.0E+02	1.96E+07	3.10E-18
4.0E+02	2.78E+07	1.90E-17
5.0E+02	3.11E+07	3.24E-17
7.0E+02	3.68E+07	8.67E-17
1.0E+03	4.39E+07	1.87E-16
2.0E+03	6.21E+07	5.96E-16
4.0E+03	8.79E+07	9.74E-16
7.0E+03	1.16E+08	1.18E-15
1.0E+04	1.39E+08	1.22E-15
2.0E+04	1.96E+08	1.09E-15
4.0E+04	2.78E+08	5.28E-16
7.0E+04	3.68E+08	1.59E-16
1.0E+05	4.39E+08	5.23E-17
2.0E+05	6.21E+08	4.63E-18
4.0E+05	8.78E+08	2.64E-19
5.0E+05	9.82E+08	1.15E-19

References: 5, 7, 21, 22, 23, 24, 25, 57, 58

Accuracy: He^{2+} E < 600 eV/amu - unknown; He^{2+} E > 600 eV/amu - 20%

Notes: (1) At energies less than 2 keV/amu the results of Fite et al. (Ref. 5) and Nutt et al. (Ref. 23) diverge by as much as a factor of 10. This is believed to be due to the fact that the results reported by Fite et al. were obtained by measuring the slow ions produced in the collision, and are less reliable than those of Nutt et al at low energies.
(2) No experimental data exist below 500 eV/amu. Cross sections below 500 eV/amu were taken from calculations of Ref. 57 and 58 using multi-state close coupling methods.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 7.0E+01 eV/amu, E_{max} = 5.0E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-78.3976	.773499	-5.12346	-.125575	-.232331	.261317	-.0479652	.189434	-.138708

The fit represents the above cross sections with an rms deviation of 3.9%.
The maximum deviation is 7.6% at 5.0E+02 eV/amu.
See appendix for Chebyshev fit details.

$He^{2+} + H \rightarrow He^{+} + H^{+}$



Electron Capture Rate Coefficients for
$He^{2+} + H \rightarrow He^{+} + H^{+}$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	Equal Temp.	H Temp. (eV) 10.	100.	500.	1000.	5000.	10000.	20000.
2.0E+01	6.28E-13	4.12E-14	1.22E-10	7.36E-09	2.36E-08	1.14E-07	1.47E-07	1.42E-07
4.0E+01	1.04E-11	2.09E-13	1.42E-10	7.50E-09	2.38E-08	1.14E-07	1.47E-07	1.42E-07
7.0E+01	6.86E-11	9.71E-13	1.74E-10	7.72E-09	2.40E-08	1.14E-07	1.47E-07	1.42E-07
1.0E+02	2.10E-10	2.70E-12	2.10E-10	7.94E-09	2.43E-08	1.14E-07	1.47E-07	1.42E-07
2.0E+02	1.49E-09	1.97E-11	3.63E-10	8.68E-09	2.51E-08	1.14E-07	1.47E-07	1.42E-07
4.0E+02	7.21E-09	1.42E-10	8.23E-10	1.02E-08	2.68E-08	1.15E-07	1.47E-07	1.42E-07
7.0E+02	1.92E-08	6.63E-10	1.89E-09	1.26E-08	2.93E-08	1.16E-07	1.47E-07	1.42E-07
1.0E+03	3.18E-08	1.65E-09	3.37E-09	1.50E-08	3.18E-08	1.17E-07	1.47E-07	1.42E-07
2.0E+03	6.80E-08	7.50E-09	1.02E-08	2.34E-08	3.99E-08	1.19E-07	1.48E-07	1.41E-07
4.0E+03	1.13E-07	2.38E-08	2.68E-08	3.99E-08	5.48E-08	1.25E-07	1.49E-07	1.40E-07
7.0E+03	1.43E-07	4.78E-08	5.05E-08	6.16E-08	7.40E-08	1.31E-07	1.49E-07	1.39E-07
1.0E+04	1.50E-07	6.83E-08	7.05E-08	7.97E-08	8.98E-08	1.36E-07	1.50E-07	1.37E-07
2.0E+04	1.32E-07	1.14E-07	1.15E-07	1.19E-07	1.25E-07	1.47E-07	1.49E-07	1.32E-07

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 2.0E+01 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

H Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
10.	-4.49742E+01	7.78326E+00	-1.00928E+00	-3.95746E-01	8.29111E-02	3.18952E-02	5.49793E-03
100.	-3.94867E+01	3.76780E+00	4.07631E-01	-4.05432E-01	-7.75944E-02	6.09541E-02	1.07817E-02
500.	-3.54181E+01	1.44543E+00	4.43041E-01	-3.74033E-02	-7.53578E-02	-1.52405E-02	7.25877E-03
1000.	-3.40218E+01	8.22262E-01	3.19564E-01	2.78648E-02	-3.66721E-02	-1.86208E-02	-1.77688E-03
5000.	-3.18356E+01	1.16105E-01	5.74535E-02	1.52669E-02	-1.09084E-03	-3.28086E-03	-1.76834E-03
10000.	-3.14549E+01	1.12080E-02	3.75806E-03	-1.00216E-03	-2.02657E-03	-1.36406E-03	-6.16274E-04
20000.	-3.15643E+01	-2.69513E-02	-1.69052E-02	-8.31717E-03	-3.31404E-03	-1.08242E-03	-2.94451E-04
Equal Temp.	-3.99202E+01	6.04131E+00	-1.99447E+00	9.39512E-02	1.34081E-02	-1.22695E-02	-2.34071E-02

See appendix for Chebyshev fit details.

$He^{2+} + H \rightarrow He^{+} + H^{+}$

Maxwellian – Maxwellian



Electron Capture Rate Coefficients for
$H + He^{2+} \rightarrow He^{+} + H^{+}$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	H Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	400000.
1.0E+00	1.69E-07	2.14E-07	1.46E-07	5.84E-08	2.30E-08	2.88E-09*	2.32E-10**
2.0E+00	1.69E-07	2.14E-07	1.46E-07	5.83E-08	2.30E-08	2.87E-09*	2.32E-10**
4.0E+00	1.69E-07	2.13E-07	1.46E-07	5.83E-08	2.30E-08	2.87E-09*	2.32E-10**
7.0E+00	1.69E-07	2.13E-07	1.46E-07	5.83E-08	2.30E-08	2.87E-09*	2.32E-10**
1.0E+01	1.69E-07	2.13E-07	1.46E-07	5.82E-08	2.30E-08	2.87E-09*	2.32E-10**
2.0E+01	1.69E-07	2.12E-07	1.46E-07	5.82E-08	2.29E-08	2.87E-09*	2.32E-10**
4.0E+01	1.69E-07	2.12E-07	1.45E-07	5.81E-08	2.29E-08	2.87E-09*	2.32E-10**
7.0E+01	1.69E-07	2.11E-07	1.45E-07	5.80E-08	2.29E-08	2.87E-09*	2.32E-10**
1.0E+02	1.68E-07	2.10E-07	1.44E-07	5.79E-08	2.29E-08	2.87E-09*	2.32E-10**
2.0E+02	1.68E-07	2.09E-07	1.44E-07	5.77E-08	2.29E-08	2.87E-09*	2.33E-10**
4.0E+02	1.68E-07	2.07E-07	1.42E-07	5.75E-08	2.30E-08	2.87E-09*	2.33E-10**
7.0E+02	1.68E-07	2.04E-07	1.41E-07	5.73E-08	2.30E-08	2.87E-09*	2.34E-10**
1.0E+03	1.67E-07	2.03E-07	1.40E-07	5.72E-08	2.31E-08	2.87E-09*	2.35E-10**
2.0E+03	1.67E-07	1.98E-07	1.38E-07	5.71E-08	2.34E-08	2.89E-09*	2.37E-10**
4.0E+03	1.67E-07	1.92E-07	1.35E-07	5.74E-08	2.40E-08	2.94E-09	2.41E-10**
7.0E+03	1.66E-07	1.84E-07	1.32E-07	5.79E-08	2.48E-08	3.02E-09	2.46E-10**
1.0E+04	1.65E-07	1.78E-07	1.29E-07	5.85E-08	2.56E-08	3.10E-09	2.50E-10**
2.0E+04	1.61E-07	1.62E-07	1.20E-07	5.95E-08	2.79E-08	3.45E-09	2.65E-10**

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

H Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-3.12062E+01	-1.74693E-02	-8.09939E-03	-4.73707E-03	-3.77313E-03	-2.52573E-03	-1.15127E-03
20000.	-3.08555E+01	-1.09982E-01	-5.81296E-02	-2.43325E-02	-9.22165E-03	-3.45740E-03	-1.26420E-03
40000.	-3.15751E+01	-7.97019E-02	-4.01409E-02	-1.65010E-02	-6.97466E-03	-3.39246E-03	-1.78170E-03
70000.	-3.33202E+01	3.45964E-04	1.03619E-02	9.16031E-03	3.80511E-03	1.91641E-04	-1.02574E-03
100000.	-3.51051E+01	6.70775E-02	4.92944E-02	2.76338E-02	1.12857E-02	2.98288E-03	1.23487E-05
200000.	-3.92776E+01	5.41764E-02	4.43066E-02	2.88380E-02	1.56821E-02	6.68430E-03	2.77948E-03
400000.	-4.43133E+01	4.82448E-02	3.15762E-02	1.67996E-02	7.00532E-03	2.75563E-03	5.24833E-04

See appendix for Chebyshev fit details.

$$H + He^{2+} \rightarrow He^{+} + H^{+}$$

Beam – Maxwellian



Electron Capture Cross Sections for
$He^{2+} + H_2 \rightarrow He^{+}$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.4E+02	1.64E+07	3.70E-17
1.7E+02	1.81E+07	5.83E-17
2.0E+02	1.96E+07	7.31E-17
2.6E+02	2.24E+07	8.77E-17
4.0E+02	2.78E+07	1.07E-16
7.0E+02	3.68E+07	1.32E-16
1.0E+03	4.39E+07	1.52E-16
2.0E+03	6.21E+07	2.20E-16
4.0E+03	8.79E+07	4.12E-16
7.0E+03	1.16E+08	6.41E-16
1.0E+04	1.39E+08	8.01E-16
2.0E+04	1.96E+08	1.07E-15
4.0E+04	2.78E+08	7.23E-16
7.0E+04	3.68E+08	3.13E-16
1.0E+05	4.39E+08	1.39E-16
2.0E+05	6.21E+08	1.46E-17
4.0E+05	8.78E+08	1.02E-18
7.0E+05	1.16E+09	9.04E-20
1.0E+06	1.39E+09	1.63E-20

References: 7, 22, 23, 24, 90, 91, 92, 93, 94, 95, 101, 142

Accuracy: 20%

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 1.4E+02 eV/amu, E_{max} = 1.0E+06 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-76.3856	-3.05275	-3.46462	-1.13751	.0656712	.340604	-.0614851	-.0248706	-.0398477

The fit represents the above cross sections with an rms deviation of 2.2%.
The maximum deviation is 4.1% at 2.0E+03 eV/amu.
See appendix for Chebyshev fit details.

$He^{2+} + H_2 \rightarrow He^{+}$



Double Electron Capture Cross Sections for
$He^{2+} + H_2 \rightarrow He + 2H^+$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
3.0E+02	2.41E+07	3.68E-17
4.0E+02	2.78E+07	6.05E-17
5.6E+02	3.29E+07	7.02E-17
7.0E+02	3.68E+07	6.63E-17
1.0E+03	4.39E+07	4.70E-17
2.0E+03	6.21E+07	1.34E-17
2.7E+03	7.22E+07	1.20E-17
4.0E+03	8.79E+07	1.56E-17
7.0E+03	1.16E+08	2.75E-17
1.0E+04	1.39E+08	3.74E-17
2.0E+04	1.96E+08	5.51E-17
2.3E+04	2.11E+08	5.71E-17
4.0E+04	2.78E+08	3.84E-17
7.0E+04	3.68E+08	9.91E-18
1.0E+05	4.39E+08	3.33E-18
2.0E+05	6.21E+08	1.62E-19
3.7E+05	8.45E+08	3.88E-21

References: 7, 22, 27, 90, 91, 92, 93, 94, 95

Accuracy: $E < 2.5 \times 10^3$ eV/amu - unknown; 2.5×10^3 eV/amu $< E < 1 \times 10^5$ eV/amu - 40%; $E > 1 \times 10^5$ eV/amu - 50%

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 3.0E+02 eV/amu, E_{max} = 3.7E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-79.5999	-3.55036	-2.20554	-1.45406	-.357248	.498461	-.146918	-.0680202	.0918861

The fit represents the above cross sections with an rms deviation of 6.8%.
The maximum deviation is 12.8% at 2.0E+03 eV/amu.
See appendix for Chebyshev fit details.

$$He^{2+} + H_2 \rightarrow He + 2H^+$$

Cross Section vs. Energy



Electron Capture Cross Sections for
$He^{2+} + He \rightarrow He^{+} + He^{+}$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.4E-04	2.15E+04	1.19E-18
4.0E-04	2.78E+04	9.33E-19
1.0E-03	4.39E+04	5.95E-19
4.0E-03	8.79E+04	3.01E-19
1.0E-02	1.39E+05	1.90E-19
4.0E-02	2.78E+05	8.18E-20
1.0E-01	4.39E+05	5.26E-20
4.0E-01	8.79E+05	3.05E-20
7.0E-01	1.16E+06	2.54E-20
1.0E+00	1.39E+06	2.29E-20
1.6E+00	1.76E+06	2.21E-20
4.0E+00	2.78E+06	3.66E-20
1.0E+01	4.39E+06	1.07E-19
4.0E+01	8.79E+06	5.15E-19
1.0E+02	1.39E+07	1.44E-18
4.0E+02	2.78E+07	6.58E-18
1.0E+03	4.39E+07	1.75E-17
4.0E+03	8.79E+07	6.42E-17
1.0E+04	1.39E+08	1.52E-16
2.0E+04	1.96E+08	2.72E-16
4.0E+04	2.78E+08	3.03E-16
7.0E+04	3.68E+08	1.89E-16
1.0E+05	4.39E+08	1.11E-16
4.0E+05	8.78E+08	3.94E-18
7.0E+05	1.16E+09	3.61E-19
1.0E+06	1.39E+09	6.99E-20
2.0E+06	1.96E+09	3.33E-21

References: 90, 91, 92, 94, 97, 117, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 265, 266, 282

Accuracy: E < 250 eV/amu - see notes; E > 250 eV/amu - 20%

Notes: (1) The data for energies less than 250 eV/amu have been obtained using the theoretical results of Cohen and Bardsley (Ref. 149). From 250 eV/amu the data have been extrapolated down to 2.5 eV/amu to join smoothly with the theoretical results.
(2) Johnson et al. (Ref. 265) have measured the rate coefficient to be 4.8×10^{-14} cm^3/s at a temperature of 300 K.
(3) Theoretical values of the rate coefficient of Bardsley et al. (Ref. 266) are: T=100 K, k=4.43×10^{-14} cm^3/s; 300 K, 4.24×10^{-14} cm^3/s; 600 K, 4.02×10^{-14} cm^3/s; 1000 K, 3.79×10^{-14} cm^3/s; 3000 K, 3.25×10^{-14} cm^3/s.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 2.4E-04 eV/amu, E_{max} = 2.0E+06 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-83.5447	.758421	-.673803	-3.68088	-2.02719	-.148166	.0237951	-.0738396	.243993

The fit represents the above cross sections with an rms deviation of 19.9%.
The maximum deviation is 52.7% at 2.0E+00 eV/amu.
See appendix for Chebyshev fit details.

$He^{2+} + He \rightarrow He^{+} + He^{+}$

Cross Section vs. Energy



Electron Capture Rate Coefficients for
$He^{2+} + He \rightarrow He^{+} + He^{+}$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	Equal Temp.	He Temp. (eV) 1.	100.	500.	1000.	5000.	10000.	20000.
1.0E+00	3.05E-14	3.05E-14	5.38E-12	7.10E-11	2.12E-10	2.25E-09	6.04E-09	1.55E-08
2.0E+00	4.41E-14	3.61E-14	5.46E-12	7.13E-11	2.12E-10	2.25E-09	6.04E-09	1.55E-08
4.0E+00	9.66E-14	5.44E-14	5.64E-12	7.17E-11	2.13E-10	2.25E-09	6.04E-09	1.55E-08
7.0E+00	2.21E-13	9.66E-14	5.90E-12	7.24E-11	2.14E-10	2.25E-09	6.05E-09	1.55E-08
1.0E+01	3.88E-13	1.53E-13	6.17E-12	7.31E-11	2.15E-10	2.26E-09	6.05E-09	1.55E-08
2.0E+01	1.19E-12	4.20E-13	7.11E-12	7.54E-11	2.18E-10	2.26E-09	6.06E-09	1.55E-08
4.0E+01	3.68E-12	1.24E-12	9.12E-12	8.01E-11	2.25E-10	2.27E-09	6.07E-09	1.55E-08
7.0E+01	9.12E-12	3.03E-12	1.25E-11	8.74E-11	2.35E-10	2.29E-09	6.10E-09	1.55E-08
1.0E+02	1.62E-11	5.38E-12	1.62E-11	9.48E-11	2.45E-10	2.31E-09	6.13E-09	1.56E-08
2.0E+02	4.94E-11	1.63E-11	3.11E-11	1.21E-10	2.79E-10	2.38E-09	6.21E-09	1.57E-08
4.0E+02	1.50E-10	4.96E-11	7.08E-11	1.80E-10	3.52E-10	2.51E-09	6.38E-09	1.59E-08
7.0E+02	3.52E-10	1.21E-10	1.50E-10	2.79E-10	4.71E-10	2.71E-09	6.64E-09	1.61E-08
1.0E+03	5.98E-10	2.12E-10	2.45E-10	3.91E-10	5.98E-10	2.92E-09	6.91E-09	1.64E-08
2.0E+03	1.63E-09	5.98E-10	6.42E-10	8.28E-10	1.08E-09	3.64E-09	7.80E-09	1.74E-08
4.0E+03	4.40E-09	1.63E-09	1.69E-09	1.93E-09	2.25E-09	5.20E-09	9.66E-09	1.93E-08
7.0E+03	9.66E-09	3.64E-09	3.71E-09	4.01E-09	4.40E-09	7.80E-09	1.25E-08	2.21E-08
1.0E+04	1.55E-08	6.04E-09	6.13E-09	6.47E-09	6.91E-09	1.06E-08	1.55E-08	2.48E-08
2.0E+04	3.28E-08	1.55E-08	1.56E-08	1.59E-08	1.64E-08	2.03E-08	2.48E-08	3.28E-08

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
1.	-5.01963E+01	6.98068E+00	4.23808E-01	-3.99615E-01	1.34075E-01	-2.04534E-02	-6.31634E-03
100.	-4.61698E+01	4.09338E+00	1.25172E+00	-1.33442E-01	-1.61511E-01	2.93643E-02	3.24079E-02
500.	-4.34226E+01	2.58703E+00	1.14624E+00	1.66058E-01	-1.03603E-01	-4.97857E-02	9.07394E-03
1000.	-4.20905E+01	1.98401E+00	1.00005E+00	2.44794E-01	-4.33832E-02	-5.53954E-02	-1.11309E-02
5000.	-3.87801E+01	8.87534E-01	5.45829E-01	2.27039E-01	4.75018E-02	-1.20021E-02	-1.64614E-02
10000.	-3.72149E+01	5.46062E-01	3.51272E-01	1.61982E-01	4.68205E-02	1.29171E-03	-8.13292E-03
20000.	-3.56466E+01	2.80620E-01	1.85802E-01	9.16706E-02	3.15681E-02	5.20354E-03	-2.25604E-03
Equal Temp.	-4.86572E+01	7.31907E+00	4.09101E-02	-2.98991E-01	1.03426E-01	-6.77656E-02	6.42732E-03

See appendix for Chebyshev fit details.

$He^{2+} + He \rightarrow He^{+} + He^{+}$

Maxwellian – Maxwellian



Electron Capture Rate Coefficients for
$He + He^{2+} \rightarrow He^{+} + He^{+}$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	2.50E-08	5.92E-08	8.41E-08	6.94E-08	4.87E-08	1.84E-08	1.49E-09
2.0E+00	2.15E-08	5.40E-08	8.40E-08	6.94E-08	4.87E-08	1.84E-08	1.49E-09
4.0E+00	2.11E-08	5.33E-08	8.40E-08	6.94E-08	4.87E-08	1.84E-08	1.49E-09
7.0E+00	2.11E-08	5.33E-08	8.39E-08	6.93E-08	4.87E-08	1.84E-08	1.49E-09
1.0E+01	2.11E-08	5.33E-08	8.39E-08	6.93E-08	4.87E-08	1.83E-08	1.49E-09
2.0E+01	2.11E-08	5.32E-08	8.38E-08	6.92E-08	4.87E-08	1.83E-08	1.49E-09
4.0E+01	2.11E-08	5.31E-08	8.36E-08	6.92E-08	4.86E-08	1.83E-08	1.49E-09
7.0E+01	2.11E-08	5.30E-08	8.34E-08	6.91E-08	4.86E-08	1.83E-08	1.49E-09
1.0E+02	2.12E-08	5.30E-08	8.33E-08	6.90E-08	4.86E-08	1.83E-08	1.49E-09
2.0E+02	2.13E-08	5.29E-08	8.29E-08	6.88E-08	4.85E-08	1.82E-08	1.49E-09
4.0E+02	2.15E-08	5.28E-08	8.23E-08	6.85E-08	4.84E-08	1.82E-08	1.50E-09
7.0E+02	2.19E-08	5.27E-08	8.18E-08	6.82E-08	4.84E-08	1.81E-08	1.50E-09
1.0E+03	2.22E-08	5.28E-08	8.13E-08	6.80E-08	4.83E-08	1.81E-08	1.50E-09
2.0E+03	2.35E-08	5.31E-08	8.01E-08	6.74E-08	4.82E-08	1.80E-08	1.51E-09
4.0E+03	2.60E-08	5.38E-08	7.84E-08	6.66E-08	4.80E-08	1.79E-08	1.53E-09
7.0E+03	2.93E-08	5.47E-08	7.63E-08	6.56E-08	4.78E-08	1.78E-08	1.56E-09
1.0E+04	3.22E-08	5.55E-08	7.47E-08	6.47E-08	4.76E-08	1.78E-08	1.58E-09
2.0E+04	3.96E-08	5.70E-08	7.03E-08	6.19E-08	4.68E-08	1.79E-08	1.67E-09

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-3.50304E+01	2.06181E-01	1.92296E-01	4.56411E-02	4.15838E-02	-1.64172E-02	5.89475E-03
20000.	-3.34502E+01	-1.69065E-03	4.25743E-02	-4.85577E-03	1.65855E-02	-1.02931E-02	4.96346E-03
40000.	-3.26729E+01	-7.16579E-02	-3.83983E-02	-1.62201E-02	-5.97032E-03	-1.96773E-03	-5.29171E-04
70000.	-3.30216E+01	-4.42089E-02	-2.40130E-02	-1.09682E-02	-4.88645E-03	-2.28859E-03	-1.09577E-03
100000.	-3.36938E+01	-1.54667E-02	-7.60687E-03	-3.31747E-03	-1.90471E-03	-1.15086E-03	-7.19346E-04
200000.	-3.56481E+01	-1.62738E-02	-3.50236E-03	1.52811E-03	2.39339E-03	1.00308E-03	6.34282E-04
500000.	-4.06040E+01	3.82554E-02	2.68882E-02	1.51307E-02	6.78931E-03	2.33434E-03	4.95076E-04

See appendix for Chebyshev fit details.

$He + He^{2+} \rightarrow He^{+} + He^{+}$

Beam – Maxwellian



Double Electron Capture Cross Sections for
He^{2+} + He -> He + He^{2+}

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
7.0E+00	3.68E+06	4.62E-16
1.0E+01	4.39E+06	4.48E-16
2.0E+01	6.21E+06	4.10E-16
4.0E+01	8.79E+06	3.83E-16
7.0E+01	1.16E+07	3.58E-16
1.0E+02	1.39E+07	3.44E-16
2.0E+02	1.96E+07	3.19E-16
4.0E+02	2.78E+07	2.88E-16
7.0E+02	3.68E+07	2.62E-16
1.0E+03	4.39E+07	2.44E-16
2.0E+03	6.21E+07	2.10E-16
4.0E+03	8.79E+07	1.73E-16
7.0E+03	1.16E+08	1.46E-16
1.0E+04	1.39E+08	1.29E-16
2.0E+04	1.96E+08	9.68E-17
4.0E+04	2.78E+08	6.08E-17
7.0E+04	3.68E+08	2.73E-17
1.0E+05	4.39E+08	9.83E-18
2.0E+05	6.21E+08	6.36E-19
4.0E+05	8.78E+08	2.09E-20

References: 90, 91, 92, 94, 140, 144, 145, 148, 150, 151, 152, 153, 282

Accuracy: 20% throughout the energy range

Note: (1) The data below 100 eV/amu are theoretical data of reference 153 (Ferguson and Moiseiwitsch).

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 7.0E+00 eV/amu, E_{max} = 4.0E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-75.3410	-3.70255	-2.16467	-1.23660	-.590825	-.179613	.0396927	.102768	.0774058

The fit represents the above cross sections with an rms deviation of 3.2%.
The maximum deviation is 6.8% at 2.0E+04 eV/amu.
See appendix for Chebyshev fit details.

$He^{2+} + He \rightarrow He + He^{2+}$



Double Electron Capture Rate Coefficients for
He^{2+} + He -> He + He^{2+}

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	Equal Temp.	He Temp. (eV) 1.	100.	500.	1000.	5000.	10000.	20000.
2.0E+00	5.21E-12	5.68E-13	2.92E-09	5.58E-09	7.14E-09	1.17E-08	1.38E-08	1.57E-08
4.0E+00	1.36E-10	1.94E-11	2.94E-09	5.58E-09	7.15E-09	1.17E-08	1.38E-08	1.57E-08
7.0E+00	5.25E-10	1.36E-10	2.98E-09	5.60E-09	7.16E-09	1.17E-08	1.38E-08	1.57E-08
1.0E+01	8.99E-10	3.22E-10	3.02E-09	5.61E-09	7.16E-09	1.17E-08	1.38E-08	1.57E-08
2.0E+01	1.73E-09	9.54E-10	3.14E-09	5.65E-09	7.19E-09	1.17E-08	1.38E-08	1.57E-08
4.0E+01	2.60E-09	1.76E-09	3.36E-09	5.73E-09	7.24E-09	1.17E-08	1.38E-08	1.57E-08
7.0E+01	3.36E-09	2.45E-09	3.65E-09	5.84E-09	7.31E-09	1.17E-08	1.38E-08	1.57E-08
1.0E+02	3.90E-09	2.90E-09	3.90E-09	5.95E-09	7.38E-09	1.17E-08	1.38E-08	1.57E-08
2.0E+02	5.12E-09	3.91E-09	4.58E-09	6.30E-09	7.60E-09	1.18E-08	1.39E-08	1.57E-08
4.0E+02	6.60E-09	5.12E-09	5.57E-09	6.88E-09	8.00E-09	1.19E-08	1.39E-08	1.58E-08
7.0E+02	8.00E-09	6.30E-09	6.60E-09	7.60E-09	8.52E-09	1.21E-08	1.40E-08	1.58E-08
1.0E+03	8.97E-09	7.14E-09	7.38E-09	8.18E-09	8.97E-09	1.22E-08	1.41E-08	1.58E-08
2.0E+03	1.10E-08	8.97E-09	9.10E-09	9.60E-09	1.01E-08	1.27E-08	1.43E-08	1.60E-08
4.0E+03	1.31E-08	1.10E-08	1.11E-08	1.14E-08	1.17E-08	1.35E-08	1.48E-08	1.61E-08
7.0E+03	1.48E-08	1.27E-08	1.28E-08	1.29E-08	1.31E-08	1.43E-08	1.53E-08	1.64E-08
1.0E+04	1.57E-08	1.38E-08	1.38E-08	1.39E-08	1.41E-08	1.50E-08	1.57E-08	1.66E-08
2.0E+04	1.69E-08	1.57E-08	1.57E-08	1.58E-08	1.58E-08	1.62E-08	1.66E-08	1.69E-08

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 2.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
1.	-4.15949E+01	4.05943E+00	-1.86222E+00	9.53695E-01	-4.31628E-01	1.13605E-01	1.67894E-03
100.	-3.79435E+01	9.09167E-01	1.91522E-01	-7.51345E-02	-3.33315E-02	9.43860E-03	3.83286E-03
500.	-3.72996E+01	5.30896E-01	1.95295E-01	-2.21372E-03	-3.37015E-02	-9.19923E-03	3.89947E-03
1000.	-3.70111E+01	3.90475E-01	1.68883E-01	1.91542E-02	-2.19861E-02	-1.21205E-02	-1.71700E-04
5000.	-3.63583E+01	1.43130E-01	8.03033E-02	2.65227E-02	7.89255E-04	-4.74954E-03	-3.10575E-03
10000.	-3.61072E+01	7.57604E-02	4.50781E-02	1.73388E-02	2.65954E-03	-1.73029E-03	-1.75170E-03
20000.	-3.59012E+01	2.91780E-02	1.75429E-02	6.74273E-03	9.12966E-04	-9.12778E-04	-9.71523E-04
Equal Temp.	-4.01576E+01	3.11099E+00	-1.35722E+00	7.35133E-01	-4.32956E-01	1.86361E-01	-6.65206E-02

See appendix for Chebyshev fit details.

$He^{2+} + He \rightarrow He + He^{2+}$

Maxwellian — Maxwellian



Double Electron Capture Rate Coefficients for
$He + He^{2+} \rightarrow He + He^{2+}$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	400000.
1.0E+00	1.79E-08	1.90E-08	1.69E-08	1.00E-08	4.31E-09	3.95E-10*	9.21E-12**
2.0E+00	1.79E-08	1.90E-08	1.69E-08	1.00E-08	4.31E-09	3.95E-10*	9.22E-12**
4.0E+00	1.79E-08	1.90E-08	1.69E-08	1.00E-08	4.31E-09	3.95E-10*	9.24E-12**
7.0E+00	1.79E-08	1.90E-08	1.68E-08	9.99E-09	4.31E-09	3.94E-10*	9.26E-12**
1.0E+01	1.79E-08	1.90E-08	1.68E-08	9.99E-09	4.31E-09	3.94E-10*	9.28E-12**
2.0E+01	1.79E-08	1.90E-08	1.68E-08	9.97E-09	4.30E-09	3.94E-10*	9.32E-12**
4.0E+01	1.79E-08	1.90E-08	1.68E-08	9.94E-09	4.29E-09	3.94E-10*	9.39E-12**
7.0E+01	1.79E-08	1.89E-08	1.67E-08	9.91E-09	4.29E-09	3.93E-10*	9.45E-12**
1.0E+02	1.79E-08	1.89E-08	1.67E-08	9.89E-09	4.28E-09	3.93E-10*	9.51E-12**
2.0E+02	1.79E-08	1.89E-08	1.66E-08	9.83E-09	4.27E-09	3.93E-10*	9.65E-12**
4.0E+02	1.79E-08	1.88E-08	1.65E-08	9.76E-09	4.26E-09	3.93E-10*	9.86E-12**
7.0E+02	1.79E-08	1.88E-08	1.64E-08	9.68E-09	4.26E-09	3.93E-10*	1.01E-11**
1.0E+03	1.78E-08	1.87E-08	1.63E-08	9.63E-09	4.26E-09	3.94E-10*	1.03E-11**
2.0E+03	1.78E-08	1.86E-08	1.61E-08	9.50E-09	4.28E-09	3.97E-10*	1.08E-11**
4.0E+03	1.78E-08	1.84E-08	1.58E-08	9.34E-09	4.34E-09	4.06E-10*	1.17E-11**
7.0E+03	1.77E-08	1.81E-08	1.54E-08	9.20E-09	4.42E-09	4.21E-10*	1.28E-11**
1.0E+04	1.76E-08	1.79E-08	1.51E-08	9.08E-09	4.49E-09	4.39E-10	1.38E-11**
2.0E+04	1.74E-08	1.71E-08	1.42E-08	8.74E-09	4.65E-09	5.07E-10	1.71E-11**

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-3.56880E+01	-1.04856E-02	-6.22291E-03	-3.54221E-03	-2.04547E-03	-1.06539E-03	-4.43802E-04
20000.	-3.56043E+01	-3.86901E-02	-2.24247E-02	-1.12950E-02	-5.53082E-03	-2.66314E-03	-1.23198E-03
40000.	-3.58798E+01	-6.74672E-02	-3.60384E-02	-1.56499E-02	-6.43318E-03	-2.74207E-03	-1.24709E-03
70000.	-3.69171E+01	-5.93014E-02	-2.54553E-02	-7.53787E-03	-2.21925E-03	-1.22692E-03	-8.87276E-04
100000.	-3.85034E+01	2.31019E-02	2.39390E-02	1.48606E-02	4.98011E-03	-1.80325E-05	-1.31207E-03
200000.	-4.32277E+01	7.43187E-02	6.13383E-02	4.01508E-02	2.17045E-02	9.21588E-03	3.71392E-03
400000.	-5.05148E+01	2.45518E-01	1.33039E-01	5.66724E-02	2.09008E-02	7.13076E-03	2.41652E-03

See appendix for Chebyshev fit details.

$He + He^{2+} \rightarrow He + He^{2+}$

Beam – Maxwellian



Electron Capture Cross Sections for
$He^{2+} + Li \rightarrow He^{+} + Li^{+}$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.0E+02	1.39E+07	4.22E-15
2.0E+02	1.96E+07	6.95E-15
4.0E+02	2.78E+07	9.64E-15
7.0E+02	3.68E+07	1.11E-14
1.0E+03	4.39E+07	1.15E-14
2.0E+03	6.21E+07	1.14E-14
4.0E+03	8.79E+07	1.06E-14
7.0E+03	1.16E+08	9.25E-15
1.0E+04	1.39E+08	7.39E-15
1.5E+04	1.70E+08	4.51E-15
2.0E+04	1.96E+08	2.24E-15
3.0E+04	2.41E+08	6.70E-16
5.0E+04	3.11E+08	2.11E-16
7.0E+04	3.68E+08	1.43E-16
1.0E+05	4.39E+08	9.57E-17
1.5E+05	5.38E+08	5.79E-17
2.0E+05	6.21E+08	3.82E-17
2.5E+05	6.94E+08	2.32E-17
3.0E+05	7.61E+08	1.38E-17
4.0E+05	8.78E+08	5.17E-18
5.0E+05	9.82E+08	2.26E-18
5.9E+05	1.07E+09	1.21E-18

References: 230, 231, 232, 234, 241, 242, 244, 246

Accuracy: $E < 2x10^5$ eV/amu - 30%; $E > 2x10^5$ eV/amu - 40%

Notes: (1) For data of the sum of the single and double (total) cross sections see Kadota et al references 243 and 245.
(2) For the electron capture cross sections for Li^{+}, Li^{2+} + He see references 249, 250, 251, and 254.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 1.0E+02 eV/amu, E_{max} = 5.9E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-69.9969	-3.86583	-2.18937	.0319758	.0613388	-.163069	-.271234	-.111623	.204819

The fit represents the above cross sections with an rms deviation of 11.0%.
The maximum deviation is 22.5% at 5.0E+04 eV/amu.
See appendix for Chebyshev fit details.

$He^{2+} + Li \rightarrow He^{+} + Li^{+}$



Electron Capture Rate Coefficients for
$Li + He^{2+} \rightarrow He^{+} + Li^{+}$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	Li Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	1.02E-06	4.40E-07	9.71E-08	5.26E-08	4.20E-08*	2.37E-08*	2.20E-09**
2.0E+00	1.02E-06	4.39E-07	9.71E-08	5.26E-08	4.20E-08*	2.37E-08*	2.23E-09**
4.0E+00	1.02E-06	4.39E-07	9.71E-08	5.26E-08	4.20E-08*	2.37E-08*	2.22E-09**
7.0E+00	1.02E-06	4.39E-07	9.71E-08	5.26E-08	4.20E-08*	2.37E-08*	2.22E-09**
1.0E+01	1.02E-06	4.39E-07	9.71E-08	5.26E-08	4.20E-08*	2.37E-08*	2.22E-09**
2.0E+01	1.02E-06	4.38E-07	9.71E-08	5.26E-08	4.20E-08*	2.37E-08*	2.22E-09**
4.0E+01	1.02E-06	4.38E-07	9.72E-08	5.26E-08	4.20E-08*	2.37E-08*	2.22E-09**
7.0E+01	1.01E-06	4.37E-07	9.72E-08	5.26E-08	4.20E-08*	2.36E-08*	2.22E-09**
1.0E+02	1.01E-06	4.37E-07	9.72E-08	5.26E-08	4.20E-08	2.36E-08*	2.22E-09**
2.0E+02	1.00E-06	4.37E-07	9.74E-08	5.26E-08	4.20E-08	2.36E-08*	2.22E-09**
4.0E+02	9.93E-07	4.39E-07	9.76E-08	5.26E-08	4.20E-08	2.35E-08*	2.22E-09**
7.0E+02	9.81E-07	4.41E-07	9.81E-08	5.26E-08	4.19E-08	2.34E-08*	2.22E-09**
1.0E+03	9.69E-07	4.42E-07	9.86E-08	5.26E-08	4.19E-08	2.34E-08*	2.23E-09**
2.0E+03	9.35E-07	4.45E-07	1.01E-07	5.27E-08	4.19E-08	2.33E-08*	2.24E-09**
4.0E+03	8.74E-07	4.45E-07	1.06E-07	5.29E-08	4.18E-08	2.31E-08*	2.25E-09**
7.0E+03	8.03E-07	4.38E-07	1.16E-07	5.36E-08	4.18E-08	2.30E-08*	2.26E-09**
1.0E+04	7.48E-07	4.29E-07	1.25E-07	5.46E-08	4.18E-08	2.28E-08	2.26E-09**
2.0E+04	6.25E-07	3.94E-07	1.48E-07	5.95E-08	4.21E-08	2.25E-08	2.28E-09**

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

Li Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-2.78062E+01	-1.87483E-01	-1.15766E-01	-5.63757E-02	-2.07864E-02	-4.18092E-03	1.20353E-03
20000.	-2.93011E+01	-2.38001E-02	-2.23600E-02	-2.23853E-02	-1.68257E-02	-8.26641E-03	-2.16146E-03
40000.	-3.21294E+01	1.47297E-01	1.06505E-01	5.96604E-02	2.42461E-02	4.76111E-03	-3.24711E-03
70000.	-3.34880E+01	3.30410E-02	2.70085E-02	2.03864E-02	1.35662E-02	7.66428E-03	3.81877E-03
100000.	-3.39729E+01	-1.63250E-03	3.97915E-04	1.54890E-03	1.48096E-03	1.08056E-03	6.82015E-04
200000.	-3.51439E+01	-2.29742E-02	-1.03372E-02	-3.67071E-03	-8.35726E-04	-5.21814E-04	-6.93460E-05
500000.	-3.98397E+01	1.28943E-02	5.77153E-03	3.54906E-03	-1.13575E-03	6.68011E-04	-9.25001E-04

See appendix for Chebyshev fit details.

$Li + He^{2+} \rightarrow He^{+} + Li^{+}$

Beam – Maxwellian



Double Electron Capture Cross Sections for
$He^{2+} + Li \rightarrow He + Li^{2+}$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
6.1E+03	1.08E+08	9.12E-18
7.0E+03	1.16E+08	1.30E-17
9.0E+03	1.32E+08	2.39E-17
1.0E+04	1.39E+08	2.99E-17
1.5E+04	1.70E+08	4.44E-17
2.0E+04	1.96E+08	4.03E-17
3.0E+04	2.41E+08	2.66E-17
5.0E+04	3.11E+08	1.42E-17
7.0E+04	3.68E+08	9.17E-18
1.0E+05	4.39E+08	5.86E-18
1.5E+05	5.38E+08	3.12E-18
2.0E+05	6.21E+08	1.17E-18
2.5E+05	6.94E+08	3.16E-19
3.0E+05	7.61E+08	1.05E-19
4.0E+05	8.78E+08	1.62E-20
5.0E+05	9.82E+08	1.03E-20

References: 231, 232, 241, 242, 244, 246

Accuracy: 30%

Notes: (1) In the region ($1\text{-}3\times10^4$ eV/amu) where the cross section is a maximum the experimental data fluctuates by a factor of $\pm$ 2. We believe that the value quoted can be relied on to 30%.

(2) For data of the sum of the single and double electron capture see Kadota et al references 243 and 245.

(3) For electron capture cross sections of Li^+, Li^{2+} + He see references 249, 250, 251, and 254.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 6.1E+03 eV/amu, E_{max} = 5.0E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-81.1888	-3.55493	-2.11208	-.149724	-.232101	.216608	.275776	.0980514	.0367760

The fit represents the above cross sections with an rms deviation of 3.1%.
The maximum deviation is 8.4% at 2.5E+05 eV/amu.
See appendix for Chebyshev fit details.

$He^{2+} + Li \rightarrow He + Li^{2+}$

Cross Section vs. Energy



Double Electron Capture Rate Coefficients for
$Li + He^{2+} \rightarrow He + Li^{2+}$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	Li Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	400000.
1.0E+00	4.14E-09	7.91E-09	5.19E-09	3.37E-09	2.57E-09	7.25E-10*	1.43E-11**
2.0E+00	4.14E-09	7.90E-09	5.19E-09	3.37E-09	2.57E-09	7.25E-10*	1.43E-11**
4.0E+00	4.13E-09	7.90E-09	5.19E-09	3.37E-09	2.57E-09	7.25E-10*	1.43E-11**
7.0E+00	4.12E-09	7.89E-09	5.19E-09	3.37E-09	2.57E-09	7.24E-10*	1.43E-11**
1.0E+01	4.12E-09	7.88E-09	5.19E-09	3.37E-09	2.57E-09	7.23E-10*	1.43E-11**
2.0E+01	4.10E-09	7.87E-09	5.19E-09	3.37E-09	2.57E-09	7.22E-10*	1.44E-11**
4.0E+01	4.09E-09	7.85E-09	5.19E-09	3.37E-09	2.57E-09	7.20E-10*	1.44E-11**
7.0E+01	4.08E-09	7.83E-09	5.19E-09	3.37E-09	2.57E-09	7.18E-10*	1.45E-11**
1.0E+02	4.07E-09	7.81E-09	5.19E-09	3.37E-09	2.57E-09	7.17E-10	1.45E-11**
2.0E+02	4.07E-09	7.76E-09	5.19E-09	3.37E-09	2.56E-09	7.14E-10	1.47E-11**
4.0E+02	4.08E-09	7.70E-09	5.19E-09	3.37E-09	2.56E-09	7.10E-10	1.49E-11**
7.0E+02	4.13E-09	7.62E-09	5.19E-09	3.38E-09	2.56E-09	7.07E-10	1.51E-11**
1.0E+03	4.18E-09	7.55E-09	5.19E-09	3.38E-09	2.55E-09	7.05E-10	1.53E-11**
2.0E+03	4.31E-09	7.34E-09	5.19E-09	3.38E-09	2.55E-09	7.03E-10	1.59E-11**
4.0E+03	4.51E-09	6.99E-09	5.19E-09	3.39E-09	2.54E-09	7.07E-10	1.68E-11**
7.0E+03	4.68E-09	6.60E-09	5.16E-09	3.40E-09	2.53E-09	7.16E-10	1.81E-11**
1.0E+04	4.79E-09	6.31E-09	5.11E-09	3.40E-09	2.52E-09	7.26E-10	1.93E-11**
2.0E+04	4.95E-09	5.71E-09	4.90E-09	3.40E-09	2.50E-09	7.51E-10	2.35E-11**

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

Li Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-3.85237E+01	7.92423E-02	5.63215E-02	1.72118E-02	-5.29456E-03	-7.47819E-03	-2.52581E-03
20000.	-3.74702E+01	-1.30340E-01	-7.61020E-02	-3.32737E-02	-9.67526E-03	-1.15041E-04	2.05695E-03
40000.	-3.81678E+01	-1.43365E-02	-1.25875E-02	-1.01317E-02	-6.93426E-03	-3.91276E-03	-1.88591E-03
70000.	-3.90109E+01	4.43065E-03	2.10527E-03	3.09061E-04	-4.54509E-04	-5.21843E-04	-3.71172E-04
100000.	-3.95717E+01	-1.18800E-02	-5.53013E-03	-2.04086E-03	-8.30113E-04	-4.87010E-04	-3.29738E-04
200000.	-4.20984E+01	1.49204E-03	1.62234E-02	1.48608E-02	6.78142E-03	1.37283E-03	-7.46341E-04
400000.	-4.97186E+01	1.87794E-01	1.11341E-01	5.29149E-02	2.28786E-02	8.25402E-03	3.38953E-03

See appendix for Chebyshev fit details.

$$Li + He^{2+} \rightarrow He + Li^{2+}$$

Beam – Maxwellian



B. Electron Capture into Excited States

(* denotes rate coefficient data also)

	Page
Electron Capture by H^+	**B-2**
$H^+ + H \rightarrow H(2s) + H^+$	B-2*
$H^+ + H \rightarrow H(2p) + H^+$	B-8*
$H^+ + H_2 \rightarrow H(2s)$	B-14
$H^+ + H_2 \rightarrow H(2p)$	B-14
$H^+ + H_2 \rightarrow H(\text{Lyman-}\alpha\ \text{Total})$	B-16
$H^+ + H_2 \rightarrow H(\text{Balmer-}\alpha)$	B-18
$H^+ + H_2 \rightarrow H(\text{Balmer-}\beta)$	B-18
$H^+ + H_2 \rightarrow H(3s)$	B-20
$H^+ + H_2 \rightarrow H(3p)$	B-20
$H^+ + H_2 \rightarrow H(3d)$	B-20
$H^+ + He \rightarrow H(2s) + He^+$	B-22*
$H^+ + He \rightarrow H(2p) + He^+$	B-28*
$H^+ + He \rightarrow H(\text{Balmer-}\alpha) + He^+$	B-34*
$H^+ + He \rightarrow H(3s) + He^+$	B-40*
$H^+ + He \rightarrow H(3p) + He^+$	B-46*
$H^+ + He \rightarrow H(3d) + He^+$	B-52*
$H^+ + Li \rightarrow H(2s) + Li^+$	B-58
$H^+ + Li \rightarrow H(2p) + Li^+$	B-58
$H^+ + Li \rightarrow H(3s) + Li^+$	B-60
$H^+ + Li \rightarrow H(3p) + Li^+$	B-60
$H^+ + Li \rightarrow H(3d) + Li^+$	B-60
Electron Capture by He^+	**B-62**
$He^+ + H_2 \rightarrow H(2^1S + 2^3S)$	B-62
$He^+ + H_2 \rightarrow He(2^1P)$	B-64
$He^+ + H_2 \rightarrow He(3^1P)$	B-64
$He^+ + H_2 \rightarrow He(\lambda = 447.1\ nm)$	B-66
$He^+ + H_2 \rightarrow He(\lambda = 587.6\ nm)$	B-66
$He^+ + H_2 \rightarrow He(\lambda = 667.8\ nm)$	B-66
$He^+ + He \rightarrow He(2^1S + 2^3S) + He^+$	B-68*
$He^+ + He \rightarrow He(2^1P) + He^+$	B-74*
$He^+ + He \rightarrow He(3^1S) + He^+$	B-80
$He^+ + He \rightarrow He(3^1P) + He^+$	B-80
$He^+ + He \rightarrow He(3^1D) + He^+$	B-80
$He^+ + He \rightarrow He(3^3S) + He^+$	B-82
$He^+ + He \rightarrow He(3^3P) + He^+$	B-82
$He^+ + He \rightarrow He(3^3D) + He^+$	B-82
$He^+ + He \rightarrow He(4^1S) + He^+$	B-84
$He^+ + He \rightarrow He(4^1P) + He^+$	B-84
$He^+ + He \rightarrow He(4^1D) + He^+$	B-84
$He^+ + He \rightarrow He(4^3S) + He^+$	B-86
$He^+ + He \rightarrow He(4^3P) + He^+$	B-86
$He^+ + He \rightarrow He(4^3D) + He^+$	B-86
$He^+ + Li \rightarrow He(53.7\ nm) + Li^+$	B-88
$He^+ + Li \rightarrow He(58.4\ nm) + Li^+$	B-88

Electron Capture by He^{2+} B-90

He^{2+} + H -> He^{+}(2s) + H^{+} B-90*
He^{2+} + H -> He^{+}(2p) + H^{+} B-96*
He^{2+} + H -> He^{+}(Lyman-α) + H^{+} B-102
He^{2+} + H -> He^{+}(Lyman-β) + H^{+} B-102
He^{2+} + H -> He^{+}(Lyman-γ) + H^{+} B-102
He^{2+} + H -> He^{+}(Balmer-α) + H^{+} B-102
He^{2+} + H -> He^{+}(3p) + H^{+} B-104
He^{2+} + H -> He^{+}(3s+3d) + H^{+} B-104
He^{2+} + H -> He^{+}(4p) + H^{+} B-104
He^{2+} + H_2 -> He^{+}(2s) B-106
He^{2+} + H_2 -> He^{+}(2p) B-106
He^{2+} + H_2 -> He^{+}(Lyman-α) B-108
He^{2+} + H_2 -> He^{+}(Lyman-β) B-108
He^{2+} + H_2 -> He^{+}(Lyman-γ) B-108
He^{2+} + H_2 -> He^{+}(Balmer-α) B-108
He^{2+} + H_2 -> He^{+}(3p) B-110
He^{2+} + H_2 -> He^{+}(3s+3d) B-110
He^{2+} + H_2 -> He^{+}(4p) B-110
He^{2+} + He -> He^{+}(2s) + He^{+} B-112*
He^{2+} + Li -> He^{+}(2p) + Li^{+} B-118
He^{2+} + Li -> He^{+}(3p) + Li^{+} B-118
He^{2+} + Li -> He^{+}(4p) + Li^{+} B-118
He^{2+} + Li -> He^{+}(3s+3d) + Li^{+} B-118
He^{2+} + Li -> He^{+}(4s+4d) + Li^{+} B-118
He^{2+} + Li -> He^{+}(Lyman-α) + Li^{+} B-120
He^{2+} + Li -> He^{+}(Lyman-β) + Li^{+} B-120
He^{2+} + Li -> He^{+}(Lyman-γ) + Li^{+} B-120
He^{2+} + Li -> He^{+}(Balmer-α) + Li^{+} B-122
He^{2+} + Li -> He^{+}(Balmer-β) + Li^{+} B-122
He^{2+} + Li -> He^{+}(Balmer-γ) + Li^{+} B-122
He^{2+} + Li -> He^{+}(Balmer-σ) + Li^{+} B-122

Electron Capture Cross Sections for
$H^+ + H \rightarrow H(2s) + H^+$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.0E+03	6.21E+07	1.41E-18
3.0E+03	7.61E+07	2.77E-18
4.0E+03	8.79E+07	4.59E-18
5.0E+03	9.82E+07	6.78E-18
6.0E+03	1.08E+08	8.91E-18
7.0E+03	1.16E+08	1.19E-17
1.0E+04	1.39E+08	1.98E-17
2.0E+04	1.96E+08	3.65E-17
4.0E+04	2.78E+08	2.17E-17
5.0E+04	3.11E+08	1.39E-17
6.0E+04	3.40E+08	9.05E-18
7.0E+04	3.68E+08	5.56E-18
1.0E+05	4.39E+08	1.93E-18
2.0E+05	6.21E+08	2.59E-19

References: 1, 11, 12, 13, 14, 15, 16, 70, 71, 72

Accuracy: Energy < $3x10^3$ eV/amu - factor 2; $3x10^3$ eV/amu < E < $1x10^5$ eV/amu - 30%; E > $1x10^5$ eV/amu - 50%

Notes: (1) For energies greater than $5x10^3$ eV/amu, the crossed beam results of references 12 and 14 are in good agreement with the data obtained using the tungsten oven H cell employed in the other experimental references.
(2) No corrections were applied for cascading but the effect is believed to be no more than 5%.
(3) Data for energies greater than $1.2x10^5$ eV/amu were taken from the theoretical paper by Shakeshaft (ref. 70) who computed the cross sections using time-dependent impact parameter coupled-state equations.
(4) Cross sections in the energy range $1x10^3$ - $5x10^3$ eV/amu rely on the experimental results of Morgan et al (ref. 15) and theoretical results of ref. 71 and 72.
(5) Cross sections for the 2s state were found by mixing the 2s and 2p states with an electric field.

For a Chebyshev fit of the above cross section it is necessary to use the following parameters.
E_{min} = 2.0E+03 eV/amu, E_{max} = 2.0E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-80.2974	-.711716	-2.02522	-.193493	.250985	.0856986	-.0237110	-.0276091	-.00357455

The fit represents the above cross sections with an rms deviation of 1.3%.
The maximum deviation is 2.4% at 7.0E+04 eV/amu.
See appendix for Chebyshev fit details.

$$H^+ + H \rightarrow H(2s) + H^+$$



Electron Capture Rate Coefficients for
$H^+ + H \rightarrow H(2s) + H^+$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	Equal Temp.	H Temp. (eV) 500.	1000.	2000.	4000.	7000.	10000.	20000.
1.0E+02	1.86E-14	1.48E-11	9.32E-11	4.11E-10	1.35E-09	2.66E-09	3.50E-09	4.23E-09
2.0E+02	2.59E-12	2.51E-11	1.16E-10	4.52E-10	1.40E-09	2.70E-09	3.52E-09	4.23E-09
4.0E+02	3.83E-11	5.41E-11	1.69E-10	5.36E-10	1.50E-09	2.77E-09	3.57E-09	4.23E-09
7.0E+02	1.69E-10	1.16E-10	2.63E-10	6.70E-10	1.65E-09	2.87E-09	3.62E-09	4.22E-09
1.0E+03	3.72E-10	1.98E-10	3.72E-10	8.12E-10	1.79E-09	2.96E-09	3.68E-09	4.22E-09
2.0E+03	1.30E-09	5.80E-10	8.12E-10	1.30E-09	2.23E-09	3.25E-09	3.83E-09	4.20E-09
4.0E+03	2.96E-09	1.55E-09	1.79E-09	2.23E-09	2.96E-09	3.68E-09	4.05E-09	4.14E-09
7.0E+03	4.05E-09	2.80E-09	2.96E-09	3.25E-09	3.68E-09	4.05E-09	4.20E-09	4.03E-09
1.0E+04	4.23E-09	3.59E-09	3.68E-09	3.83E-09	4.05E-09	4.20E-09	4.23E-09	3.90E-09
2.0E+04	3.42E-09	4.22E-09	4.22E-09	4.20E-09	4.14E-09	4.03E-09	3.90E-09	3.42E-09

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+02 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

H Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
500.	-4.38672E+01	3.12293E+00	-1.60004E-01	-3.10412E-01	-2.36074E-02	1.60083E-02	6.19524E-03
1000.	-4.24130E+01	2.14668E+00	6.37440E-02	-2.57968E-01	-5.56388E-02	1.72468E-02	8.38626E-03
2000.	-4.10369E+01	1.31982E+00	1.21498E-01	-1.62370E-01	-6.16572E-02	4.43261E-03	8.17248E-03
4000.	-3.98120E+01	6.46066E-01	8.59976E-02	-8.09184E-02	-4.48832E-02	-5.52064E-03	3.62665E-03
7000.	-3.90771E+01	2.57541E-01	2.91176E-02	-4.35279E-02	-2.76600E-02	-6.63736E-03	5.07727E-04
10000.	-3.87811E+01	9.11341E-02	-6.21649E-03	-3.19458E-02	-1.89226E-02	-5.58779E-03	-2.28503E-04
20000.	-3.86539E+01	-7.79801E-02	-5.02818E-02	-2.49832E-02	-9.47407E-03	-2.56718E-03	-3.98810E-04
Equal Temp.	-4.60750E+01	5.50322E+00	-2.23799E+00	4.38735E-01	-2.58100E-01	1.18690E-01	-1.94375E-02

See appendix for Chebyshev fit details.

$H^+ + H \rightarrow H(2s) + H^+$

Maxwellian – Maxwellian



Electron Capture Rate Coefficients for
$H + H^+ \rightarrow H(2s) + H^+$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	H Energy (eV/amu) 10000.	20000.	40000.	60000.	80000.	100000.	150000.
1.0E+00	2.74E-09	7.12E-09	6.01E-09*	3.07E-09*	1.47E-09**	8.48E-10**	3.21E-10**
2.0E+00	2.74E-09	7.11E-09	6.00E-09*	3.07E-09*	1.47E-09**	8.48E-10**	3.21E-10**
4.0E+00	2.74E-09	7.08E-09	5.99E-09*	3.07E-09*	1.47E-09*	8.48E-10**	3.21E-10**
7.0E+00	2.74E-09	7.05E-09	5.97E-09*	3.07E-09*	1.47E-09*	8.48E-10**	3.21E-10**
1.0E+01	2.73E-09	7.03E-09	5.96E-09	3.06E-09*	1.47E-09*	8.48E-10**	3.21E-10**
2.0E+01	2.73E-09	6.97E-09	5.94E-09	3.06E-09*	1.47E-09*	8.49E-10**	3.21E-10**
4.0E+01	2.74E-09	6.89E-09	5.90E-09	3.05E-09*	1.47E-09*	8.50E-10*	3.21E-10**
7.0E+01	2.75E-09	6.81E-09	5.86E-09	3.05E-09*	1.48E-09*	8.51E-10*	3.21E-10**
1.0E+02	2.76E-09	6.75E-09	5.83E-09	3.04E-09*	1.48E-09*	8.53E-10*	3.22E-10**
2.0E+02	2.82E-09	6.60E-09	5.75E-09	3.04E-09*	1.49E-09*	8.57E-10*	3.23E-10**
4.0E+02	2.95E-09	6.40E-09	5.64E-09	3.04E-09	1.51E-09*	8.66E-10*	3.25E-10**
7.0E+02	3.14E-09	6.21E-09	5.52E-09	3.05E-09	1.54E-09*	8.80E-10*	3.28E-10**
1.0E+03	3.30E-09	6.08E-09	5.42E-09	3.05E-09	1.57E-09*	8.94E-10*	3.30E-10**
2.0E+03	3.69E-09	5.77E-09	5.14E-09	3.05E-09	1.65E-09*	9.42E-10*	3.37E-10**
4.0E+03	4.10E-09	5.35E-09	4.69E-09	3.00E-09	1.75E-09*	1.03E-09*	3.52E-10**
7.0E+03	4.35E-09	4.90E-09	4.19E-09	2.88E-09	1.82E-09*	1.14E-09*	3.87E-10**
1.0E+04	4.40E-09	4.57E-09	3.82E-09	2.74E-09	1.84E-09	1.20E-09*	4.28E-10*
2.0E+04	4.08E-09	3.80E-09	3.07E-09	2.35E-09	1.74E-09	1.26E-09*	5.48E-10*

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

H Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-3.91129E+01	2.48347E-01	9.68613E-02	-2.22662E-02	-5.16334E-02	-2.86624E-02	-4.22512E-03
20000.	-3.78678E+01	-2.64262E-01	-1.22968E-01	-4.41549E-02	-1.57924E-02	-6.00867E-03	-1.43873E-03
40000.	-3.81878E+01	-2.65205E-01	-1.50964E-01	-6.70004E-02	-2.27907E-02	-3.96914E-03	2.12259E-03
60000.	-3.92919E+01	-7.86994E-02	-5.90240E-02	-4.39866E-02	-2.75365E-02	-1.19525E-02	-2.20171E-03
80000.	-4.05364E+01	1.08994E-01	4.62123E-02	-5.38404E-03	-2.37480E-02	-1.87047E-02	-8.06980E-03
100000.	-4.15580E+01	1.82433E-01	1.07666E-01	3.51890E-02	-4.95054E-03	-1.53694E-02	-1.20555E-02
150000.	-4.35210E+01	1.78485E-01	1.28691E-01	7.63149E-02	3.78040E-02	1.49449E-02	2.43305E-03

See appendix for Chebyshev fit details.

$$H + H^+ \rightarrow H(2s) + H^+$$



Electron Capture Cross Sections for
$H^+ + H \rightarrow H(2p) + H^+$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
6.0E+02	3.40E+07	6.05E-18
7.0E+02	3.68E+07	8.69E-18
1.0E+03	4.39E+07	1.85E-17
1.5E+03	5.38E+07	3.10E-17
2.0E+03	6.21E+07	3.05E-17
3.0E+03	7.61E+07	2.87E-17
4.0E+03	8.79E+07	2.89E-17
7.0E+03	1.16E+08	3.06E-17
1.0E+04	1.39E+08	2.99E-17
2.0E+04	1.96E+08	1.79E-17
3.0E+04	2.41E+08	1.01E-17
4.0E+04	2.78E+08	5.83E-18
7.0E+04	3.68E+08	1.46E-18
1.0E+05	4.39E+08	6.18E-19
2.0E+05	6.21E+08	1.41E-19
2.4E+05	6.80E+08	1.00E-19

References: 17, 18, 19, 20, 70

Accuracy: $E < 2x10^3$ eV/amu - 40%; $2x10^3$ eV/amu $< E < 3x10^4$ eV/amu - 25%; $E > 3x10^4$ eV/amu - unknown

Notes: (1) Thomas in ref. 20 has applied a correction to ref. 19.
(2) All cross sections have been normalized to either the $H^+ + H_2 \rightarrow H(2p) + H_2^+$ or $e^- + H(1s) \rightarrow e^- + H(2p)$ cross sections.
(3) Cross sections for H impact energies greater than $3x10^4$ eV/amu rely on computations of Shakeshaft (ref. 70). Data were obtained using time-dependent impact parameter coupled-state equations.
(4) The cross section (σ_i) for capture into state i is determined by measuring the emission cross section (σ_{ij}) for a transition from state i to j, correcting for any branching from state i, and correcting for cascading from upper states.
(5) Cascading from upper states to the 2p state is less than 5%. Thus $\sigma(2p) \simeq \sigma$(Lyman-α) emission cross section.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 6.0E+02 eV/amu, E_{max} = 2.4E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-79.9812	-2.30968	-1.73621	.110265	.0489823	.217654	-.0576541	-.0617916	.0399669

The fit represents the above cross sections with an rms deviation of 2.8%.
The maximum deviation is 5.8% at 2.0E+04 eV/amu.
See appendix for Chebyshev fit details.

$$H^+ + H \rightarrow H(2p) + H^+$$



Electron Capture Rate Coefficients for
$H^+ + H \rightarrow H(2p) + H^+$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	Equal Temp.	H Temp. (eV) 50.	100.	500.	1000.	5000.	10000.	20000.
4.0E+01	5.58E-13	1.28E-12	1.45E-11	5.48E-10	1.16E-09	2.88E-09	3.03E-09	2.47E-09
7.0E+01	1.45E-11	6.91E-12	3.22E-11	5.92E-10	1.19E-09	2.89E-09	3.03E-09	2.47E-09
1.0E+02	5.74E-11	1.95E-11	5.74E-11	6.36E-10	1.22E-09	2.89E-09	3.03E-09	2.46E-09
2.0E+02	3.33E-10	1.13E-10	1.81E-10	7.73E-10	1.32E-09	2.91E-09	3.03E-09	2.46E-09
4.0E+02	8.99E-10	4.11E-10	4.88E-10	1.02E-09	1.49E-09	2.93E-09	3.02E-09	2.45E-09
7.0E+02	1.49E-09	8.38E-10	8.99E-10	1.32E-09	1.71E-09	2.96E-09	3.01E-09	2.43E-09
1.0E+03	1.90E-09	1.17E-09	1.22E-09	1.56E-09	1.90E-09	2.99E-09	3.00E-09	2.41E-09
2.0E+03	2.68E-09	1.93E-09	1.96E-09	2.17E-09	2.38E-09	3.05E-09	2.95E-09	2.36E-09
4.0E+03	3.07E-09	2.70E-09	2.71E-09	2.79E-09	2.88E-09	3.06E-09	2.84E-09	2.25E-09
7.0E+03	2.84E-09	3.05E-09	3.05E-09	3.06E-09	3.07E-09	2.95E-09	2.65E-09	2.09E-09
1.0E+04	2.47E-09	3.03E-09	3.03E-09	3.02E-09	3.00E-09	2.78E-09	2.47E-09	1.96E-09
2.0E+04	1.59E-09	2.47E-09	2.46E-09	2.44E-09	2.41E-09	2.19E-09	1.96E-09	1.59E-09

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 4.0E+01 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

H Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
50.	-4.42667E+01	3.68258E+00	-1.49113E+00	1.59924E-01	3.18208E-03	-6.21884E-02	2.01934E-02
100.	-4.30926E+01	2.69099E+00	-8.96227E-01	-7.60448E-02	4.64849E-02	-4.79878E-02	6.77301E-03
500.	-4.08437E+01	9.29788E-01	-1.25719E-01	-1.89331E-01	-3.21599E-02	5.88363E-03	2.45244E-03
1000.	-4.02252E+01	5.06698E-01	-5.16460E-02	-1.44926E-01	-4.91357E-02	2.77891E-03	6.06624E-03
5000.	-3.93773E+01	-6.84093E-02	-8.56458E-02	-6.23155E-02	-2.73651E-02	-6.00122E-03	5.77484E-04
10000.	-3.94214E+01	-1.65783E-01	-1.04137E-01	-4.96524E-02	-1.77141E-02	-4.10624E-03	-1.51335E-04
20000.	-3.98550E+01	-1.76608E-01	-9.99144E-02	-4.14670E-02	-1.28283E-02	-2.76171E-03	-2.52981E-04
Equal Temp.	-4.38515E+01	3.35460E+00	-2.04640E+00	5.49669E-01	-2.67503E-01	7.12101E-02	2.83428E-03

See appendix for Chebyshev fit details.

$H^+ + H \rightarrow H(2p) + H^+$

Maxwellian — Maxwellian



Electron Capture Rate Coefficients for
$H + H^+ \rightarrow H(2p) + H^+$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	H Energy (eV/amu) 10000.	20000.	40000.	60000.	80000.	100000.	200000.
1.0E+00	4.14E-09	3.51E-09	1.62E-09*	7.27E-10*	4.16E-10**	2.72E-10**	8.76E-11**
2.0E+00	4.13E-09	3.50E-09	1.62E-09*	7.27E-10*	4.16E-10**	2.72E-10**	8.76E-11**
4.0E+00	4.12E-09	3.50E-09	1.61E-09*	7.28E-10*	4.16E-10**	2.72E-10**	8.76E-11**
7.0E+00	4.11E-09	3.49E-09	1.61E-09*	7.28E-10*	4.16E-10**	2.72E-10**	8.77E-11**
1.0E+01	4.10E-09	3.49E-09	1.61E-09*	7.28E-10*	4.16E-10*	2.72E-10**	8.77E-11**
2.0E+01	4.08E-09	3.47E-09	1.61E-09	7.28E-10*	4.16E-10*	2.72E-10**	8.77E-11**
4.0E+01	4.06E-09	3.46E-09	1.61E-09	7.29E-10*	4.16E-10*	2.73E-10**	8.78E-11**
7.0E+01	4.03E-09	3.44E-09	1.60E-09	7.30E-10*	4.17E-10*	2.73E-10*	8.79E-11**
1.0E+02	4.00E-09	3.42E-09	1.60E-09	7.31E-10*	4.17E-10*	2.73E-10*	8.79E-11**
2.0E+02	3.94E-09	3.39E-09	1.60E-09	7.35E-10*	4.19E-10*	2.75E-10*	8.81E-11**
4.0E+02	3.86E-09	3.34E-09	1.60E-09	7.42E-10*	4.22E-10*	2.77E-10*	8.83E-11**
7.0E+02	3.77E-09	3.27E-09	1.60E-09	7.53E-10*	4.27E-10*	2.80E-10*	8.80E-11**
1.0E+03	3.69E-09	3.22E-09	1.60E-09	7.64E-10*	4.32E-10*	2.82E-10*	8.71E-11**
2.0E+03	3.50E-09	3.05E-09	1.59E-09	7.95E-10	4.49E-10*	2.92E-10*	8.40E-11**
4.0E+03	3.23E-09	2.77E-09	1.57E-09	8.40E-10	4.85E-10*	3.12E-10*	8.06E-11**
7.0E+03	2.93E-09	2.46E-09	1.50E-09	8.76E-10	5.29E-10*	3.42E-10*	8.00E-11**
1.0E+04	2.69E-09	2.25E-09	1.43E-09	8.88E-10	5.60E-10*	3.69E-10*	8.18E-11**
2.0E+04	2.10E-09	1.76E-09	1.23E-09	8.56E-10	5.99E-10	4.24E-10*	9.59E-11**

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

H Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-3.89330E+01	-2.63479E-01	-1.46563E-01	-6.61908E-02	-2.64497E-02	-9.58772E-03	-3.32633E-03
20000.	-3.92573E+01	-2.65213E-01	-1.59204E-01	-7.45512E-02	-2.59202E-02	-4.40931E-03	1.71443E-03
40000.	-4.05806E+01	-8.23737E-02	-5.96947E-02	-4.22268E-02	-2.58735E-02	-1.17419E-02	-3.02153E-03
60000.	-4.19604E+01	9.70829E-02	4.52599E-02	7.29529E-04	-1.72682E-02	-1.54561E-02	-7.90347E-03
80000.	-4.30171E+01	1.56735E-01	9.77827E-02	3.77245E-02	1.87118E-03	-9.85709E-03	-9.46753E-03
100000.	-4.38524E+01	1.71818E-01	1.11913E-01	5.35817E-02	1.64028E-02	-2.85661E-04	-5.46080E-03
200000.	-4.63327E+01	-9.29799E-03	6.37363E-03	2.65872E-02	3.44937E-02	2.70598E-02	1.28430E-02

See appendix for Chebyshev fit details.

$$H + H^+ \rightarrow H(2p) + H^+$$

Beam – Maxwellian



Electron Capture Cross Sections for

$H^+ + H_2 \rightarrow H(2s,2p)$

H(2s) Energy eV/amu	H(2s) Cross Section cm^2	H(2p) Energy eV/amu	H(2p) Cross Section cm^2
2.0E+03	2.64E-18	4.0E+03	1.99E-17
4.0E+03	5.08E-18	7.0E+03	2.37E-17
7.0E+03	9.24E-18	1.0E+04	2.64E-17
1.0E+04	1.37E-17	1.4E+04	2.72E-17
2.0E+04	2.69E-17	2.0E+04	2.35E-17
2.5E+04	2.79E-17	4.0E+04	8.74E-18
4.0E+04	2.21E-17	7.0E+04	3.39E-18
7.0E+04	9.53E-18	1.0E+05	1.75E-18
1.0E+05	4.20E-18	1.5E+05	1.00E-18
2.0E+05	4.24E-19		

References: H(2s) - 13, 16, 18, 121, 122, 123, 124, 125, 126; H(2p) - 18, 122, 124, 128

Accuracy: See notes

Notes: (1) Large errors are usually associated with measuring emission from excited states to obtain either electron capture, dissociative or excitation emission cross sections. The cross sections reported here are emission cross sections from electron capture collisions. The H(2s) cross sections are determined by observing the excited H atom in an electric field which mixes the 2s state with the 2p state. The scatter of the data in the references is a factor of 2 for H(2s) capture and 20% for H(2p). We estimate that the data presented here have a confidence level to within 50%.
(2) For the collisions in an electromagnetic field sufficient to mix the 2s and 2p, the Lyman-α emission cross section is obtained from $\sigma(2s)+\sigma(2p)$.
(3) The cross section (σ_i) for capture into state i is determined by measuring the emission cross section for a transition from state i to j, correcting for any branching from state i and correcting for cascading from upper states.
(4) Since cascading to the 2p state is less than 5% $\sigma(2p)=\sigma$(Lyman-α) emission.

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

H(2s) E_{min} = 2.0E+03 eV/amu, E_{max} = 2.0E+05 eV/amu
H(2p) E_{min} = 4.0E+03 eV/amu, E_{max} = 1.5E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

	A0	A1	A2	A3	A4	A5	A6	A7	A8
H(2s)	-79.7252	-.533265	-1.56108	-.431130	.0810709	.0519074	-.0478876	-.00197365	
H(2p)	-78.7561	-1.60375	-.734004	.106262	.120750	-.0254921	.00605546	.0276214	.0341018

The fit represents the H(2s) cross section with an rms deviation of 1.3%.
The maximum deviation is 2.5% at 1.0E+04 eV/amu.

The fit represents the H(2p) cross section with an rms deviation of 1.5%.
The maximum deviation is 8.5% at 1.3E+05 eV/amu.

See appendix for Chebyshev fit details.

$H^+ + H_2 \rightarrow H(2s, 2p)$



Line Emission Cross Sections of Electron Capture Collisions for
$H^+ + H_2 \rightarrow H(Ly\alpha)$ Total

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
8.2E+01	1.26E+07	7.15E-20
1.0E+02	1.39E+07	1.37E-19
2.0E+02	1.96E+07	9.34E-19
4.0E+02	2.78E+07	4.56E-18
7.0E+02	3.68E+07	1.20E-17
1.0E+03	4.39E+07	1.69E-17
1.5E+03	5.38E+07	1.97E-17
2.0E+03	6.21E+07	2.11E-17
3.0E+03	7.61E+07	2.39E-17
4.0E+03	8.79E+07	2.94E-17
7.0E+03	1.16E+08	4.30E-17
1.0E+04	1.39E+08	5.22E-17
1.4E+04	1.64E+08	5.61E-17
2.0E+04	1.96E+08	4.87E-17
4.0E+04	2.78E+08	2.97E-17
7.0E+04	3.68E+08	1.94E-17
1.0E+05	4.39E+08	1.46E-17
1.4E+05	5.20E+08	1.19E-17

References: 124, 128, 187, 188, 193, 195, 197

Accuracy: 40%

Notes: (1) The cross section σ_i for capture into state i is determined by measuring the emission cross section (σ_{ij}) for a transition from state i to j, correcting for any branching from state i and correcting for cascading from upper states.
(2) The capture cross section into excited states is measured by making use of the Doppler shift. The total $\sigma(Ly\alpha)$ was determined from the total of the shifted and unshifted radiation.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 8.2E+01 eV/amu, E_{max} = 1.4E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-79.2785	2.29244	-1.84261	.243522	-.113691	.0377226	.155522	-.0201741	-.0977164

The fit represents the above cross sections with an rms deviation of 3.4%.
The maximum deviation is 6.0% at 2.0E+04 eV/amu.
See appendix for Chebyshev fit details.

$H^+ + H_2 \rightarrow H(\text{Lyman}-\alpha)\text{Total}$



Line Emission Cross Sections of Electron Capture Collisions for

$H^+ + H_2 \rightarrow H(H_\alpha, H_\beta)$

(H_α) Energy eV/amu	Cross Section cm^2
5.0E+02	2.44E-19
7.0E+02	6.10E-19
1.0E+03	1.13E-18
2.0E+03	2.13E-18
4.0E+03	3.11E-18
7.0E+03	4.17E-18
1.0E+04	4.95E-18
2.0E+04	7.18E-18
3.0E+04	8.52E-18
4.0E+04	7.68E-18
7.0E+04	3.34E-18
1.0E+05	1.25E-18
2.0E+05	1.06E-19
4.0E+05	4.17E-21

(H_β) Energy eV/amu	Cross Section cm^2
5.0E+02	4.24E-20
7.0E+02	9.36E-20
1.0E+03	1.61E-19
2.0E+03	3.12E-19
4.0E+03	4.27E-19
7.0E+03	4.96E-19
1.0E+04	5.17E-19
2.0E+04	4.82E-19
4.0E+04	2.94E-19
7.0E+04	1.11E-19
1.0E+05	4.45E-20
1.5E+05	1.74E-20

References: (H_α) - 131, 133, 134, 135, 136, 137, 138, 139; (H_β) - 137, 138, 139

Accuracy: See notes

Notes: (1) Large errors are usually associated with measuring optical emission from excited states which arise from electron capture, dissociative or excitation collisions. The cross sections reported here are for electron capture collisions resulting in Balmer-α emission (n=3->2), λ=6562.79 Å) and Balmer-β emission (n=4->2, λ=4861.33 Å). Errors arise mainly from the detecter calibration which for the visible spectrum varies over many orders of magnitude. We can only estimate the error to be more than a factor of 2 for H_α.

(2) Tne H_α emission cross section is found from the following relation - $\sigma(H_\alpha) = \sigma(3s) + .118\sigma(3p) + \sigma(3d)$.

(3) Cascading from upper levels contribute no more than 5% to the measured cross sections.

(4) H_α cross sections below $3x10^3$ eV/amu were found by normalizing Hess's results (ref. 137 and 138) to other results at $3x10^3$ eV/amu.

(5) Very little data are available for Balmer-β (H_β) emission cross sections.

(6) H_β cross sections below $3x10^3$ eV/amu were found by normalizing Hess's data (refs. 137 and 138) at $3x10^3$ eV/amu to Hughes et al data (ref. 139).

(7) No estimate is given for errors in the H_β data.

(8) Data are available for capture in the 4s state (see refs. 130 and 132).

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

(H_α) E_{min} = 5.0E+02 eV/amu, E_{max} = 4.0E+05 eV/amu
(H_β) E_{min} = 5.0E+02 eV/amu, E_{max} = 1.5E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

	A0	A1	A2	A3	A4	A5	A6	A7	A8
(H_α)	-83.9403	-1.52055	-2.66829	-.703430	-.220654	.245822	.0371514	-.0525595	-.0727916
(H_β)	-86.8907	-.394318	-1.51191	-.164906	-.148036	.0816213	.0463644	.0309715	.00687359

The fit represents the (H_α) cross section with an rms deviation of 2.2%.
The maximum deviation is 5.1% at 2.0E+04 eV/amu.
The fit represents the (H_β) cross section with an rms deviation of 0.6%.
The maximum deviation is 1.1% at 1.0E+03 eV/amu.
See appendix for Chebyshev fit details.

$$H^+ + H_2 \rightarrow H(H_\alpha, H_\beta)$$



Electron Capture Cross Sections for

$H^+ + H_2 \rightarrow H(3s,3p,3d) + H_2^+$

H(3s)		H(3p)		H(3d)	
Energy eV/amu	Cross Section cm^2	Energy eV/amu	Cross Section cm^2	Energy eV/amu	Cross Section cm^2
1.0E+03	3.33E-19	2.2E+03	1.18E-18	1.4E+03	4.27E-19
2.0E+03	5.25E-19	4.0E+03	3.49E-18	2.0E+03	5.56E-19
4.0E+03	8.67E-19	5.9E+03	4.19E-18	4.0E+03	8.89E-19
7.0E+03	1.45E-18	7.0E+03	4.17E-18	7.0E+03	1.27E-18
1.0E+04	2.10E-18	1.0E+04	3.86E-18	1.0E+04	1.57E-18
2.0E+04	4.90E-18	2.0E+04	2.83E-18	1.1E+04	1.60E-18
3.2E+04	7.21E-18	3.2E+04	1.63E-18	2.0E+04	1.04E-18
4.0E+04	6.54E-18	4.0E+04	1.07E-18	4.0E+04	3.94E-19
7.0E+04	2.88E-18	7.0E+04	2.96E-19	7.0E+04	1.03E-19
1.0E+05	1.16E-18	1.0E+05	1.35E-19	1.0E+05	3.09E-20
2.0E+05	9.99E-20	1.6E+05	4.76E-20	1.4E+05	8.93E-21
3.0E+05	1.83E-20	2.0E+05	3.13E-20	2.0E+05	4.39E-21
		3.4E+05	1.01E-20	3.0E+05	2.47E-21

References: H(3s) 129, 130, 131, 132, 133, 134, 135; H(3p) 122, 131, 133, 134, 136; H(3d) 131, 134, 136

Accuracy: See notes

Notes: (1) Large errors are usually associated with measuring optical emission from excited states which arise either from electron capture, dissociative, or excitation collisions. The cross sections reported here are for electron capture of the fast proton into a fast H atom. Errors arise mainly from absolute calibration of the detector which for the visible spectrum varies over orders of magnitude from blue to red. We estimate the data presented to be accurate to a factor of 2.

(2) The cross section (σ_i) for capture into state i is determined by measuring the emission cross section (σ_{ij}) for a transition from state i to j, correcting for any branching from state i and correcting for cascading from upper states.

(3) For the n=3 levels cascading is less than 5%.

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

H(3s) E_{min} = 1.0E+03 eV/amu, E_{max} = 3.0E+05 eV/amu
H(3p) E_{min} = 2.2E+03 eV/amu, E_{max} = 3.4E+05 eV/amu
H(3d) E_{min} = 1.4E+03 eV/amu, E_{max} = 3.0E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

	A0	A1	A2	A3	A4	A5	A6	A7	A8
H(3s)	-84.0392	-.728083	-2.03457	-.860699	.0489259	.212089	.0319211	-.0748778	-.0241098
H(3p)	-84.5768	-2.73107	-1.38696	.326474	.0471408	.0427739	-.0758510	-.0181499	.0430074
H(3d)	-86.9132	-2.76179	-1.74766	.0813682	.392354	.101286	.0203703	.00824074	-.0805033

The fit represents the H(3s) cross section with an rms deviation of 2.3%.
The maximum deviation is 4.0% at 7.0E+04 eV/amu.

The fit represents the H(3p) cross section with an rms deviation of 1.3%.
The maximum deviation is 2.6% at 1.6E+05 eV/amu.

The fit represents the H(3d) cross section with an rms deviation of 5.0%.
The maximum deviation is 10.4% at 7.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$H^+ + H_2 \rightarrow H(3s, 3p, 3d)$



Electron Capture Cross Sections for
$H^+ + He \rightarrow H(2s) + He^+$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.8E+03	7.35E+07	5.16E-20
4.0E+03	8.79E+07	1.45E-19
7.0E+03	1.16E+08	5.91E-19
1.0E+04	1.39E+08	1.03E-18
1.4E+04	1.64E+08	1.47E-18
2.0E+04	1.96E+08	2.44E-18
3.0E+04	2.41E+08	5.61E-18
4.0E+04	2.78E+08	7.74E-18
5.0E+04	3.11E+08	8.08E-18
7.0E+04	3.68E+08	6.60E-18
1.0E+05	4.39E+08	3.75E-18
2.0E+05	6.21E+08	4.23E-19

References: 16, 121, 125, 126, 183, 184, 189, 190, 191

Accuracy: 50%

Note: (1) The $\sigma(2s)$ cross section is found by mixing the 2s state with the 2p state by an electric field.

For Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 2.8E+03 eV/amu, E_{max} = 2.0E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-83.0478	1.54023	-1.55597	-.364564	-.186200	.0423931	-.0198717	-.166625	-.0726702

The fit represents the above cross sections with an rms deviation of 2.5%.
The maximum deviation is 4.2% at 5.0E+04 eV/amu.
See appendix for Chebyshev fit details.

$H^+ + He \rightarrow H(2s) + He^+$



Electron Capture Rate Coefficients for
$H^+ + He \rightarrow H(2s) + He^+$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	Equal Temp.	He Temp. (eV) 500.	1000.	2000.	4000.	7000.	10000.	20000.
2.0E+02	2.85E-16	3.73E-15	4.23E-14	4.60E-13	3.64E-12	1.48E-11	3.22E-11	1.30E-10
4.0E+02	8.08E-14	1.07E-13	3.25E-13	1.32E-12	5.90E-12	1.89E-11	3.78E-11	1.40E-10
7.0E+02	1.18E-12	9.32E-13	1.61E-12	3.64E-12	1.03E-11	2.58E-11	4.68E-11	1.57E-10
1.0E+03	4.16E-12	2.94E-12	4.16E-12	7.24E-12	1.58E-11	3.35E-11	5.68E-11	1.74E-10
2.0E+03	2.70E-11	1.83E-11	2.11E-11	2.70E-11	4.07E-11	6.58E-11	9.64E-11	2.35E-10
4.0E+03	1.20E-10	8.03E-11	8.55E-11	9.64E-11	1.20E-10	1.59E-10	2.03E-10	3.72E-10
7.0E+03	3.54E-10	2.43E-10	2.51E-10	2.68E-10	3.02E-10	3.54E-10	4.08E-10	5.86E-10
1.0E+04	6.20E-10	4.53E-10	4.62E-10	4.80E-10	5.15E-10	5.68E-10	6.20E-10	7.81E-10
2.0E+04	1.21E-09	1.04E-09	1.05E-09	1.06E-09	1.08E-09	1.11E-09	1.13E-09	1.21E-09

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 2.0E+02 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
500.	-5.15547E+01	6.12867E+00	-1.10703E+00	1.83503E-01	-6.31506E-02	-4.18133E-02	-5.16887E-03
1000.	-5.02988E+01	5.12782E+00	-5.67743E-01	-2.34476E-02	-1.29792E-02	-4.58512E-02	-5.85869E-03
2000.	-4.88467E+01	4.04429E+00	-9.26042E-02	-1.42687E-01	-1.51195E-02	-3.17168E-02	-6.62145E-03
4000.	-4.73254E+01	3.01860E+00	2.18123E-01	-1.57987E-01	-4.76884E-02	-1.64944E-02	-1.20711E-03
7000.	-4.60971E+01	2.29486E+00	3.28511E-01	-1.26781E-01	-6.47340E-02	-1.05302E-02	4.12821E-03
10000.	-4.53140E+01	1.88943E+00	3.36913E-01	-1.00227E-01	-6.44137E-02	-9.37217E-03	4.93922E-03
20000.	-4.37344E+01	1.17101E+00	2.58458E-01	-4.78379E-02	-4.40287E-02	-8.69295E-03	2.45757E-03
Equal Temp.	-5.19683E+01	7.03024E+00	-1.90084E+00	5.72322E-01	-2.68910E-01	2.66381E-02	-1.15409E-02

See appendix for Chebyshev fit details.

$H^+ + He \rightarrow H(2s) + He^+$

Maxwellian – Maxwellian



Electron Capture Rate Coefficients for
$He + H^+ \rightarrow H(2s) + He^+$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	60000.	80000.	100000.	120000.
1.0E+00	1.43E-10**	4.81E-10**	2.14E-09*	2.46E-09*	2.10E-09**	1.64E-09**	1.02E-09**
2.0E+00	1.43E-10**	4.81E-10**	2.14E-09*	2.46E-09*	2.10E-09**	1.64E-09**	1.02E-09**
4.0E+00	1.43E-10**	4.82E-10**	2.14E-09*	2.46E-09*	2.10E-09**	1.64E-09**	1.02E-09**
7.0E+00	1.42E-10**	4.84E-10*	2.14E-09*	2.46E-09*	2.10E-09*	1.64E-09**	1.02E-09**
1.0E+01	1.42E-10**	4.85E-10*	2.13E-09*	2.46E-09*	2.10E-09*	1.63E-09**	1.02E-09**
2.0E+01	1.42E-10**	4.88E-10*	2.13E-09*	2.46E-09*	2.10E-09*	1.63E-09*	1.02E-09**
4.0E+01	1.43E-10*	4.93E-10*	2.12E-09*	2.46E-09*	2.10E-09*	1.62E-09*	1.02E-09**
7.0E+01	1.44E-10*	5.00E-10*	2.11E-09*	2.46E-09*	2.10E-09*	1.61E-09*	1.02E-09**
1.0E+02	1.45E-10*	5.07E-10*	2.11E-09*	2.46E-09*	2.10E-09*	1.61E-09*	1.02E-09**
2.0E+02	1.49E-10*	5.28E-10*	2.09E-09*	2.46E-09*	2.10E-09*	1.59E-09*	1.03E-09**
4.0E+02	1.59E-10*	5.68E-10*	2.07E-09*	2.45E-09*	2.09E-09*	1.57E-09*	1.03E-09*
7.0E+02	1.77E-10*	6.23E-10*	2.05E-09*	2.42E-09*	2.08E-09*	1.56E-09*	1.04E-09*
1.0E+03	1.97E-10*	6.73E-10*	2.03E-09*	2.39E-09*	2.06E-09*	1.54E-09*	1.05E-09*
2.0E+03	2.79E-10*	8.12E-10*	1.98E-09*	2.29E-09*	1.99E-09*	1.51E-09*	1.06E-09*
4.0E+03	4.63E-10*	1.01E-09*	1.90E-09	2.12E-09*	1.87E-09*	1.46E-09*	1.07E-09*
7.0E+03	7.11E-10	1.19E-09	1.81E-09	1.92E-09*	1.72E-09*	1.38E-09*	1.05E-09*
1.0E+04	9.07E-10	1.29E-09	1.73E-09	1.78E-09*	1.59E-09*	1.31E-09*	1.02E-09*
2.0E+04	1.27E-09	1.42E-09	1.54E-09	1.47E-09*	1.31E-09*	1.12E-09*	9.16E-10*

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-4.41473E+01	1.00293E+00	5.77264E-01	1.69345E-01	-4.20278E-02	-7.82819E-02	-4.04936E-02
20000.	-4.22126E+01	5.46817E-01	2.44571E-01	2.72848E-02	-4.37757E-02	-3.23001E-02	-7.56613E-03
40000.	-4.00821E+01	-1.29340E-01	-7.16121E-02	-3.33138E-02	-1.32072E-02	-4.05181E-03	-1.23221E-03
60000.	-3.98602E+01	-1.91480E-01	-1.30757E-01	-6.51877E-02	-2.04843E-02	-1.38003E-03	2.12462E-03
80000.	-4.01499E+01	-1.66211E-01	-1.18429E-01	-6.40569E-02	-2.40368E-02	-4.53618E-03	8.86157E-04
100000.	-4.06316E+01	-1.42002E-01	-8.01346E-02	-4.14812E-02	-2.15460E-02	-9.87334E-03	-2.92063E-03
120000.	-4.14146E+01	-1.03802E-02	-2.66264E-02	-3.13729E-02	-2.24983E-02	-1.04122E-02	-2.88295E-03

See appendix for Chebyshev fit details.

$$He + H^+ \rightarrow H(2s) + He^+$$



Electron Capture Cross Sections for
$H^+ + He \rightarrow H(2p) + He^+$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
7.0E+02	3.68E+07	2.02E-20
1.0E+03	4.39E+07	4.48E-20
2.0E+03	6.21E+07	2.10E-19
4.0E+03	8.79E+07	6.83E-19
7.0E+03	1.16E+08	1.57E-18
1.0E+04	1.39E+08	2.32E-18
2.0E+04	1.96E+08	3.14E-18
3.0E+04	2.41E+08	2.88E-18
4.0E+04	2.78E+08	2.17E-18
7.0E+04	3.68E+08	8.52E-19
1.0E+05	4.39E+08	3.93E-19
2.0E+05	6.21E+08	3.86E-20
3.0E+05	7.61E+08	9.74E-21

References: 24, 46, 125, 163, 164, 165, 166, 183, 184, 185, 186, 187, 188, 245

Accuracy: 30%

Notes: (1) If cascade is neglected $\sigma(H\ Ly_\alpha) = \sigma(H\ 2p)$. Cascade contributions are less than 5%.
(2) The cross sections for capture into the 2p state or Ly_α emission have large discrepancies at energies less than $1x10^4$ eV/amu. These discrepancies are thought to be due to a small amount of H^o present in the H^+ beam during measurements. The cross section for excitation of $H + He \rightarrow H(2p)$ is approximatly 2 orders of magnitude greater than $H^+ + He \rightarrow H(2p) + He^+$.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 7.0E+02 eV/amu, E_{max} = 3.0E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-86.2710	-.0743636	-2.70622	-.368010	.0743916	.0747662	.0195366	.00225497	.0337284

The fit represents the above cross sections with an rms deviation of 1.9%.
The maximum deviation is 4.1% at 7.0E+04 eV/amu.
See appendix for Chebyshev fit details.

$H^+ + He \rightarrow H(2p) + He^+$

Cross Section vs. Energy



Electron Capture Rate Coefficients for
$H^+ + He \rightarrow H(2p) + He^+$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	Equal Temp.	He Temp. (eV) 100.	1000.	2000.	4000.	7000.	10000.	20000.
4.0E+01	3.31E-18	7.92E-17	4.90E-13	2.90E-12	1.36E-11	4.11E-11	7.62E-11	1.98E-10
7.0E+01	1.22E-15	2.29E-15	6.69E-13	3.33E-12	1.45E-11	4.25E-11	7.77E-11	1.99E-10
1.0E+02	1.36E-14	1.36E-14	8.78E-13	3.79E-12	1.54E-11	4.38E-11	7.92E-11	2.00E-10
2.0E+02	2.97E-13	2.03E-13	1.79E-12	5.52E-12	1.85E-11	4.82E-11	8.41E-11	2.05E-10
4.0E+02	2.38E-12	1.53E-12	4.61E-12	9.89E-12	2.55E-11	5.74E-11	9.42E-11	2.13E-10
7.0E+02	9.28E-12	6.00E-12	1.12E-11	1.85E-11	3.73E-11	7.18E-11	1.09E-10	2.26E-10
1.0E+03	2.02E-11	1.32E-11	2.02E-11	2.93E-11	5.05E-11	8.66E-11	1.24E-10	2.38E-10
2.0E+03	7.43E-11	5.16E-11	6.21E-11	7.43E-11	9.92E-11	1.37E-10	1.73E-10	2.75E-10
4.0E+03	1.96E-10	1.50E-10	1.61E-10	1.73E-10	1.96E-10	2.28E-10	2.57E-10	3.34E-10
7.0E+03	3.27E-10	2.76E-10	2.83E-10	2.92E-10	3.07E-10	3.27E-10	3.45E-10	3.91E-10
1.0E+04	3.98E-10	3.57E-10	3.62E-10	3.66E-10	3.75E-10	3.87E-10	3.98E-10	4.22E-10
2.0E+04	4.33E-10	4.38E-10	4.39E-10	4.39E-10	4.39E-10	4.38E-10	4.38E-10	4.33E-10

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 4.0E+01 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
100.	-5.42320E+01	7.44291E+00	-2.03521E+00	2.94202E-01	-1.85784E-01	2.90590E-02	2.43336E-02
1000.	-4.97005E+01	3.78186E+00	-7.01113E-02	-4.11509E-01	-2.83920E-02	2.87633E-02	2.44756E-03
2000.	-4.83079E+01	2.82754E+00	1.63410E-01	-3.46227E-01	-7.46954E-02	2.85450E-02	8.40255E-03
4000.	-4.69020E+01	1.95707E+00	2.50111E-01	-2.30851E-01	-9.41042E-02	9.45349E-03	1.19992E-02
7000.	-4.57899E+01	1.32780E+00	2.41351E-01	-1.38821E-01	-8.61534E-02	-6.49649E-03	8.83775E-03
10000.	-4.51295E+01	9.78457E-01	2.08879E-01	-9.24217E-02	-7.29486E-02	-1.27985E-02	5.37477E-03
20000.	-4.40453E+01	4.43773E-01	1.12537E-01	-3.78696E-02	-4.20886E-02	-1.39142E-02	-2.74451E-04
Equal Temp.	-5.47397E+01	8.28139E+00	-2.97971E+00	8.65229E-01	-5.12815E-01	1.97141E-01	-4.18893E-02

See appendix for Chebyshev fit details.

$H^+ + He \rightarrow H(2p) + He^+$

Maxwellian — Maxwellian



Electron Capture Rate Coefficients for
$He + H^+ \rightarrow H(2p) + He^+$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	60000.	80000.	100000.	200000.
1.0E+00	3.21E-10	6.15E-10	6.02E-10	3.75E-10	2.51E-10*	1.72E-10*	2.40E-11**
2.0E+00	3.21E-10	6.15E-10	6.01E-10	3.75E-10*	2.51E-10*	1.72E-10*	2.40E-11**
4.0E+00	3.20E-10	6.14E-10	6.01E-10	3.75E-10*	2.51E-10*	1.72E-10*	2.40E-11**
7.0E+00	3.20E-10	6.13E-10	6.00E-10	3.75E-10	2.51E-10*	1.72E-10*	2.40E-11**
1.0E+01	3.19E-10	6.12E-10	5.99E-10	3.75E-10	2.51E-10*	1.72E-10*	2.40E-11**
2.0E+01	3.18E-10	6.10E-10	5.98E-10	3.75E-10	2.51E-10*	1.71E-10*	2.40E-11**
4.0E+01	3.18E-10	6.08E-10	5.96E-10	3.75E-10	2.51E-10	1.71E-10*	2.40E-11**
7.0E+01	3.17E-10	6.05E-10	5.93E-10	3.75E-10	2.51E-10	1.70E-10*	2.40E-11**
1.0E+02	3.17E-10	6.03E-10	5.92E-10	3.76E-10	2.51E-10	1.70E-10*	2.40E-11**
2.0E+02	3.19E-10	5.99E-10	5.87E-10	3.76E-10	2.52E-10	1.69E-10*	2.41E-11**
4.0E+02	3.25E-10	5.94E-10	5.81E-10	3.76E-10	2.52E-10	1.68E-10	2.43E-11**
7.0E+02	3.35E-10	5.88E-10	5.72E-10	3.77E-10	2.52E-10	1.68E-10	2.45E-11**
1.0E+03	3.46E-10	5.82E-10	5.64E-10	3.77E-10	2.53E-10	1.68E-10	2.48E-11**
2.0E+03	3.75E-10	5.64E-10	5.41E-10	3.75E-10	2.53E-10	1.68E-10	2.56E-11*
4.0E+03	4.13E-10	5.34E-10	5.02E-10	3.66E-10	2.52E-10	1.71E-10	2.75E-11*
7.0E+03	4.40E-10	5.01E-10	4.58E-10	3.49E-10	2.49E-10	1.74E-10	3.04E-11*
1.0E+04	4.50E-10	4.75E-10	4.24E-10	3.31E-10	2.44E-10	1.75E-10	3.34E-11*
2.0E+04	4.33E-10	4.12E-10	3.51E-10	2.83E-10	2.22E-10	1.71E-10	4.25E-11*

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-4.35175E+01	1.70837E-01	8.90710E-02	3.52059E-03	-3.19243E-02	-2.56283E-02	-9.06765E-03
20000.	-4.26087E+01	-1.54555E-01	-8.94632E-02	-4.26611E-02	-1.61171E-02	-3.24065E-03	7.74472E-04
40000.	-4.27112E+01	-2.07196E-01	-1.26305E-01	-5.98268E-02	-2.07064E-02	-3.39655E-03	1.43741E-03
60000.	-4.34961E+01	-8.32780E-02	-6.90413E-02	-4.75586E-02	-2.57998E-02	-1.00636E-02	-1.67943E-03
80000.	-4.42403E+01	-2.77368E-02	-2.78472E-02	-2.36213E-02	-1.56284E-02	-8.38900E-03	-4.20444E-03
100000.	-4.49754E+01	-4.83485E-04	1.15859E-02	4.48823E-03	-4.64650E-03	-6.95457E-03	-5.16220E-03
200000.	-4.86802E+01	2.01513E-01	1.40611E-01	7.66020E-02	3.17782E-02	8.85564E-03	7.47137E-05

See appendix for Chebyshev fit details.

$He + H^+ \rightarrow H(2p) + He^+$

Beam – Maxwellian



Line Emission Cross Sections of Electron Capture Collisions for
$H^+ + He \rightarrow H(H_\alpha) + He^+$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.3E+03	5.01E+07	5.94E-21
2.0E+03	6.21E+07	1.62E-20
4.0E+03	8.79E+07	6.73E-20
7.0E+03	1.16E+08	2.14E-19
1.0E+04	1.39E+08	4.29E-19
2.0E+04	1.96E+08	1.45E-18
4.0E+04	2.78E+08	2.92E-18
7.1E+04	3.70E+08	1.90E-18
1.0E+05	4.39E+08	1.13E-18
2.0E+05	6.21E+08	1.74E-19
3.0E+05	7.61E+08	4.13E-20

References: 122, 134, 136, 136, 156, 160, 192

Accuracy: 40%

Notes: (1) H Balmer-α emission is H_α with λ=656.3 nm.
(2) Balmer-α emission is the n=3->2 transition and is equal to $\sigma(3s)$ + .118 $\sigma(3p)$ + $\sigma(3d)$ where $\sigma(31)$ is the cross section for capture into each n=3 angular momentum state. The factor .118 is for the branching ratio of the 3p state.
(3) If the cross sections from pages B-40, B-46 and B-52 are summed using this formula then the H_α cross sections are within 10-20% of the evaluated sums.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 1.3E+03 eV/amu, E_{max} = 3.0E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-86.7431	1.47945	-2.26092	-.605332	.0536620	.125835	-.00401562	-.0314533	-.0205645

The fit represents the above cross sections with an rms deviation of 2.4%.
The maximum deviation is 5.5% at 7.1E+04 eV/amu.
See appendix for Chebyshev fit details.

$$H^+ + He \rightarrow H(H_\alpha) + He^+$$

Cross Section vs. Energy



Line Emission Rate Coefficients of Electron Capture Collisions for
$H^+ + He \rightarrow H(H_\alpha) + He^+$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	Equal Temp.	He Temp. (eV) 100.	500.	1000.	5000.	10000.	15000.	20000.
1.0E+02	4.62E-17	4.62E-17	4.63E-15	4.11E-14	2.61E-12	1.33E-11	3.27E-11	5.90E-11
2.0E+02	8.39E-15	4.63E-15	3.00E-14	1.07E-13	3.14E-12	1.45E-11	3.45E-11	6.13E-11
4.0E+02	1.54E-13	8.75E-14	1.81E-13	3.53E-13	4.36E-12	1.72E-11	3.84E-11	6.61E-11
7.0E+02	8.25E-13	4.86E-13	7.01E-13	1.03E-12	6.62E-12	2.16E-11	4.46E-11	7.34E-11
1.0E+03	2.14E-12	1.27E-12	1.62E-12	2.14E-12	9.40E-12	2.64E-11	5.11E-11	8.10E-11
2.0E+03	1.21E-11	7.26E-12	8.18E-12	9.40E-12	2.24E-11	4.56E-11	7.47E-11	1.07E-10
4.0E+03	5.67E-11	3.60E-11	3.79E-11	4.04E-11	6.25E-11	9.38E-11	1.27E-10	1.61E-10
7.0E+03	1.54E-10	1.08E-10	1.10E-10	1.14E-10	1.41E-10	1.75E-10	2.07E-10	2.38E-10
1.0E+04	2.49E-10	1.88E-10	1.91E-10	1.94E-10	2.20E-10	2.49E-10	2.77E-10	3.02E-10
2.0E+04	4.27E-10	3.80E-10	3.81E-10	3.83E-10	3.94E-10	4.07E-10	4.18E-10	4.27E-10

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+02 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
100.	-5.56271E+01	7.54225E+00	-1.58664E+00	3.76818E-01	-2.66300E-01	4.31633E-02	1.35706E-02
500.	-5.35452E+01	5.80636E+00	-5.23960E-01	-1.37429E-01	-6.79972E-02	-9.79835E-03	1.77516E-02
1000.	-5.22967E+01	4.84928E+00	-6.69532E-02	-2.78274E-01	-4.61871E-02	-1.88369E-03	9.07721E-03
5000.	-4.89693E+01	2.71099E+00	4.02894E-01	-2.04786E-01	-8.92702E-02	1.65791E-03	8.93841E-03
10000.	-4.72849E+01	1.82717E+00	3.81313E-01	-1.07442E-01	-7.83908E-02	-9.54437E-03	6.43829E-03
15000.	-4.62748E+01	1.34714E+00	3.25406E-01	-6.01114E-02	-6.20496E-02	-1.28087E-02	3.45512E-03
20000.	-4.55796E+01	1.03752E+00	2.73271E-01	-3.47245E-02	-4.87660E-02	-1.29500E-02	1.49966E-03
Equal Temp.	-5.48859E+01	7.46410E+00	-1.79561E+00	4.52870E-01	-3.47418E-01	1.03439E-01	-8.24563E-03

See appendix for Chebyshev fit details.

$$H^+ + He \rightarrow H(H_\alpha) + He^+$$

Maxwellian – Maxwellian



Line Emission Rate Coefficients of Electron Capture Collisions for
$He + H^+ \rightarrow H(H_\alpha) + He^+$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	60000.	80000.	100000.	200000.
1.0E+00	5.96E-11*	2.84E-10*	8.07E-10*	7.33E-10*	6.23E-10*	4.95E-10*	1.08E-10**
2.0E+00	5.96E-11*	2.84E-10*	8.06E-10*	7.33E-10*	6.23E-10*	4.95E-10*	1.08E-10**
4.0E+00	5.96E-11*	2.83E-10*	8.03E-10*	7.33E-10*	6.23E-10*	4.94E-10*	1.08E-10**
7.0E+00	5.97E-11*	2.83E-10*	8.01E-10*	7.33E-10*	6.23E-10*	4.94E-10*	1.08E-10**
1.0E+01	5.98E-11*	2.83E-10*	7.99E-10	7.33E-10*	6.23E-10*	4.93E-10*	1.08E-10**
2.0E+01	6.01E-11*	2.82E-10	7.94E-10	7.33E-10*	6.23E-10*	4.92E-10*	1.08E-10**
4.0E+01	6.08E-11*	2.82E-10	7.87E-10	7.33E-10	6.23E-10*	4.90E-10*	1.07E-10**
7.0E+01	6.18E-11*	2.82E-10	7.80E-10	7.33E-10	6.23E-10*	4.88E-10*	1.07E-10**
1.0E+02	6.29E-11*	2.83E-10	7.74E-10	7.33E-10	6.23E-10*	4.87E-10*	1.07E-10**
2.0E+02	6.67E-11	2.85E-10	7.60E-10	7.33E-10	6.23E-10	4.84E-10*	1.07E-10**
4.0E+02	7.45E-11	2.92E-10	7.42E-10	7.31E-10	6.21E-10	4.79E-10*	1.07E-10**
7.0E+02	8.65E-11	3.04E-10	7.24E-10	7.27E-10	6.17E-10	4.75E-10*	1.07E-10**
1.0E+03	9.83E-11	3.16E-10	7.11E-10	7.22E-10	6.12E-10	4.72E-10*	1.07E-10**
2.0E+03	1.36E-10	3.52E-10	6.80E-10	7.02E-10	5.97E-10	4.64E-10*	1.09E-10*
4.0E+03	2.05E-10	4.00E-10	6.43E-10	6.60E-10	5.69E-10	4.51E-10	1.12E-10*
7.0E+03	2.87E-10	4.40E-10	6.04E-10	6.08E-10	5.33E-10	4.33E-10	1.17E-10*
1.0E+04	3.46E-10	4.63E-10	5.75E-10	5.68E-10	5.02E-10	4.16E-10	1.22E-10*
2.0E+04	4.46E-10	4.86E-10	5.07E-10	4.79E-10	4.27E-10	3.67E-10	1.33E-10*

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-4.58848E+01	9.71807E-01	4.83279E-01	9.11138E-02	-6.12690E-02	-5.76723E-02	-1.79106E-02
20000.	-4.36507E+01	2.64211E-01	1.43555E-01	2.70221E-02	-2.48764E-02	-2.40082E-02	-7.54802E-03
40000.	-4.21414E+01	-2.00867E-01	-8.98633E-02	-2.87027E-02	-8.54506E-03	-3.26747E-03	-1.21561E-03
60000.	-4.22369E+01	-1.51900E-01	-1.08731E-01	-5.90440E-02	-2.19357E-02	-3.21591E-03	2.30144E-03
80000.	-4.25414E+01	-1.32445E-01	-9.43661E-02	-5.17183E-02	-2.05460E-02	-5.15132E-03	-3.15003E-04
100000.	-4.29896E+01	-1.10475E-01	-6.22264E-02	-3.20171E-02	-1.65763E-02	-7.76766E-03	-2.70010E-03
200000.	-4.58241E+01	7.16963E-02	5.83098E-02	3.32118E-02	1.21459E-02	1.47440E-03	-1.87694E-03

See appendix for Chebyshev fit details.

$$He + H^+ \rightarrow H(H_\alpha) + He^+$$

Beam – Maxwellian



Electron Capture Cross Sections for
$H^+ + He \rightarrow H(3s) + He^+$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.2E+03	4.81E+07	7.90E-20
2.0E+03	6.21E+07	9.72E-20
4.0E+03	8.79E+07	1.39E-19
7.0E+03	1.16E+08	1.94E-19
1.0E+04	1.39E+08	2.59E-19
2.0E+04	1.96E+08	6.57E-19
4.0E+04	2.78E+08	1.65E-18
4.4E+04	2.91E+08	1.67E-18
7.0E+04	3.68E+08	1.20E-18
1.0E+05	4.39E+08	7.58E-19
2.0E+05	6.21E+08	1.65E-19
4.0E+05	8.78E+08	1.10E-20
7.0E+05	1.16E+09	1.16E-21
1.0E+06	1.39E+09	2.40E-22
1.3E+06	1.58E+09	6.65E-23

References: 20, 129, 130, 134, 154, 155, 156, 157, 158, 159, 160, 161

Accuracy: unknown

Notes: (1) Very few papers report absolute cross sections. The overall assessment of the data is in doubt due to the validity of the different normalization schemes.
(2) Cascade from higher states is usually neglected and can contribute up to a 5% error in the reported cross sections.
(3) The cross section (σ_i) for capture into state i is determined by measuring the emission cross section (σ_{ij}) for a transition from state i to j, correcting for any branching and correcting for cascading from upper states. For Balmer-α emission from the n=3 state, the cross section $\sigma(H_\alpha)=\sigma(3s) + .188\ \sigma(3p) + \sigma(3d)$.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 1.2E+03 eV/amu, E_{max} = 1.3E+06 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-89.3963	-2.75333	-3.10976	-.952686	.352124	.248416	-.119805	-.0706000	.0607563

The fit represents the above cross sections with an rms deviation of 7.1%.
The maximum deviation is 12.3% at 7.0E+04 eV/amu.
See appendix for Chebyshev fit details.

$H^+ + He \rightarrow H(3s) + He^+$



Electron Capture Rate Coefficients for
$H^+ + He \rightarrow H(3s) + He^+$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	Equal Temp.	He Temp. (eV) 100.	500.	1000.	5000.	10000.	15000.	20000.
1.0E+02	1.04E-15	1.04E-15	6.14E-14	3.76E-13	5.53E-12	1.30E-11	2.28E-11	3.55E-11
2.0E+02	1.02E-13	6.14E-14	2.92E-13	7.83E-13	6.07E-12	1.36E-11	2.37E-11	3.67E-11
4.0E+02	1.02E-12	6.72E-13	1.14E-12	1.78E-12	7.17E-12	1.51E-11	2.56E-11	3.90E-11
7.0E+02	2.98E-12	2.17E-12	2.71E-12	3.38E-12	8.87E-12	1.73E-11	2.86E-11	4.26E-11
1.0E+03	4.99E-12	3.78E-12	4.32E-12	4.99E-12	1.07E-11	1.97E-11	3.17E-11	4.63E-11
2.0E+03	1.23E-11	9.31E-12	9.91E-12	1.07E-11	1.77E-11	2.91E-11	4.32E-11	5.94E-11
4.0E+03	3.44E-11	2.44E-11	2.54E-11	2.66E-11	3.72E-11	5.27E-11	6.98E-11	8.78E-11
7.0E+03	8.42E-11	5.98E-11	6.12E-11	6.29E-11	7.70E-11	9.50E-11	1.13E-10	1.30E-10
1.0E+04	1.37E-10	1.03E-10	1.04E-10	1.06E-10	1.20E-10	1.37E-10	1.53E-10	1.68E-10
2.0E+04	2.48E-10	2.17E-10	2.17E-10	2.18E-10	2.26E-10	2.34E-10	2.42E-10	2.48E-10

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+02 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
100.	-5.37012E+01	5.51736E+00	-1.26631E+00	6.41819E-01	-2.59717E-01	-3.50690E-02	-8.79675E-04
500.	-5.19390E+01	4.01666E+00	-2.93656E-01	1.47807E-01	-7.57652E-02	-7.86568E-02	3.19866E-03
1000.	-5.09725E+01	3.25191E+00	1.17635E-01	-4.41960E-03	-5.45641E-02	-6.54437E-02	-4.41120E-03
5000.	-4.89296E+01	1.94263E+00	4.86039E-01	-5.63401E-02	-8.77510E-02	-3.17734E-02	-1.68441E-04
10000.	-4.79520E+01	1.50755E+00	4.27186E-01	-3.56552E-02	-7.51944E-02	-2.47079E-02	1.68707E-03
15000.	-4.72456E+01	1.22338E+00	3.57663E-01	-2.27359E-02	-5.99425E-02	-2.00776E-02	7.63417E-04
20000.	-4.66818E+01	1.00201E+00	2.99649E-01	-1.36330E-02	-4.72262E-02	-1.65699E-02	-1.65257E-05
Equal Temp.	-5.31267E+01	5.47937E+00	-1.36275E+00	6.93916E-01	-3.79924E-01	1.82789E-02	-3.07574E-03

See appendix for Chebyshev fit details.

$H^+ + He \rightarrow H(3s) + He^+$

Maxwellian — Maxwellian



Electron Capture Rate Coefficients for
$He + H^+ \rightarrow H(3s) + He^+$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	3.61E-11	1.29E-10	4.57E-10	4.40E-10	3.32E-10	1.02E-10	4.40E-12*
2.0E+00	3.62E-11	1.29E-10	4.56E-10	4.40E-10	3.32E-10	1.02E-10	4.40E-12*
4.0E+00	3.62E-11	1.29E-10	4.55E-10	4.40E-10	3.32E-10	1.02E-10	4.40E-12*
7.0E+00	3.63E-11	1.29E-10	4.54E-10	4.40E-10	3.32E-10	1.02E-10	4.40E-12*
1.0E+01	3.64E-11	1.29E-10	4.54E-10	4.39E-10	3.31E-10	1.02E-10	4.40E-12*
2.0E+01	3.67E-11	1.30E-10	4.52E-10	4.39E-10	3.31E-10	1.02E-10	4.41E-12*
4.0E+01	3.72E-11	1.30E-10	4.50E-10	4.38E-10	3.30E-10	1.01E-10	4.41E-12*
7.0E+01	3.78E-11	1.31E-10	4.47E-10	4.37E-10	3.29E-10	1.01E-10	4.41E-12*
1.0E+02	3.83E-11	1.32E-10	4.44E-10	4.36E-10	3.28E-10	1.01E-10	4.41E-12*
2.0E+02	4.01E-11	1.35E-10	4.38E-10	4.33E-10	3.26E-10	9.99E-11	4.42E-12
4.0E+02	4.35E-11	1.40E-10	4.29E-10	4.30E-10	3.24E-10	9.92E-11	4.44E-12
7.0E+02	4.86E-11	1.49E-10	4.19E-10	4.27E-10	3.21E-10	9.86E-11	4.47E-12
1.0E+03	5.37E-11	1.57E-10	4.12E-10	4.24E-10	3.19E-10	9.82E-11	4.50E-12
2.0E+03	7.17E-11	1.82E-10	3.96E-10	4.14E-10	3.14E-10	9.78E-11	4.60E-12
4.0E+03	1.08E-10	2.16E-10	3.78E-10	3.93E-10	3.06E-10	9.84E-11	4.81E-12
7.0E+03	1.56E-10	2.47E-10	3.59E-10	3.66E-10	2.93E-10	1.00E-10	5.14E-12
1.0E+04	1.93E-10	2.66E-10	3.46E-10	3.44E-10	2.82E-10	1.02E-10	5.51E-12
2.0E+04	2.61E-10	2.90E-10	3.13E-10	2.94E-10	2.48E-10	1.06E-10	7.04E-12

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-4.69590E+01	9.17776E-01	4.84822E-01	1.28148E-01	-3.38227E-02	-5.58964E-02	-2.93167E-02
20000.	-4.50524E+01	3.95677E-01	1.97669E-01	3.56569E-02	-3.01554E-02	-2.88658E-02	-8.74226E-03
40000.	-4.32289E+01	-1.65074E-01	-7.60355E-02	-2.21081E-02	-4.24972E-03	-1.82033E-03	-1.77544E-03
70000.	-4.32620E+01	-1.48289E-01	-9.51993E-02	-4.97051E-02	-2.06125E-02	-5.32942E-03	1.01964E-03
100000.	-4.37786E+01	-1.05527E-01	-6.21604E-02	-3.28418E-02	-1.67902E-02	-7.71384E-03	-2.68575E-03
200000.	-4.60232E+01	3.14283E-05	2.12471E-02	1.95060E-02	8.72740E-03	1.21515E-03	-1.62989E-03
500000.	-5.21281E+01	1.52077E-01	1.10093E-01	6.67186E-02	3.41707E-02	1.49151E-02	5.83661E-03

See appendix for Chebyshev fit details.

$He + H^+ \rightarrow H(3s) + He^+$

Beam – Maxwellian



Electron Capture Cross Sections for
$H^+ + He \rightarrow H(3p) + He^+$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.7E+03	5.73E+07	1.35E-19
2.0E+03	6.21E+07	1.48E-19
4.0E+03	8.79E+07	2.45E-19
7.0E+03	1.16E+08	3.84E-19
1.0E+04	1.39E+08	5.13E-19
2.0E+04	1.96E+08	8.57E-19
3.6E+04	2.64E+08	1.14E-18
4.0E+04	2.78E+08	1.14E-18
7.0E+04	3.68E+08	7.28E-19
1.0E+05	4.39E+08	3.49E-19
2.0E+05	6.21E+08	3.95E-20
4.0E+05	8.78E+08	1.75E-21
7.0E+05	1.16E+09	1.02E-22
1.0E+06	1.39E+09	1.44E-23
1.2E+06	1.52E+09	4.95E-24

References: 134, 136, 156, 157, 159, 160, 163, 164, 165, 166, 167

Accuracy: Unknown

Notes: (1) Very few papers report absolute cross sections. The overall assessment of the data is in doubt due to the validity of the different normalization schemes.
(2) Cascade from higher states is usually neglected but can contribute up to 5% error in the reported cross section measurements.
(3) The cross section (σ_i) for capture into state i is determined by measuring the emission cross section (σ_{ij}) for a transition from state i to j, correcting for cascading from upper states and correcting for any branching. For Balmer-α emission from the n=3 state the cross section $\sigma(H_\alpha)=\sigma(3s) + .118\sigma(3p) + \sigma(3d)$.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 1.7E+03 eV/amu, E_{max} = 1.2E+06 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-90.6049	-4.52585	-3.52507	-.678448	.325826	.143519	-.0657813	-.0500666	.0154546

The fit represents the above cross sections with an rms deviation of 1.7%.
The maximum deviation is 3.6% at 2.0E+04 eV/amu.
See appendix for Chebyshev fit details.

$H^+ + He \rightarrow H(3p) + He^+$

Cross Section vs. Energy



Electron Capture Rate Coefficients for
$H^+ + He \rightarrow H(3p) + He^+$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	Equal Temp.	He Temp. (eV) 100.	500.	1000.	5000.	10000.	15000.	20000.
1.0E+02	4.48E-17	4.48E-17	1.54E-14	2.06E-13	8.41E-12	2.24E-11	3.77E-11	5.37E-11
2.0E+02	3.19E-14	1.54E-14	1.44E-13	5.85E-13	9.46E-12	2.36E-11	3.90E-11	5.49E-11
4.0E+02	8.46E-13	4.71E-13	9.92E-13	1.85E-12	1.16E-11	2.60E-11	4.15E-11	5.75E-11
7.0E+02	3.74E-12	2.44E-12	3.29E-12	4.43E-12	1.49E-11	2.96E-11	4.54E-11	6.14E-11
1.0E+03	7.38E-12	5.15E-12	6.13E-12	7.38E-12	1.83E-11	3.34E-11	4.92E-11	6.52E-11
2.0E+03	2.12E-11	1.57E-11	1.69E-11	1.83E-11	3.02E-11	4.60E-11	6.20E-11	7.77E-11
4.0E+03	5.24E-11	3.99E-11	4.12E-11	4.28E-11	5.56E-11	7.15E-11	8.68E-11	1.01E-10
7.0E+03	9.84E-11	7.80E-11	7.93E-11	8.08E-11	9.26E-11	1.07E-10	1.19E-10	1.31E-10
1.0E+04	1.35E-10	1.12E-10	1.13E-10	1.14E-10	1.24E-10	1.35E-10	1.45E-10	1.53E-10
2.0E+04	1.90E-10	1.78E-10	1.78E-10	1.79E-10	1.82E-10	1.85E-10	1.88E-10	1.90E-10

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+02 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
100.	-5.48603E+01	6.73175E+00	-2.33750E+00	8.57032E-01	-2.92258E-01	8.89193E-03	1.19267E-02
500.	-5.23540E+01	4.57847E+00	-9.34127E-01	1.52574E-01	-2.98508E-02	-5.41726E-02	1.56047E-02
1000.	-5.09960E+01	3.48139E+00	-3.33404E-01	-6.24733E-02	4.55378E-04	-3.70467E-02	2.75925E-03
5000.	-4.83469E+01	1.64909E+00	2.61394E-01	-1.05216E-01	-5.38273E-02	-7.49908E-03	7.09196E-04
10000.	-4.73425E+01	1.12338E+00	2.55051E-01	-5.63104E-02	-5.16563E-02	-1.13825E-02	1.38299E-03
15000.	-4.67459E+01	8.47733E-01	2.16661E-01	-3.36664E-02	-4.24929E-02	-1.19700E-02	4.13866E-04
20000.	-4.63234E+01	6.64850E-01	1.80257E-01	-2.17729E-02	-3.47781E-02	-1.13742E-02	-1.56934E-04
Equal Temp.	-5.42194E+01	6.56075E+00	-2.45975E+00	9.87265E-01	-4.35539E-01	8.21003E-02	-9.92180E-03

See appendix for Chebyshev fit details.

$H^+ + He \rightarrow H(3p) + He^+$

Maxwellian — Maxwellian



Electron Capture Rate Coefficients for
$He + H^+ \rightarrow H(3p) + He^+$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	7.13E-11	1.68E-10	3.16E-10	2.67E-10	1.53E-10	2.45E-11	5.53E-13*
2.0E+00	7.13E-11	1.68E-10	3.16E-10	2.67E-10	1.53E-10	2.45E-11	5.53E-13*
4.0E+00	7.13E-11	1.68E-10	3.15E-10	2.66E-10	1.53E-10	2.45E-11	5.53E-13*
7.0E+00	7.13E-11	1.68E-10	3.15E-10	2.66E-10	1.53E-10	2.44E-11	5.53E-13*
1.0E+01	7.13E-11	1.68E-10	3.14E-10	2.65E-10	1.52E-10	2.44E-11	5.53E-13*
2.0E+01	7.15E-11	1.68E-10	3.14E-10	2.64E-10	1.52E-10	2.44E-11	5.54E-13*
4.0E+01	7.17E-11	1.68E-10	3.12E-10	2.63E-10	1.52E-10	2.43E-11	5.54E-13*
7.0E+01	7.21E-11	1.68E-10	3.11E-10	2.62E-10	1.51E-10	2.43E-11	5.55E-13
1.0E+02	7.26E-11	1.68E-10	3.09E-10	2.61E-10	1.51E-10	2.42E-11	5.55E-13
2.0E+02	7.41E-11	1.69E-10	3.06E-10	2.58E-10	1.50E-10	2.42E-11	5.57E-13
4.0E+02	7.71E-11	1.71E-10	3.01E-10	2.54E-10	1.49E-10	2.42E-11	5.61E-13
7.0E+02	8.16E-11	1.74E-10	2.95E-10	2.50E-10	1.49E-10	2.42E-11	5.68E-13
1.0E+03	8.61E-11	1.77E-10	2.90E-10	2.47E-10	1.49E-10	2.43E-11	5.74E-13
2.0E+03	1.00E-10	1.86E-10	2.79E-10	2.39E-10	1.48E-10	2.49E-11	5.97E-13
4.0E+03	1.24E-10	1.96E-10	2.63E-10	2.25E-10	1.46E-10	2.63E-11	6.43E-13
7.0E+03	1.51E-10	2.04E-10	2.45E-10	2.08E-10	1.42E-10	2.88E-11	7.17E-13
1.0E+04	1.70E-10	2.07E-10	2.30E-10	1.95E-10	1.38E-10	3.12E-11	8.02E-13
2.0E+04	1.95E-10	2.03E-10	1.97E-10	1.64E-10	1.24E-10	3.75E-11	1.18E-12

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-4.61593E+01	4.69768E-01	2.55845E-01	6.75806E-02	-2.06018E-02	-3.21005E-02	-1.71678E-02
20000.	-4.48832E+01	1.05388E-01	5.13750E-02	1.87880E-03	-1.77315E-02	-1.43595E-02	-5.26979E-03
40000.	-4.39844E+01	-1.88716E-01	-1.03793E-01	-4.31001E-02	-1.44695E-02	-4.32136E-03	-8.32803E-04
70000.	-4.43219E+01	-1.88451E-01	-1.07258E-01	-4.99127E-02	-2.00817E-02	-6.25364E-03	-3.66359E-04
100000.	-4.52889E+01	-7.00699E-02	-4.15959E-02	-2.66009E-02	-1.74699E-02	-9.19908E-03	-3.00670E-03
200000.	-4.87156E+01	1.42634E-01	1.14850E-01	6.70006E-02	2.67000E-02	5.12084E-03	-2.83726E-03
500000.	-5.61687E+01	2.49112E-01	1.78331E-01	1.05979E-01	5.28963E-02	2.25156E-02	8.83834E-03

See appendix for Chebyshev fit details.

$He + H^+ \rightarrow H(3p) + He^+$

Beam – Maxwellian



Electron Capture Cross Sections for
$H^+ + He \rightarrow H(3d) + He^+$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
3.0E+03	7.61E+07	1.16E-19
4.0E+03	8.79E+07	1.36E-19
7.0E+03	1.16E+08	1.65E-19
1.0E+04	1.39E+08	1.71E-19
2.0E+04	1.96E+08	1.43E-19
4.0E+04	2.78E+08	8.33E-20
7.0E+04	3.68E+08	3.95E-20
1.0E+05	4.39E+08	1.53E-20
2.0E+05	6.21E+08	1.44E-21
2.7E+05	7.22E+08	4.00E-22
3.2E+05	7.86E+08	1.67E-22

References: 134, 136, 156, 157, 158, 159, 160

Accuracy: Unknown

Notes: (1) Very few papers report the absolute cross sections. The overall assessment of the data is in doubt due to the validity of the different normalization schemes.
(2) Cascade from higher states is usually neglected, but can contribute up to a 5% error in the reported cross section.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 3.0E+03 eV/amu, E_{max} = 3.2E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-90.4702	-2.99013	-1.60913	-.296858	-.0140200	.0353005	.0153108	-.0221875	-.0321765

The fit represents the above cross sections with an rms deviation of 1.9%.
The maximum deviation is 3.1% at 1.0E+05 eV/amu.
See appendix for Chebyshev fit details.

$H^+ + He \rightarrow H(3d) + He^+$



Electron Capture Rate Coefficients for
$H^+ + He \rightarrow H(3d) + He^+$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	Equal Temp.	He Temp. (eV) 100.	500.	1000.	5000.	10000.	15000.	20000.
2.0E+02	2.41E-16	6.60E-17	3.49E-15	4.08E-14	3.27E-12	8.54E-12	1.24E-11	1.52E-11
4.0E+02	7.71E-14	2.80E-14	1.01E-13	2.88E-13	4.19E-12	9.26E-12	1.29E-11	1.55E-11
7.0E+02	8.89E-13	4.53E-13	7.29E-13	1.15E-12	5.53E-12	1.03E-11	1.36E-11	1.60E-11
1.0E+03	2.38E-12	1.43E-12	1.84E-12	2.38E-12	6.80E-12	1.12E-11	1.43E-11	1.65E-11
2.0E+03	7.79E-12	5.85E-12	6.28E-12	6.80E-12	1.04E-11	1.37E-11	1.61E-11	1.78E-11
4.0E+03	1.48E-11	1.26E-11	1.29E-11	1.32E-11	1.52E-11	1.72E-11	1.85E-11	1.95E-11
7.0E+03	1.94E-11	1.78E-11	1.79E-11	1.80E-11	1.90E-11	1.98E-11	2.04E-11	2.08E-11
1.0E+04	2.10E-11	2.01E-11	2.02E-11	2.02E-11	2.06E-11	2.10E-11	2.12E-11	2.12E-11
2.0E+04	2.01E-11	2.09E-11	2.09E-11	2.09E-11	2.08E-11	2.05E-11	2.03E-11	2.01E-11

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 2.0E+02 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
100.	-5.63688E+01	5.53526E+00	-2.53031E+00	7.68047E-01	-2.09210E-01	3.02777E-02	8.99288E-04
500.	-5.46783E+01	4.07558E+00	-1.58318E+00	2.94914E-01	-2.76037E-02	-2.07878E-02	1.01392E-02
1000.	-5.34861E+01	3.08285E+00	-1.00328E+00	6.05624E-02	3.07528E-02	-2.37156E-02	4.91264E-03
5000.	-5.08092E+01	1.02925E+00	-1.15489E-01	-1.11263E-01	1.29497E-04	5.31512E-03	-1.44689E-03
10000.	-5.00260E+01	5.09612E-01	-1.91665E-02	-7.34488E-02	-1.56542E-02	2.56893E-03	7.98864E-04
15000.	-4.96848E+01	2.98178E-01	-7.68530E-03	-5.18119E-02	-1.65953E-02	-3.35169E-04	1.07168E-03
20000.	-4.94956E+01	1.81879E-01	-1.04948E-02	-3.99118E-02	-1.50920E-02	-1.49095E-03	1.11348E-03
Equal Temp.	-5.53820E+01	4.85397E+00	-2.35173E+00	7.57520E-01	-2.46645E-01	5.44226E-02	-8.31810E-03

See appendix for Chebyshev fit details.

$H^+ + He \rightarrow H(3d) + He^+$

Maxwellian — Maxwellian



Electron Capture Rate Coefficients for
$He + H^+ \rightarrow H(3d) + He^+$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	60000.	80000.	100000.	200000.
1.0E+00	2.37E-11	2.80E-11	2.31E-11	1.65E-11	1.09E-11	6.71E-12*	8.94E-13**
2.0E+00	2.37E-11	2.80E-11	2.31E-11	1.65E-11	1.09E-11	6.71E-12*	8.93E-13**
4.0E+00	2.37E-11	2.80E-11	2.31E-11	1.65E-11	1.09E-11	6.70E-12*	8.93E-13**
7.0E+00	2.36E-11	2.79E-11	2.30E-11	1.65E-11	1.09E-11	6.70E-12*	8.92E-13**
1.0E+01	2.36E-11	2.79E-11	2.30E-11	1.65E-11	1.09E-11	6.70E-12*	8.92E-13**
2.0E+01	2.36E-11	2.78E-11	2.30E-11	1.65E-11	1.09E-11	6.69E-12	8.91E-13**
4.0E+01	2.35E-11	2.77E-11	2.29E-11	1.65E-11	1.09E-11	6.68E-12	8.90E-13**
7.0E+01	2.34E-11	2.76E-11	2.28E-11	1.65E-11	1.09E-11	6.67E-12	8.89E-13*
1.0E+02	2.34E-11	2.75E-11	2.28E-11	1.65E-11	1.09E-11	6.67E-12	8.89E-13*
2.0E+02	2.33E-11	2.73E-11	2.26E-11	1.65E-11	1.10E-11	6.66E-12	8.89E-13*
4.0E+02	2.31E-11	2.70E-11	2.24E-11	1.65E-11	1.10E-11	6.68E-12	8.92E-13*
7.0E+02	2.30E-11	2.66E-11	2.22E-11	1.64E-11	1.10E-11	6.72E-12	8.98E-13*
1.0E+03	2.29E-11	2.63E-11	2.20E-11	1.63E-11	1.09E-11	6.76E-12	9.05E-13*
2.0E+03	2.26E-11	2.55E-11	2.15E-11	1.59E-11	1.08E-11	6.90E-12	9.35E-13*
4.0E+03	2.24E-11	2.41E-11	2.05E-11	1.53E-11	1.06E-11	7.06E-12	1.01E-12*
7.0E+03	2.22E-11	2.26E-11	1.92E-11	1.45E-11	1.04E-11	7.19E-12	1.13E-12*
1.0E+04	2.18E-11	2.14E-11	1.80E-11	1.38E-11	1.01E-11	7.22E-12	1.26E-12*
2.0E+04	1.98E-11	1.83E-11	1.51E-11	1.20E-11	9.26E-12	7.06E-12	1.66E-12*

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-4.90115E+01	-6.26583E-02	-3.14715E-02	-1.74368E-02	-1.18746E-02	-8.00345E-03	-5.51328E-03
20000.	-4.87965E+01	-1.63847E-01	-9.40039E-02	-4.32626E-02	-1.65508E-02	-4.78447E-03	-6.28354E-04
40000.	-4.91665E+01	-1.54448E-01	-9.66541E-02	-5.08695E-02	-2.30319E-02	-8.09916E-03	-1.06973E-03
60000.	-4.97788E+01	-1.11497E-01	-7.93398E-02	-4.38613E-02	-1.80124E-02	-5.25241E-03	-1.28373E-03
80000.	-5.05382E+01	-4.92366E-02	-4.11913E-02	-2.64723E-02	-1.18809E-02	-3.95220E-03	-2.08141E-03
100000.	-5.14178E+01	3.35427E-02	2.01516E-02	3.56710E-04	-9.44387E-03	-8.21678E-03	-3.58171E-03
200000.	-5.52583E+01	2.09996E-01	1.57946E-01	9.13682E-02	3.94669E-02	1.12103E-02	-2.26555E-04

See appendix for Chebyshev fit details.

$He + H^+ \rightarrow H(3d) + He^+$



Electron Capture Cross Sections for

$H^+ + Li \rightarrow H(2s,2p) + Li^+$

H(2s) Energy eV/amu	H(2s) Cross Section cm^2	H(2p) Energy eV/amu	H(2p) Cross Section cm^2
9.6E+02	1.09E-15	9.8E+02	1.44E-15
1.9E+03	2.22E-15	2.0E+03	2.37E-15
2.4E+03	2.52E-15	2.9E+03	2.95E-15
2.9E+03	2.56E-15	3.4E+03	3.05E-15
4.8E+03	2.06E-15	4.9E+03	2.75E-15
6.7E+03	1.39E-15	6.8E+03	2.02E-15
8.6E+03	9.00E-16	9.7E+03	1.13E-15
9.5E+03	7.26E-16	1.4E+04	3.01E-16
1.4E+04	2.73E-16	1.9E+04	9.71E-17
1.9E+04	1.23E-16	2.4E+04	3.76E-17
2.4E+04	6.20E-17	2.9E+04	1.76E-17
3.8E+04	1.40E-17	3.9E+04	3.98E-18
4.9E+04	6.29E-18	4.9E+04	1.21E-18
5.9E+04	3.55E-18	5.9E+04	4.47E-19
6.8E+04	2.22E-18	6.9E+04	1.94E-19
8.8E+04	9.76E-19	8.9E+04	4.96E-20
9.8E+04	6.77E-19	1.1E+05	1.69E-20
1.3E+05	2.72E-19		

References: 255, 256, 257

Accuracy: $E < 2x10^4$ eV/amu - 60%; $E > 2x10^4$ eV/amu - unknown

Notes: (1) The cross section data for energies greater than $2x10^4$ eV/amu are theoretical results of Ermolaev (ref. 257); for excitation of the Li(2p) level by protons see ref. 258.
(2) Cascading from higher states is less than 5%, thus $\sigma(2p) \simeq \sigma(Ly_\alpha)$.

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

H(2s) E_{min} = 9.55E+02 eV/amu, E_{max} = 1.3E+05 eV/amu
H(2p) E_{min} = 9.8E+02 eV/amu, E_{max} = 1.1E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

	A0	A1	A2	A3	A4	A5	A6	A7	A8
H(2s)	-73.9306	-4.42348	-1.68271	.337287	.0753407	-.0624317	-.0223686	6.11347E-04	-.00573178
H(2p)	-75.0922	-5.75721	-2.57021	.0939967	.255624	-.0329880	-.0184241	.0249338	.0333479

The fit represents the H(2s) cross section with an rms deviation of 0.9%.
The maximum deviation is 1.7% at 4.8E+04 eV/amu.

The fit represents the H(2p) cross section with an rms deviation of 3.2%.
The maximum deviation is 5.5% at 1.9E+04 eV/amu.

See appendix for Chebyshev fit details.

$H^+ + Li \rightarrow H(2s,2p) + Li^+$

Cross Section vs. Energy



Electron Capture Cross Sections for

$H^+ + Li \rightarrow H(3s,3p,3d) + Li^+$

(3s) Energy eV/amu	(3s) Cross Section cm^2	(3p) Energy eV/amu	(3p) Cross Section cm^2	(3d) Energy eV/amu	(3d) Cross Section cm^2
1.3E+03	7.96E-17	9.8E+02	1.28E-16	3.1E+03	7.47E-17
2.0E+03	9.60E-17	1.5E+03	1.51E-16	3.9E+03	8.78E-17
2.9E+03	1.23E-16	2.0E+03	1.76E-16	4.8E+03	9.79E-17
3.9E+03	1.55E-16	2.9E+03	2.33E-16	6.8E+03	1.06E-16
4.9E+03	2.07E-16	3.9E+03	3.17E-16	9.7E+03	8.39E-17
6.0E+03	3.46E-16	4.9E+03	3.65E-16	1.3E+04	4.31E-17
6.9E+03	4.43E-16	6.9E+03	3.45E-16	1.5E+04	2.78E-17
7.9E+03	4.55E-16	8.9E+03	2.81E-16	2.0E+04	1.16E-17
8.9E+03	4.29E-16	9.9E+03	2.47E-16	2.9E+04	2.97E-18
1.0E+04	3.83E-16	1.5E+04	1.23E-16	3.9E+04	1.11E-18
1.5E+04	1.64E-16	2.0E+04	5.12E-17	4.9E+04	4.65E-19
2.0E+04	6.88E-17	2.5E+04	2.18E-17	7.0E+04	1.44E-19
2.5E+04	3.15E-17	3.0E+04	1.05E-17	9.9E+04	3.63E-20
3.5E+04	7.97E-18	3.5E+04	6.04E-18	1.3E+05	1.20E-20
4.9E+04	1.96E-18	4.5E+04	2.25E-18	1.5E+05	7.57E-21
6.0E+04	1.03E-18	5.0E+04	1.43E-18		
8.0E+04	3.69E-19	6.0E+04	6.60E-19		
1.0E+05	1.81E-19	7.0E+04	3.86E-19		
1.3E+05	8.92E-20	9.0E+04	1.48E-19		
		1.0E+05	9.25E-20		
		1.3E+05	3.87E-20		

References: 256, 257

Accuracy: $E < 2x10^4$ eV/amu - 60%; $E > 2x10^4$ eV/amu - unknown

Notes: (1) The cross section data for energies greater than $2x10^4$ eV/amu are the theoretical results of Ermolaev (ref. 257).

(2) The cross section (σ_i) for capture into state i is determined by measuring the emission cross section (σ_{ij}) for a transition from state i to j, correcting for any branching from state i and correcting for cascading from upper states.

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

(3s) E_{min} = 1.3E+03 eV/amu, E_{max} = 1.2E+05 eV/amu
(3p) E_{min} = 9.8E+02 eV/amu, E_{max} = 1.2E+05 eV/amu
(3d) E_{min} = 3.1E+03 eV/amu, E_{max} = 1.5E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

	A0	A1	A2	A3	A4	A5	A6	A7	A8
(3s)	-77.4321	-3.51092	-2.10742	.102860	.543346	-.0941225	-.179427	.114301	-.00463843
(3p)	-77.4555	-4.03388	-2.19950	.00628310	.364180	-.0330367	-.0799875	.00340643	-.00766823
(3d)	-80.9668	-4.92271	-1.27692	.391264	-.0254482	-.0874189	.0722031	.0109053	-.0235328

The fit represents the (3s) cross section with an rms deviation of 6.8%.
The maximum deviation is 19.4% at 4.9E+03 eV/amu.

The fit represents the (3p) cross section with an rms deviation of 3.7%.
The maximum deviation is 6.6% at 3.0E+04 eV/amu.

The fit represents the (3d) cross section with an rms deviation of 3.2%.
The maximum deviation is 6.5% at 4.9E+04 eV/amu.

See appendix for Chebyshev fit details.

$H^+ + Li \rightarrow H(3s,3p,3d) + Li^+$

Cross Section vs. Energy



Electron Capture Cross Sections for
$He^+ + H_2 \rightarrow He(2^1S + 2^3S)$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.5E+03	6.95E+07	6.75E-18
3.0E+03	7.61E+07	9.57E-18
4.0E+03	8.79E+07	1.58E-17
5.0E+03	9.82E+07	2.18E-17
7.0E+03	1.16E+08	3.14E-17
9.0E+03	1.32E+08	4.14E-17
1.0E+04	1.39E+08	4.63E-17
1.5E+04	1.70E+08	6.54E-17
2.0E+04	1.96E+08	7.00E-17
2.5E+04	2.20E+08	6.69E-17
3.5E+04	2.60E+08	5.28E-17
4.5E+04	2.95E+08	4.03E-17
5.0E+04	3.11E+08	3.52E-17
6.0E+04	3.40E+08	2.75E-17
6.4E+04	3.51E+08	2.48E-17

References: 98, 102, 196, 197, 198

Accuracy: 40%

Note: The cross sections are for capture into the metastable states 2^1S and 2^3S.

For Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 2.5E+03 eV/amu, E_{max} = 6.4E+04 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-76.2905	.717609	-.746846	-.0903391	1.71246E-03	.0416862	-7.59576E-04	-.0179523	6.07163E-03

The fit represents the above cross sections with an rms deviation of 0.5%.
The maximum deviation is 0.9% at 4.0E+03 eV/amu.
See appendix for Chebyshev fit details.

$$He^+ + H_2 \rightarrow He(2^1S + 2^3S)$$

Cross Section vs. Energy



Electron Capture Cross Sections for

$He^+ + H_2 \rightarrow He(2^1P, 3^1P)$

(2^1P) Energy eV/amu	Cross Section cm^2
1.3E+03	2.95E-17
1.5E+03	3.02E-17
2.0E+03	3.37E-17
3.0E+03	3.67E-17
3.5E+03	3.90E-17
4.3E+03	4.12E-17
5.0E+03	4.22E-17
5.8E+03	4.19E-17
6.5E+03	4.13E-17
7.0E+03	4.11E-17
8.0E+03	4.05E-17
8.5E+03	3.96E-17
8.8E+03	3.92E-17

(3^1P) Energy eV/amu	Cross Section cm^2
2.5E+03	2.05E-18
3.0E+03	1.89E-18
3.5E+03	1.61E-18
4.3E+03	1.45E-18
5.0E+03	1.33E-18
5.8E+03	1.28E-18
6.5E+03	1.33E-18
7.0E+03	1.47E-18
8.0E+03	1.64E-18
8.5E+03	1.83E-18
9.0E+03	1.91E-18

References: 20, 163

Accuracy: 60%

Notes: (1) The cross section (σ_i) for capture into state i is determined by measuring the emission cross section (σ_{ij}) for a transition from state i to j, correcting for any branching from state i and correcting for cascading from upper states.
(2) In these measurements cascade was neglected in the analysis of the cross section.

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

(2^1P) E_{min} = 1.3E+03 eV/amu, E_{max} = 8.8E+03 eV/amu

(3^1P) E_{min} = 2.5E+03 eV/amu, E_{max} = 9.0E+03 eV/amu

Chebyshev Fitting Parameters for Cross Sections

	A0	A1	A2	A3	A4	A5	A6	A7	A8
(2^1P)	-75.7187	.173380	-.0602524	-.0334767	.00459756	.00195946	.00158847	-4.27148E-04	-.00726785
(3^1P)	-81.8842	-.0794869	.196956	.0554748	-.00702131	-.0164034	-.0116162	.00548035	1.56409E-04

The fit represents the (2^1P) cross sections with an rms deviation of 0.2%.
The maximum deviation is 0.2% at 6.4E+03 eV/amu.

The fit represents the (3^1P) cross sections with an rms deviation of 0.2%.
The maximum deviation is 0.4% at 7.9E+03 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + H_2 \rightarrow He(2^1P, 3^1P)$



Line Emission Cross Sections of Electron Capture Collisions for

$He^+ + H_2 \rightarrow He(587.6, 447.1, 667.8\ nm)$

(587.6 nm)		(667.8 nm)		(447.1 nm)	
Energy eV/amu	Cross Section cm^2	Energy eV/amu	Cross Section cm^2	Energy eV/amu	Cross Section cm^2
1.7E+01	1.12E-21	3.7E+01	9.97E-22	2.0E+01	9.98E-22
2.0E+01	3.58E-21	5.0E+01	7.92E-21	2.5E+01	2.67E-21
2.5E+01	7.00E-21	6.0E+01	1.47E-20	3.0E+01	3.75E-21
3.0E+01	8.74E-21	8.0E+01	4.70E-20	4.0E+01	5.02E-21
4.0E+01	1.26E-20	1.0E+02	1.04E-19	5.0E+01	6.21E-21
5.0E+01	1.81E-20	1.5E+02	2.70E-19	6.0E+01	7.65E-21
6.0E+01	2.83E-20	1.8E+02	3.35E-19	7.0E+01	1.00E-20
8.2E+01	7.80E-20	2.0E+02	3.95E-19	8.0E+01	1.77E-20
1.0E+02	1.72E-19			1.0E+02	6.11E-20
1.5E+02	6.40E-19			1.2E+02	9.53E-20
2.0E+02	1.12E-18			1.5E+02	1.30E-19
2.4E+02	1.17E-18			2.0E+02	1.60E-19
3.0E+02	1.16E-18			2.5E+02	1.75E-19
5.2E+02	1.02E-18			3.0E+02	1.82E-19
8.3E+02	8.33E-19			4.0E+02	1.81E-19
1.0E+03	7.69E-19			6.0E+02	1.63E-19
1.6E+03	7.00E-19			8.0E+02	1.45E-19
2.0E+03	6.93E-19			1.0E+03	1.31E-19
3.0E+03	7.25E-19			1.5E+03	1.07E-19
4.0E+03	7.68E-19			2.0E+03	9.32E-20
5.2E+03	8.05E-19			3.0E+03	8.58E-20
7.8E+03	8.90E-19			4.0E+03	8.98E-20
				5.0E+03	9.72E-20
				6.0E+03	1.06E-19
				7.2E+03	1.16E-19
				8.2E+03	1.27E-19
				1.0E+04	1.48E-19

References: 199, 200

Accuracy: 40%

Notes: (1) Relative data of Polyakova et al (ref. 200) have been normalized to data of Isler et al (ref. 199) at 125 eV/amu.
(2) The 587.6 nm transition is the strongest He emission from He^+ capture collisions in H_2.
(3) The $4^3D \rightarrow 2^3P$ transition is 447.1 nm, the $3^3D \rightarrow 2^3P$ transition is 587.6 nm, and the $3^1D \rightarrow 2^1P$ transition is 667.8 nm.

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

(587.6 nm) E_{min} = 1.7E+01 eV/amu, E_{max} = 7.8E+03 eV/amu
(667.8 nm) E_{min} = 3.7E+01 eV/amu, E_{max} = 2.0E+02 eV/amu
(447.1 nm) E_{min} = 2.0E+01 eV/amu, E_{max} = 1.0E+04 eV/amu

Chebyshev Fitting Parameters for Cross Sections

	A0	A1	A2	A3	A4	A5	A6	A7	A8
(587.6 nm)	-86.6646	2.83888	-1.58217	.357959	.371758	-.133932	-.263367	.286400	-.104165
(667.8 nm)	-89.3928	2.77428	-.477421	.0466864	-.123082	.139543	-.0694321	.0304386	
(447.1 nm)	-89.4303	1.99037	-1.26304	.473572	.362531	-.183668	-.0812802	.198425	-.197268

The fit represents the 587.6 nm cross section with an rms deviation of 4.7%.
The maximum deviation is 10.8% at 3.1E+02 eV/amu.
The fit represents the 667.8 nm cross section with an rms deviation of 0.0%.
The maximum deviation is 0.0%.
The fit represents the 447.1 nm cross section with an rms deviation of 14.7%.
The maximum deviation is 42.6% at 7.1E+01 eV/amu.
See appendix for Chebyshev fit details.

$He^+ + H_2 \rightarrow He(587.6, 667.8, 447.1\ nm)$



Electron Capture Cross Sections for
$He^+ + He \rightarrow He(2^1S + 2^3S)$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.0E+01	4.39E+06	1.00E-19
1.5E+01	5.38E+06	1.94E-19
3.0E+01	7.61E+06	7.49E-19
5.0E+01	9.82E+06	1.69E-18
8.0E+01	1.24E+07	2.33E-18
1.5E+02	1.70E+07	2.81E-18
3.0E+02	2.41E+07	2.99E-18
6.0E+02	3.40E+07	3.35E-18
1.5E+03	5.38E+07	4.54E-18
2.5E+03	6.95E+07	6.54E-18
4.0E+03	8.79E+07	9.49E-18
6.0E+03	1.08E+08	1.46E-17
9.0E+03	1.32E+08	2.40E-17
1.0E+04	1.39E+08	2.61E-17
1.5E+04	1.70E+08	2.41E-17
3.3E+04	2.50E+08	1.85E-17
5.0E+04	3.11E+08	1.52E-17

References: 98, 196, 197, 201, 202

Accuracy: 40%

Note: (1) Theoretical values of Sural et al (ref. 202) are used to extrapolate the data of Gilbody et al (ref. 98) at 5×10^3 eV/amu to those of MacVicar and Barst (ref. 201) at 55 eV/amu.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 1.0E+01 eV/amu, E_{max} = 5.0E+04 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-80.6529	2.39212	-.649912	.218666	-.468812	-.126058	.149096	.0275996	.0661909

The fit represents the above cross sections with an rms deviation of 5.2%.
The maximum deviation is 12.8% at 1.5E+04 eV/amu.
See appendix for Chebyshev fit details.

$He^+ + He \rightarrow He(2^1S + 2^3S)$

Cross Section vs. Energy



Electron Capture Rate Coefficients for
$He^+ + He \rightarrow He(2^1S + 2^3S)$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

He^+ Temp. (eV)	Equal Temp.	He Temp. (eV) 10.	100.	500.	1000.	5000.	10000.	20000.
2.0E+00	9.32E-17	7.50E-14	1.05E-11	4.83E-11	7.97E-11	3.80E-10	9.44E-10	1.97E-09
4.0E+00	1.30E-14	1.29E-13	1.08E-11	4.84E-11	7.98E-11	3.80E-10	9.44E-10	1.97E-09
7.0E+00	1.29E-13	2.39E-13	1.12E-11	4.86E-11	8.00E-11	3.81E-10	9.45E-10	1.97E-09
1.0E+01	3.83E-13	3.83E-13	1.16E-11	4.88E-11	8.02E-11	3.81E-10	9.45E-10	1.97E-09
2.0E+01	2.09E-12	1.09E-12	1.30E-11	4.95E-11	8.08E-11	3.82E-10	9.46E-10	1.97E-09
4.0E+01	7.41E-12	3.29E-12	1.56E-11	5.09E-11	8.19E-11	3.84E-10	9.48E-10	1.97E-09
7.0E+01	1.56E-11	7.41E-12	1.93E-11	5.29E-11	8.37E-11	3.87E-10	9.52E-10	1.97E-09
1.0E+02	2.27E-11	1.16E-11	2.27E-11	5.49E-11	8.55E-11	3.90E-10	9.56E-10	1.98E-09
2.0E+02	4.08E-11	2.38E-11	3.25E-11	6.13E-11	9.14E-11	3.99E-10	9.67E-10	1.98E-09
4.0E+02	6.75E-11	4.15E-11	4.81E-11	7.36E-11	1.03E-10	4.19E-10	9.91E-10	2.00E-09
7.0E+02	1.03E-10	6.20E-11	6.75E-11	9.14E-11	1.21E-10	4.50E-10	1.03E-09	2.03E-09
1.0E+03	1.40E-10	8.02E-11	8.55E-11	1.09E-10	1.40E-10	4.82E-10	1.06E-09	2.05E-09
2.0E+03	2.88E-10	1.41E-10	1.46E-10	1.73E-10	2.09E-10	5.92E-10	1.18E-09	2.13E-09
4.0E+03	7.07E-10	2.89E-10	2.97E-10	3.33E-10	3.80E-10	8.25E-10	1.40E-09	2.27E-09
7.0E+03	1.40E-09	5.93E-10	6.03E-10	6.49E-10	7.07E-10	1.18E-09	1.70E-09	2.46E-09
1.0E+04	1.97E-09	9.45E-10	9.56E-10	1.00E-09	1.06E-09	1.50E-09	1.97E-09	2.62E-09
2.0E+04	2.98E-09	1.97E-09	1.98E-09	2.01E-09	2.05E-09	2.33E-09	2.62E-09	2.98E-09

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 2.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
10.	-5.00830E+01	5.12336E+00	-2.62808E-01	8.40742E-02	2.48250E-01	-1.23169E-01	-7.59116E-02
100.	-4.68137E+01	2.59124E+00	7.95345E-01	5.35033E-02	1.89675E-03	-1.71822E-02	-5.25035E-02
500.	-4.53295E+01	1.72678E+00	8.43475E-01	1.98073E-01	-3.44676E-02	-5.47555E-02	-3.44800E-02
1000.	-4.46993E+01	1.46770E+00	7.78110E-01	2.19162E-01	-2.15385E-02	-5.87801E-02	-3.66707E-02
5000.	-4.24353E+01	7.86347E-01	4.49607E-01	1.53041E-01	5.53429E-03	-2.94156E-02	-2.10243E-02
10000.	-4.10534E+01	4.26270E-01	2.51921E-01	9.52564E-02	1.32834E-02	-1.03321E-02	-9.66498E-03
20000.	-3.98951E+01	1.67503E-01	1.03069E-01	4.32423E-02	9.86815E-03	-1.82969E-03	-3.51334E-03
Equal Temp.	-5.11417E+01	7.32023E+00	-2.04605E+00	1.24837E+00	-5.12909E-01	3.38169E-02	-1.21200E-01

See appendix for Chebyshev fit details.

$He^+ + He \rightarrow He(2^1S + 2^3S)$

Maxwellian – Maxwellian



Electron Capture Rate Coefficients for
$He + He^+ \rightarrow He(2^1S + 2^3S)$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^+ Temp. (eV)	He Energy (eV/amu) 10000.	15000.	20000.	25000.	30000.	35000.	40000.
1.0E+00	3.62E-09**	4.10E-09**	4.29E-09**	4.44E-09**	4.57E-09**	4.65E-09**	4.68E-09**
2.0E+00	3.61E-09**	4.10E-09**	4.29E-09**	4.44E-09**	4.57E-09**	4.65E-09**	4.68E-09**
4.0E+00	3.61E-09**	4.10E-09**	4.29E-09**	4.44E-09**	4.57E-09**	4.65E-09**	4.68E-09**
7.0E+00	3.60E-09**	4.10E-09**	4.29E-09**	4.44E-09**	4.57E-09**	4.65E-09**	4.68E-09**
1.0E+01	3.59E-09**	4.10E-09**	4.29E-09**	4.44E-09**	4.57E-09**	4.65E-09**	4.68E-09**
2.0E+01	3.58E-09**	4.09E-09**	4.29E-09**	4.44E-09**	4.57E-09**	4.65E-09**	4.68E-09**
4.0E+01	3.57E-09**	4.09E-09**	4.29E-09**	4.44E-09**	4.57E-09**	4.65E-09**	4.68E-09**
7.0E+01	3.55E-09**	4.09E-09**	4.29E-09**	4.45E-09**	4.57E-09**	4.65E-09**	4.68E-09**
1.0E+02	3.53E-09**	4.09E-09**	4.29E-09**	4.45E-09**	4.57E-09**	4.65E-09**	4.68E-09**
2.0E+02	3.50E-09**	4.08E-09**	4.29E-09**	4.45E-09**	4.57E-09**	4.65E-09**	4.68E-09**
4.0E+02	3.45E-09**	4.08E-09**	4.29E-09**	4.45E-09**	4.57E-09**	4.64E-09**	4.67E-09**
7.0E+02	3.40E-09**	4.07E-09**	4.29E-09**	4.45E-09**	4.57E-09**	4.64E-09**	4.64E-09**
1.0E+03	3.37E-09**	4.06E-09**	4.29E-09**	4.45E-09**	4.57E-09**	4.64E-09**	4.58E-09**
2.0E+03	3.31E-09**	4.03E-09**	4.29E-09**	4.45E-09**	4.56E-09**	4.57E-09**	4.30E-09**
4.0E+03	3.28E-09*	3.98E-09**	4.28E-09**	4.43E-09**	4.47E-09**	4.32E-09**	3.86E-09**
7.0E+03	3.30E-09*	3.93E-09**	4.23E-09**	4.34E-09**	4.25E-09**	3.96E-09**	3.45E-09**
1.0E+04	3.34E-09*	3.89E-09**	4.14E-09**	4.18E-09**	4.02E-09**	3.67E-09**	3.18E-09**
2.0E+04	3.43E-09*	3.71E-09*	3.77E-09**	3.65E-09**	3.40E-09**	3.07E-09**	2.68E-09**

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-3.89516E+01	-4.51244E-02	1.69953E-03	1.65062E-02	1.06870E-02	2.64417E-03	-1.07811E-03
15000.	-3.86669E+01	-3.54627E-02	-2.26753E-02	-1.21671E-02	-5.34610E-03	-1.80140E-03	-7.78336E-04
20000.	-3.85671E+01	-3.27909E-02	-2.82327E-02	-2.24375E-02	-1.53665E-02	-8.81047E-03	-4.39280E-03
25000.	-3.85158E+01	-5.09890E-02	-4.36393E-02	-3.40163E-02	-2.27913E-02	-1.25967E-02	-5.50588E-03
30000.	-3.84991E+01	-8.72591E-02	-7.04596E-02	-4.87052E-02	-2.79930E-02	-1.24934E-02	-2.97299E-03
35000.	-3.85280E+01	-1.39891E-01	-1.06634E-01	-6.39054E-02	-2.79358E-02	-6.05892E-03	3.51648E-03
40000.	-3.86081E+01	-2.17036E-01	-1.48805E-01	-7.02850E-02	-1.44258E-02	8.74155E-03	8.88909E-03

See appendix for Chebyshev fit details.

$He + He^{+} \rightarrow He(2^{1}S + 2^{3}S)$

Beam – Maxwellian



Electron Capture Cross Sections for
$He^+ + He \rightarrow He(2^1P) + He^+$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.3E+03	4.91E+07	1.25E-18
2.0E+03	6.21E+07	1.18E-18
3.0E+03	7.61E+07	1.07E-18
5.0E+03	9.82E+07	9.00E-19
8.0E+03	1.24E+08	7.67E-19
1.0E+04	1.39E+08	7.03E-19
2.0E+04	1.96E+08	5.17E-19
4.0E+04	2.78E+08	3.52E-19
7.0E+04	3.68E+08	2.29E-19
1.0E+05	4.39E+08	1.46E-19
1.5E+05	5.38E+08	4.86E-20
2.0E+05	6.21E+08	1.60E-20
2.5E+05	6.94E+08	6.01E-21

References: 163, 203

Accuracy: Unknown

Notes: (1) Data between 1.2 - 9×10^3 eV/amu were never published in a referred journal.
(2) Cross sections between 9×10^3 and 1×10^5 eV/amu were obtained by interpolation between the low and high energy data.
(3) For energies less than 1×10^5 eV/amu the cross sections measured by Hippler et al (ref. 203) diverge from the data presented here.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 1.3E+03 eV/amu, E_{max} = 2.5E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-85.5425	-2.19593	-.936988	-.450041	-.224752	-.0551700	.0123849	.0330118	.0293219

The fit represents the above cross sections with an rms deviation of 1.0%.
The maximum deviation is 2.0% at 7.0E+04 eV/amu.
See appendix for Chebyshev fit details.

$He^+ + He \rightarrow He(2^1P) + He^+$

Cross Section vs. Energy



Electron Capture Rate Coefficients for
$He^+ + He \rightarrow He(2^1P) + He^+$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

He^+ Temp. (eV)	Equal Temp.	He Temp. (eV) 500.	1000.	2000.	4000.	7000.	10000.	20000.
4.0E+02	3.80E-13	7.28E-13	4.54E-12	1.72E-11	3.90E-11	5.75E-11	6.77E-11	8.29E-11
7.0E+02	4.54E-12	2.64E-12	8.03E-12	2.11E-11	4.15E-11	5.87E-11	6.85E-11	8.32E-11
1.0E+03	1.19E-11	5.64E-12	1.19E-11	2.48E-11	4.38E-11	5.99E-11	6.92E-11	8.34E-11
2.0E+03	3.54E-11	1.85E-11	2.48E-11	3.54E-11	5.03E-11	6.36E-11	7.15E-11	8.42E-11
4.0E+03	5.99E-11	3.98E-11	4.38E-11	5.03E-11	5.99E-11	6.92E-11	7.52E-11	8.56E-11
7.0E+03	7.52E-11	5.79E-11	5.99E-11	6.36E-11	6.92E-11	7.52E-11	7.95E-11	8.73E-11
1.0E+04	8.26E-11	6.80E-11	6.92E-11	7.15E-11	7.52E-11	7.95E-11	8.26E-11	8.87E-11
2.0E+04	9.16E-11	8.30E-11	8.34E-11	8.42E-11	8.56E-11	8.73E-11	8.87E-11	9.16E-11

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 4.0E+02 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
500.	-4.99295E+01	2.33819E+00	-6.50090E-01	4.21924E-02	3.49426E-02	-1.24395E-02	-1.11193E-04
1000.	-4.88301E+01	1.49821E+00	-2.72937E-01	-4.12913E-02	2.79783E-02	-2.24988E-03	-2.38793E-03
2000.	-4.78893E+01	8.38347E-01	-5.35952E-02	-4.83802E-02	7.58342E-03	3.24901E-03	-8.41972E-04
4000.	-4.71958E+01	4.15561E-01	2.60934E-02	-2.42784E-02	-3.22992E-03	1.67069E-03	3.02996E-04
7000.	-4.68029E+01	2.17494E-01	3.41131E-02	-8.30328E-03	-3.82190E-03	-5.16068E-05	2.23242E-04
10000.	-4.66140E+01	1.38147E-01	2.86268E-02	-2.78051E-03	-2.76127E-03	-4.58542E-04	5.63634E-05
20000.	-4.63524E+01	4.94719E-02	1.41098E-02	5.11657E-04	-9.99744E-04	-2.34546E-04	-2.76502E-04
Equal Temp.	-4.94385E+01	2.39001E+00	-1.04807E+00	3.37080E-01	-8.59497E-02	1.50447E-02	-2.44681E-03

See appendix for Chebyshev fit details.

$He^+ + He \rightarrow He(2^1P) + He^+$

Maxwellian – Maxwellian



Electron Capture Rate Coefficients for
$He + He^+ \rightarrow He(2^1P) + He^+$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^+ Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	60000.	80000.	100000.	120000.
1.0E+00	9.76E-11	1.02E-10*	9.78E-11*	8.77E-11*	7.60E-11*	6.40E-11*	4.28E-11**
2.0E+00	9.76E-11	1.02E-10	9.78E-11*	8.77E-11*	7.60E-11*	6.40E-11*	4.28E-11**
4.0E+00	9.76E-11	1.02E-10	9.77E-11*	8.77E-11*	7.60E-11*	6.40E-11*	4.28E-11**
7.0E+00	9.76E-11	1.02E-10	9.77E-11*	8.77E-11*	7.60E-11*	6.39E-11*	4.28E-11**
1.0E+01	9.76E-11	1.01E-10	9.77E-11*	8.77E-11*	7.60E-11*	6.39E-11*	4.28E-11**
2.0E+01	9.76E-11	1.01E-10	9.77E-11*	8.77E-11*	7.60E-11*	6.38E-11*	4.28E-11**
4.0E+01	9.76E-11	1.01E-10	9.76E-11*	8.77E-11*	7.60E-11*	6.36E-11*	4.29E-11*
7.0E+01	9.75E-11	1.01E-10	9.75E-11	8.77E-11*	7.60E-11*	6.35E-11*	4.29E-11*
1.0E+02	9.75E-11	1.01E-10	9.75E-11	8.77E-11*	7.60E-11*	6.33E-11*	4.29E-11*
2.0E+02	9.75E-11	1.01E-10	9.74E-11	8.77E-11	7.60E-11*	6.30E-11*	4.29E-11*
4.0E+02	9.74E-11	1.01E-10	9.72E-11	8.77E-11	7.60E-11*	6.26E-11*	4.30E-11*
7.0E+02	9.73E-11	1.01E-10	9.70E-11	8.77E-11	7.60E-11*	6.21E-11*	4.31E-11*
1.0E+03	9.72E-11	1.01E-10	9.68E-11	8.76E-11	7.60E-11*	6.18E-11*	4.32E-11*
2.0E+03	9.70E-11	1.01E-10	9.64E-11	8.74E-11	7.58E-11*	6.10E-11*	4.34E-11*
4.0E+03	9.68E-11	1.00E-10	9.58E-11	8.68E-11	7.50E-11	5.99E-11*	4.36E-11*
7.0E+03	9.63E-11	9.96E-11	9.51E-11	8.59E-11	7.38E-11	5.88E-11*	4.36E-11*
1.0E+04	9.59E-11	9.90E-11	9.43E-11	8.50E-11	7.26E-11	5.80E-11*	4.35E-11*
2.0E+04	9.52E-11	9.71E-11	9.18E-11	8.18E-11	6.92E-11	5.56E-11*	4.27E-11*

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-4.61121E+01	-1.01540E-02	-5.73295E-03	-2.50999E-03	-8.40721E-04	-1.26867E-04	1.76279E-04
20000.	-4.60412E+01	-1.63968E-02	-9.47895E-03	-4.78055E-03	-2.35744E-03	-1.14505E-03	-5.31594E-04
40000.	-4.61253E+01	-2.31383E-02	-1.33331E-02	-6.62458E-03	-3.18902E-03	-1.53689E-03	-7.59534E-04
60000.	-4.63375E+01	-2.16963E-02	-1.67017E-02	-1.07850E-02	-5.66807E-03	-2.33604E-03	-7.64767E-04
80000.	-4.66323E+01	-2.96500E-02	-2.31326E-02	-1.47960E-02	-7.40205E-03	-2.57953E-03	-2.97132E-04
100000.	-4.70209E+01	-5.97314E-02	-2.79167E-02	-9.52672E-03	-2.96002E-03	-1.17463E-03	-6.06512E-04
120000.	-4.77385E+01	5.70178E-03	-3.34664E-04	-4.37679E-03	-4.62136E-03	-2.70887E-03	-7.41640E-04

See appendix for Chebyshev fit details.

$He + He^+ \rightarrow He(2^1P) + He^+$



Electron Capture Cross Sections for

$He^+ + He \rightarrow He(3^1S, 3^1P, 3^1D) + He^+$

(3^1S) Energy eV/amu	(3^1S) Cross Section cm^2	(3^1P) Energy eV/amu	(3^1P) Cross Section cm^2	(3^1D) Energy eV/amu	(3^1D) Cross Section cm^2
2.5E+03	2.17E-19	7.5E+02	2.34E-19	2.0E+03	5.30E-19
3.0E+03	2.02E-19	1.0E+03	2.53E-19	2.5E+03	4.91E-19
3.8E+03	1.94E-19	1.5E+03	3.57E-19	3.0E+03	5.42E-19
5.0E+03	2.13E-19	2.0E+03	6.16E-19	3.5E+03	6.64E-19
7.0E+03	2.79E-19	2.5E+03	7.91E-19	4.0E+03	7.76E-19
8.0E+03	3.30E-19	2.6E+03	7.96E-19	5.0E+03	8.93E-19
1.0E+04	4.93E-19	3.0E+03	7.34E-19	6.0E+03	9.23E-19
1.5E+04	8.44E-19	3.5E+03	6.28E-19	7.0E+03	9.13E-19
2.0E+04	1.13E-18	5.0E+03	5.00E-19	1.0E+04	8.17E-19
2.5E+04	1.27E-18	6.5E+03	4.72E-19	1.5E+04	6.28E-19
3.5E+04	1.38E-18	8.0E+03	5.03E-19	2.0E+04	4.69E-19
4.0E+04	1.40E-18	1.0E+04	5.97E-19	2.5E+04	3.48E-19
		1.3E+04	8.63E-19	3.0E+04	2.50E-19
		1.5E+04	1.12E-18	3.7E+04	1.41E-19
		2.0E+04	1.44E-18		
		2.2E+04	1.45E-18		
		2.5E+04	1.40E-18		
		3.0E+04	1.24E-18		
		3.5E+04	1.02E-18		
		3.8E+04	8.69E-19		

References: 3^1S - 204; 3^1P - 20, 204, 205, 206; 3^1D - 20, 204, 206

Accuracy: 40%

Notes: (1) The quoted accuracy is only an estimate. The disagreement between the data sets is as large as a factor of 2 or more.
(2) The cross section (σ_i) for capture into state i is determined by measuring the emission cross section (σ_{ij}) for a transition from state i to j, correcting for any branching from state i and correcting for cascading from upper states.

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

(3^1S) E_{min} = 2.5E+03 eV/amu, E_{max} = 4.0E+04 eV/amu
(3^1P) E_{min} = 7.5E+02 eV/amu, E_{max} = 3.8E+04 eV/amu
(3^1D) E_{min} = 2.0E+03 eV/amu, E_{max} = 3.7E+04 eV/amu

Chebyshev Fitting Parameters for Cross Sections

	A0	A1	A2	A3	A4	A5	A6	A7	A8
(3^1S)	-84.2567	1.14450	.0774864	-.248351	.00546389	.0475617	-6.68831E-04	-.0125101	.00445413
(3^1P)	-83.9443	.784951	-.124752	.0431388	-.173909	-.262303	.0929503	.0881888	-.0630714
(3^1D)	-84.3847	-.485772	-.595547	-.0904580	.0360540	-.0805642	.0223178	-.00673603	-.0137040

The fit represents the (3^1S) cross section with an rms deviation of 1.2%.
The maximum deviation is 1.9% at 8.0E+03 eV/amu.

The fit represents the (3^1P) cross section with an rms deviation of 3.7%.
The maximum deviation is 9.7% at 3.5E+03 eV/amu.

The fit represents the (3^1D) cross section with an rms deviation of 1.0%.
The maximum deviation is 2.5% at 3.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + He \rightarrow He(3^1S, 3^1P, 3^1D) + He^+$

Cross Section vs. Energy



Electron Capture Cross Sections for

$He^+ + He \rightarrow He(3^3S, 3^3P, 3^3D) + He^+$

(3^3S) Energy eV/amu	(3^3S) Cross Section cm^2	(3^3P) Energy eV/amu	(3^3P) Cross Section cm^2	(3^3D) Energy eV/amu	(3^3D) Cross Section cm^2
5.0E+03	4.52E-19	2.5E+02	2.05E-19	2.5E+02	1.34E-18
6.0E+03	4.65E-19	3.0E+02	2.02E-19	3.5E+02	1.21E-18
7.0E+03	4.89E-19	4.0E+02	2.01E-19	5.0E+02	1.07E-18
9.0E+03	5.87E-19	6.0E+02	2.33E-19	7.0E+02	8.97E-19
1.0E+04	6.64E-19	8.0E+02	3.03E-19	8.0E+02	8.81E-19
1.5E+04	1.26E-18	1.0E+03	3.79E-19	9.0E+02	8.94E-19
2.0E+04	1.93E-18	1.5E+03	6.16E-19	1.0E+03	9.43E-19
2.5E+04	2.68E-18	2.0E+03	8.05E-19	1.4E+03	1.35E-18
3.0E+04	3.06E-18	2.5E+03	6.58E-19	1.7E+03	1.08E-18
		3.0E+03	6.31E-19	2.0E+03	7.98E-19
		4.0E+03	6.60E-19	2.5E+03	6.61E-19
		6.0E+03	7.15E-19	2.8E+03	7.27E-19
		8.0E+03	7.71E-19	3.4E+03	8.22E-19
		9.9E+03	8.35E-19	3.8E+03	8.50E-19
		1.5E+04	9.99E-19	4.5E+03	8.12E-19
		2.0E+04	1.11E-18	5.4E+03	6.94E-19
		3.0E+04	1.25E-18	5.9E+03	6.69E-19
		3.7E+04	1.31E-18	6.5E+03	7.03E-19
				7.1E+03	8.22E-19
				7.7E+03	9.52E-19
				8.5E+03	9.90E-19
				1.0E+04	9.53E-19
				1.2E+04	8.48E-19
				1.5E+04	6.73E-19
				1.7E+04	5.33E-19
				2.2E+04	3.49E-19
				2.9E+04	2.34E-19
				3.8E+04	1.63E-19

References: (3^3S) - 20, 206; (3^3P) - 20, 204, 205, 206; (3^3D) - 20, 204, 205, 206

Accuracy: 40%

Notes: (1) The quoted accuracy is only an estimate. The disagreement between the data sets is as large as a factor of 2 or more.
(2) The cross section (σ_i) for capture into state i is determined by measuring the emission cross section (σ_{ij}) for a transition from state i to j, correcting for any branching from state i and correcting for cascading from upper states.

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

(3^3S) E_{min} = 5.0E+03 eV/amu, E_{max} = 3.0E+04 eV/amu
(3^3P) E_{min} = 2.5E+02 eV/amu, E_{max} = 3.7E+04 eV/amu
(3^3D) E_{min} = 2.5E+02 eV/amu, E_{max} = 3.8E+04 eV/amu

Chebyshev Fitting Parameters for Cross Sections

	A0	A1	A2	A3	A4	A5	A6	A7	A8
(3^3S)	-82.7952	1.05996	.146880	-.0978993	-.0162972	.00664821	-.0131980	-.0124651	-.00414755
(3^3P)	-84.1155	.979750	-.132417	.00802795	.121042	-.129354	-.0123880	.0827080	-.0303641
(3^3D)	-83.6677	-.841000	-.315720	-.312650	-.0741395	-.00902084	.0814994	.122383	-.0655599

The fit represents the (3^3S) cross section with an rms deviation of 0.0%.
The maximum deviation is 0.0% at 6.0E+03 eV/amu.
The fit represents the (3^3P) cross section with an rms deviation of 5.3%.
The maximum deviation is 9.0% at 3.0E+03 eV/amu.
The fit represents the (3^3D) cross section with an rms deviation of 11.0%.
The maximum deviation is 23.9% at 2.5E+03 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + He \rightarrow He(3^3S, 3^3P, 3^3D) + He^+$



Electron Capture Cross Sections for
$He^+ + He \rightarrow He(4^1S, 4^1P, 4^1D) + He^+$

(4^1S) Energy eV/amu	(4^1S) Cross Section cm^2	(4^1P) Energy eV/amu	(4^1P) Cross Section cm^2	(4^1D) Energy eV/amu	(4^1D) Cross Section cm^2
6.2E+02	8.66E-21	1.0E+03	8.65E-20	2.5E+02	7.24E-20
7.3E+02	1.11E-20	1.1E+03	6.69E-20	3.0E+02	9.38E-20
9.4E+02	1.29E-20	1.2E+03	5.33E-20	3.5E+02	1.03E-19
1.0E+03	1.30E-20	1.3E+03	4.82E-20	4.1E+02	8.76E-20
1.2E+03	1.23E-20	1.3E+03	4.83E-20	5.0E+02	5.10E-20
1.3E+03	1.16E-20	1.4E+03	5.17E-20	6.2E+02	3.55E-20
1.4E+03	1.18E-20	1.6E+03	6.04E-20	7.2E+02	3.99E-20
1.5E+03	1.33E-20	1.8E+03	7.45E-20	8.1E+02	4.86E-20
1.6E+03	1.65E-20	2.2E+03	9.50E-20	9.1E+02	6.14E-20
1.8E+03	2.32E-20	2.6E+03	1.16E-19	1.0E+03	7.60E-20
2.0E+03	2.87E-20	3.2E+03	1.41E-19	1.1E+03	7.86E-20
2.3E+03	3.62E-20	3.6E+03	1.61E-19	1.3E+03	7.94E-20
2.6E+03	3.87E-20	3.9E+03	1.63E-19	1.6E+03	8.07E-20
2.7E+03	3.93E-20	4.3E+03	1.61E-19	1.9E+03	8.23E-20
3.0E+03	3.83E-20	4.9E+03	1.52E-19	2.1E+03	8.65E-20
3.2E+03	3.72E-20	5.4E+03	1.52E-19	2.2E+03	1.04E-19
3.7E+03	3.62E-20	6.4E+03	1.64E-19	2.4E+03	1.16E-19
4.1E+03	3.69E-20	7.5E+03	1.86E-19	2.5E+03	1.18E-19
4.9E+03	4.02E-20	8.6E+03	2.32E-19	2.7E+03	1.10E-19
6.5E+03	5.18E-20	1.0E+04	2.90E-19	2.9E+03	8.95E-20
7.9E+03	6.19E-20	1.1E+04	3.50E-19	3.0E+03	7.13E-20
9.6E+03	7.73E-20	1.3E+04	4.29E-19	3.1E+03	6.36E-20
1.2E+04	1.00E-19	1.6E+04	5.01E-19	3.3E+03	6.24E-20
1.4E+04	1.39E-19	1.8E+04	5.80E-19	3.5E+03	6.40E-20
1.6E+04	1.74E-19	2.0E+04	6.08E-19	3.7E+03	7.01E-20
1.9E+04	2.02E-19	2.4E+04	5.83E-19	3.9E+03	8.14E-20
2.2E+04	2.20E-19	2.8E+04	5.20E-19	4.1E+03	9.94E-20
2.8E+04	2.29E-19	3.2E+04	4.48E-19	4.4E+03	1.18E-19
3.3E+04	2.25E-19	3.7E+04	3.78E-19	4.8E+03	1.42E-19
3.8E+04	2.14E-19	3.9E+04	3.53E-19	5.6E+03	1.71E-19
				7.0E+03	2.11E-19
				8.3E+03	2.42E-19
				9.7E+03	2.75E-19
				1.2E+04	3.12E-19
				2.0E+04	2.73E-19
				3.0E+04	8.42E-20
				3.8E+04	4.33E-20

References: (4^1S) - 20, 204, 205, 206, 207; (4^1P) - 20, 166, 204, 205; (4^1D) - 20, 166, 204, 205, 206

Accuracy: 40%

Notes: (1) The quoted accuracy is only an estimate. The disagreement between the data sets is as large as a factor of 2 or more.
(2) The cross section (σ_i) for capture into state i is determined by measuring the emission cross section (σ_{ij}) for a transition from state i to j, correcting for any branching from state i and correcting for cascading from upper states.

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

(4^1S) E_{min} = 6.2E+02 eV/amu, E_{max} = 3.8E+04 eV/amu
(4^1P) E_{min} = 1.0E+03 eV/amu, E_{max} = 3.9E+04 eV/amu
(4^1D) E_{min} = 2.5E+02 eV/amu, E_{max} = 3.8E+04 eV/amu

Chebyshev Fitting Parameters for Cross Sections

	A0	A1	A2	A3	A4	A5	A6	A7	A8
(4^1S)	-88.9247	1.69591	.0188804	-.0955575	-.0823400	-.127347	-.00167808	.149767	-.0939805
(4^1P)	-86.2453	1.12538	-.147764	-.263207	-.0637178	-.168724	.168441	.00791441	-.0240953
(4^1D)	-87.4670	.304065	-.224271	-.683543	-.287424	-.118320	-.0304248	.229919	-.141272

The fit represents the (4^1S) cross section with an rms deviation of 8.5%.
The maximum deviation is 17.4% at 7.3E+02 eV/amu.
The fit represents the (4^1P) cross section with an rms deviation of 3.5%.
The maximum deviation is 7.1% at 2.6E+03 eV/amu.
The fit represents the (4^1D) cross section with an rms deviation of 16.5%.
The maximum deviation is 37.8% at 3.3E+03 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + He \rightarrow He(4^1S, 4^1P, 4^1D) + He^+$



Electron Capture Cross Sections for
$He^+ + He \rightarrow He(4^3S, 4^3P, 4^3D) + He^+$

(4^3S) Energy eV/amu	(4^3S) Cross Section cm^2	(4^3P) Energy eV/amu	(4^3P) Cross Section cm^2	(4^3D) Energy eV/amu	(4^3D) Cross Section cm^2
3.7E+02	3.99E-20	2.5E+02	3.05E-20	2.5E+02	2.84E-19
4.0E+02	3.25E-20	3.0E+02	2.38E-20	3.0E+02	3.39E-19
5.0E+02	2.09E-20	4.0E+02	1.78E-20	3.5E+02	3.62E-19
6.0E+02	2.12E-20	5.0E+02	1.62E-20	3.8E+02	3.56E-19
7.0E+02	3.06E-20	5.3E+02	1.64E-20	4.2E+02	3.26E-19
7.7E+02	3.58E-20	5.6E+02	1.95E-20	4.9E+02	2.51E-19
8.6E+02	3.44E-20	5.7E+02	3.00E-20	5.7E+02	2.06E-19
9.7E+02	3.17E-20	6.3E+02	4.53E-20	6.8E+02	1.76E-19
1.1E+03	3.08E-20	6.7E+02	3.97E-20	7.7E+02	1.66E-19
1.3E+03	3.30E-20	7.5E+02	3.37E-20	8.1E+02	1.71E-19
1.5E+03	4.12E-20	8.3E+02	2.95E-20	8.9E+02	1.97E-19
1.7E+03	4.94E-20	9.2E+02	2.72E-20	9.9E+02	2.33E-19
2.0E+03	6.01E-20	1.0E+03	2.64E-20	1.1E+03	2.68E-19
2.4E+03	6.91E-20	1.1E+03	2.85E-20	1.2E+03	2.89E-19
2.6E+03	7.00E-20	1.4E+03	3.84E-20	1.3E+03	2.89E-19
3.0E+03	6.90E-20	1.7E+03	4.83E-20	1.5E+03	2.64E-19
3.6E+03	6.54E-20	2.0E+03	6.20E-20	1.7E+03	2.26E-19
4.1E+03	6.71E-20	2.3E+03	8.50E-20	2.0E+03	1.85E-19
5.5E+03	7.97E-20	2.6E+03	9.79E-20	2.4E+03	1.50E-19
7.6E+03	1.09E-19	3.0E+03	1.02E-19	2.6E+03	1.35E-19
9.7E+03	1.48E-19	3.5E+03	9.51E-20	2.9E+03	1.21E-19
1.2E+04	1.96E-19	4.0E+03	8.49E-20	3.2E+03	1.20E-19
1.5E+04	2.73E-19	4.7E+03	7.84E-20	3.4E+03	1.23E-19
1.7E+04	3.88E-19	4.8E+03	7.76E-20	3.8E+03	1.37E-19
2.1E+04	5.52E-19	5.2E+03	8.09E-20	4.4E+03	1.60E-19
2.6E+04	7.13E-19	6.3E+03	9.53E-20	5.1E+03	1.91E-19
3.0E+04	8.28E-19	8.8E+03	1.28E-19	6.1E+03	2.30E-19
3.4E+04	8.61E-19	1.4E+04	1.88E-19	7.3E+03	2.77E-19
3.7E+04	8.65E-19	2.0E+04	2.47E-19	8.4E+03	3.08E-19
		2.7E+04	3.26E-19	9.8E+03	3.28E-19
		3.3E+04	3.87E-19	1.1E+04	3.24E-19
		3.8E+04	4.07E-19	1.3E+04	2.96E-19
				1.4E+04	2.63E-19
				1.6E+04	2.29E-19
				2.0E+04	1.78E-19
				2.5E+04	1.31E-19
				3.0E+04	1.07E-19
				3.4E+04	8.18E-20
				3.8E+04	6.62E-20

<u>References:</u> (4^3S) - 20, 166, 204, 205, 206, 207; (4^3P) - 20, 166, 204, 205, 207; (4^3D) - 20, 166, 204, 205, 206, 207

<u>Accuracy:</u> 40%

<u>Notes:</u> (1) The quoted accuracy is only an estimate. The disagreement between the data sets is as large as a factor of 2 or more.
(2) The cross section (σ_i) for capture into state i is determined by measuring the emission cross section (σ_{ij}) for a transition from state i to j, correcting for any branching from state i and correcting for cascading from upper states.

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

(4^3S) E_{min} = 3.7E+02 eV/amu, E_{max} = 3.7E+04 eV/amu
(4^3P) E_{min} = 2.5E+02 eV/amu, E_{max} = 3.7E+04 eV/amu
(4^3D) E_{min} = 2.5E+02 eV/amu, E_{max} = 3.8E+04 eV/amu

<u>Chebyshev Fitting Parameters for Cross Sections</u>

	A0	A1	A2	A3	A4	A5	A6	A7	A8
(4^3S)	-87.3205	1.78746	.468525	-.0126958	.0197677	-.212447	6.18482E-04	-.0153550	.0609384
(4^3P)	-87.9948	1.50655	.179990	-.0510960	.136680	-.148566	-.0170060	-.0293030	.130037
(4^3D)	-86.1432	-.506204	-.0953983	-.353322	-.207541	.0157713	.0723366	.167609	-.197508

The fit represents the (4^3S) cross section with an rms deviation of 9.9%.
The maximum deviation is 20.9% at 1.3E+03 eV/amu.
The fit represents the (4^3P) cross section with an rms deviation of 19.8%.
The maximum deviation is 45.5% at 5.3E+02 eV/amu.
The fit represents the (4^3D) cross section with an rms deviation of 11.3%.
The maximum deviation is 34.8% at 3.0E+02 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + He \rightarrow He(4^3S, 4^3P, 4^3D) + He^+$



Line Emission Cross Sections of Electron Capture Collisions for

$He^+ + Li \rightarrow He(58.4,53.7) + Li^+$

(58.4 nm) Energy eV/amu	Cross Section cm^2	(53.7 nm) Energy eV/amu	Cross Section cm^2
5.0E+02	1.54E-16	5.0E+02	1.29E-18
1.0E+03	2.45E-16	8.4E+02	4.50E-18
1.3E+03	2.63E-16	1.0E+03	6.81E-18
1.5E+03	2.68E-16	1.5E+03	1.05E-17
2.0E+03	2.87E-16	2.0E+03	1.47E-17
2.5E+03	3.31E-16	2.5E+03	2.06E-17
3.0E+03	4.38E-16	3.0E+03	2.93E-17
3.5E+03	5.02E-16	3.5E+03	3.98E-17
4.0E+03	5.21E-16	4.0E+03	4.53E-17
4.5E+03	5.38E-16	4.5E+03	4.77E-17
5.0E+03	5.53E-16	5.0E+03	4.80E-17

References: 267

Accuracy: Unknown

Note: (1) The 3p $^1P^0$ - $1s^2$ ^1S transition gives rise to the 53.7 nm line with the 58.4 line radiation coming from the 2p $^1P^0$ - $1s^2$ ^1S transition.

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

(58.4 nm) E_{min} = 5.0E+02 eV/amu, E_{max} = 5.0E+03 eV/amu
(53.7 nm) E_{min} = 5.0E+02 eV/amu, E_{max} = 5.0E+03 eV/amu

Chebyshev Fitting Parameters for Cross Sections

	A0	A1	A2	A3	A4	A5	A6	A7	A8
(58.4 nm)	-70.6362	-.107414	.390900	.106996	-.435807	.414268	-.355868	.226467	-.0503910
(53.7 nm)	-78.5786	2.06788	-.344623	.0776140	.0263737	-.246781	.172594	-.0896915	.0520900

The fit represents the 58.4 nm cross section with an rms deviation of 1.3%.
The maximum deviation is 2.6% at 2.5E+03 eV/amu.

The fit represents the 53.7 nm cross section with an rms deviation of 0.6%.
The maximum deviation is 1.4% at 4.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + Li \rightarrow He(58.4,53.7) + Li^+$



Electron Capture Cross Sections for
$He^{2+} + H \rightarrow He^{+}(2s) + H^{+}$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.3E+01	6.66E+06	1.01E-23
4.0E+01	8.79E+06	1.93E-22
7.0E+01	1.16E+07	2.96E-21
1.0E+02	1.39E+07	1.64E-20
2.0E+02	1.96E+07	5.15E-19
4.0E+02	2.78E+07	5.71E-18
7.0E+02	3.68E+07	2.19E-17
1.0E+03	4.39E+07	3.92E-17
2.0E+03	6.21E+07	9.06E-17
4.0E+03	8.79E+07	1.35E-16
7.0E+03	1.16E+08	1.57E-16
1.0E+04	1.39E+08	1.63E-16
2.0E+04	1.96E+08	1.40E-16
4.0E+04	2.78E+08	6.73E-17
7.0E+04	3.68E+08	1.95E-17
1.0E+05	4.39E+08	7.15E-18
2.0E+05	6.21E+08	6.07E-19
4.0E+05	8.78E+08	3.70E-20
7.0E+05	1.16E+09	3.23E-21
1.0E+06	1.39E+09	6.01E-22
2.0E+06	1.96E+09	1.56E-23
2.5E+06	2.19E+09	4.77E-24

References: 21, 24, 25, 56, 57, 168, 169, 170, 171, 172, 173

Accuracy: $E < 1.5\times10^3$ eV/amu - unknown; 1.5×10^3 eV/amu $< E < 1.2\times10^5$ eV/amu - 40%; $E > 1.2\times10^5$ eV/amu - unknown

Notes: (1) Data for energies less than 1.5×10^3 eV/amu and greater than 1.2×10^5 eV/amu are theoretical data.
(2) The $He^{+}(2s)$ population is obtained by quenching in an electric field and measuring the additional 30.4 nm photons observed.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 2.3E+01 eV/amu, E_{max} = 2.5E+06 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-89.4268	-.438776	-8.73454	.130314	-.0101210	.0343355	.227699	-.188121	-.130396

The fit represents the above cross sections with an rms deviation of 8.2%.
The maximum deviation is 19.6% at 2.0E+06 eV/amu.
See appendix for Chebyshev fit details.

$He^{2+} + H \rightarrow He^{+}(2s) + H^{+}$



Electron Capture Rate Coefficients for
$He^{2+} + H \rightarrow He^{+}(2s) + H^{+}$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	Equal Temp.	H Temp. (eV) 10.	100.	500.	1000.	5000.	10000.	20000.
7.0E+00	2.97E-16	1.49E-15	2.81E-11	1.30E-09	3.62E-09	1.53E-08	1.93E-08	1.84E-08
1.0E+01	2.08E-15	2.08E-15	2.88E-11	1.30E-09	3.63E-09	1.53E-08	1.93E-08	1.84E-08
2.0E+01	8.24E-14	5.56E-15	3.11E-11	1.31E-09	3.64E-09	1.53E-08	1.93E-08	1.84E-08
4.0E+01	2.04E-12	2.59E-14	3.60E-11	1.33E-09	3.66E-09	1.53E-08	1.93E-08	1.84E-08
7.0E+01	1.70E-11	1.33E-13	4.42E-11	1.37E-09	3.69E-09	1.53E-08	1.93E-08	1.84E-08
1.0E+02	5.34E-11	4.24E-13	5.34E-11	1.40E-09	3.73E-09	1.53E-08	1.93E-08	1.84E-08
2.0E+02	3.27E-10	4.25E-12	9.04E-11	1.51E-09	3.84E-09	1.53E-08	1.93E-08	1.84E-08
4.0E+02	1.29E-09	3.60E-11	1.93E-10	1.74E-09	4.07E-09	1.54E-08	1.93E-08	1.84E-08
7.0E+02	3.03E-09	1.58E-10	4.04E-10	2.09E-09	4.41E-09	1.55E-08	1.93E-08	1.84E-08
1.0E+03	4.75E-09	3.57E-10	6.67E-10	2.44E-09	4.75E-09	1.56E-08	1.94E-08	1.84E-08
2.0E+03	9.49E-09	1.33E-09	1.74E-09	3.61E-09	5.81E-09	1.60E-08	1.94E-08	1.83E-08
4.0E+03	1.53E-08	3.66E-09	4.07E-09	5.81E-09	7.77E-09	1.66E-08	1.95E-08	1.82E-08
7.0E+03	1.88E-08	6.87E-09	7.21E-09	8.66E-09	1.03E-08	1.74E-08	1.96E-08	1.80E-08
1.0E+04	1.96E-08	9.52E-09	9.80E-09	1.10E-08	1.23E-08	1.81E-08	1.96E-08	1.78E-08
2.0E+04	1.71E-08	1.53E-08	1.54E-08	1.60E-08	1.66E-08	1.93E-08	1.95E-08	1.71E-08

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 7.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

H Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
10.	-5.09308E+01	8.87617E+00	-8.39518E-01	-8.53899E-01	3.05164E-01	4.68396E-02	-6.90001E-02
100.	-4.33040E+01	3.46156E+00	6.25707E-01	-3.52271E-01	-1.41060E-01	4.76365E-02	2.50861E-02
500.	-3.92464E+01	1.26410E+00	4.89137E-01	2.19874E-02	-7.43310E-02	-3.13817E-02	1.62905E-03
1000.	-3.79446E+01	7.26279E-01	3.33593E-01	6.06574E-02	-2.84279E-02	-2.45434E-02	-6.72865E-03
5000.	-3.58707E+01	1.02283E-01	5.59105E-02	1.82827E-02	7.91325E-04	-3.18886E-03	-2.27033E-03
10000.	-3.55175E+01	7.32263E-03	2.24624E-03	-1.32249E-03	-2.19045E-03	-1.61111E-03	-8.21561E-04
20000.	-3.56488E+01	-2.68120E-02	-1.78967E-02	-9.56011E-03	-4.19249E-03	-1.52887E-03	-4.56494E-04
Equal Temp.	-4.73886E+01	8.61752E+00	-3.14861E+00	4.07425E-01	-9.02464E-03	-8.82451E-02	3.10401E-02

See appendix for Chebyshev fit details.

$He^{2+} + H \rightarrow He^{+}(2s) + H^{+}$

Maxwellian – Maxwellian



Electron Capture Rate Coefficients for
$H + He^{2+} \rightarrow He^{+}(2s) + H^{+}$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	H Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	2.26E-08	2.75E-08	1.87E-08	7.16E-09	3.14E-09	3.77E-10	1.37E-11
2.0E+00	2.26E-08	2.74E-08	1.87E-08	7.16E-09	3.14E-09	3.77E-10	1.37E-11
4.0E+00	2.26E-08	2.74E-08	1.86E-08	7.16E-09	3.14E-09	3.77E-10	1.37E-11
7.0E+00	2.26E-08	2.74E-08	1.86E-08	7.15E-09	3.14E-09	3.77E-10	1.37E-11
1.0E+01	2.26E-08	2.74E-08	1.86E-08	7.15E-09	3.13E-09	3.77E-10	1.37E-11
2.0E+01	2.26E-08	2.73E-08	1.86E-08	7.15E-09	3.13E-09	3.77E-10	1.37E-11
4.0E+01	2.25E-08	2.72E-08	1.85E-08	7.14E-09	3.13E-09	3.76E-10	1.37E-11
7.0E+01	2.25E-08	2.71E-08	1.85E-08	7.13E-09	3.13E-09	3.76E-10	1.37E-11
1.0E+02	2.25E-08	2.70E-08	1.84E-08	7.13E-09	3.12E-09	3.76E-10	1.37E-11
2.0E+02	2.24E-08	2.69E-08	1.83E-08	7.12E-09	3.12E-09	3.76E-10	1.38E-11
4.0E+02	2.23E-08	2.66E-08	1.81E-08	7.11E-09	3.12E-09	3.76E-10	1.38E-11
7.0E+02	2.23E-08	2.63E-08	1.80E-08	7.10E-09	3.12E-09	3.77E-10	1.38E-11
1.0E+03	2.22E-08	2.61E-08	1.78E-08	7.10E-09	3.12E-09	3.78E-10	1.38E-11
2.0E+03	2.21E-08	2.55E-08	1.75E-08	7.11E-09	3.14E-09	3.81E-10	1.39E-11
4.0E+03	2.20E-08	2.47E-08	1.72E-08	7.18E-09	3.19E-09	3.87E-10	1.41E-11
7.0E+03	2.19E-08	2.38E-08	1.67E-08	7.28E-09	3.27E-09	3.99E-10	1.44E-11
1.0E+04	2.17E-08	2.30E-08	1.64E-08	7.37E-09	3.35E-09	4.11E-10	1.47E-11
2.0E+04	2.11E-08	2.09E-08	1.53E-08	7.54E-09	3.61E-09	4.58E-10	1.57E-11

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

H Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-3.52453E+01	-2.72966E-02	-1.25046E-02	-5.94386E-03	-3.93360E-03	-2.52788E-03	-1.15226E-03
20000.	-3.49559E+01	-1.07234E-01	-5.74126E-02	-2.46077E-02	-9.61541E-03	-3.68199E-03	-1.33552E-03
40000.	-3.56977E+01	-8.13307E-02	-4.02505E-02	-1.59626E-02	-6.46672E-03	-3.09676E-03	-1.64827E-03
70000.	-3.74954E+01	1.58892E-02	1.66995E-02	1.04165E-02	3.58841E-03	-1.14383E-04	-1.18569E-03
100000.	-3.91164E+01	4.26359E-02	3.67186E-02	2.35687E-02	1.10156E-02	3.70448E-03	6.54932E-04
200000.	-4.33336E+01	5.99177E-02	4.72882E-02	2.99739E-02	1.60765E-02	6.77381E-03	2.80729E-03
500000.	-4.99733E+01	4.33562E-02	3.10236E-02	1.83053E-02	8.97845E-03	3.66858E-03	1.27965E-03

See appendix for Chebyshev fit details.

$$H + He^{2+} \rightarrow He^{+}(2s) + H^{+}$$

Beam – Maxwellian



Electron Capture Cross Sections for
$He^{2+} + H \rightarrow He^{+}(2p) + H^{+}$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
3.0E+02	2.41E+07	2.55E-17
4.0E+02	2.78E+07	4.12E-17
5.0E+02	3.11E+07	5.94E-17
7.0E+02	3.68E+07	1.04E-16
9.0E+02	4.17E+07	1.61E-16
1.0E+03	4.39E+07	1.91E-16
1.5E+03	5.38E+07	3.44E-16
2.0E+03	6.21E+07	4.71E-16
2.5E+03	6.95E+07	5.68E-16
3.0E+03	7.61E+07	6.57E-16
4.0E+03	8.79E+07	7.80E-16
5.0E+03	9.82E+07	8.63E-16
7.0E+03	1.16E+08	9.31E-16
9.0E+03	1.32E+08	9.39E-16
1.5E+04	1.70E+08	8.73E-16
2.0E+04	1.96E+08	7.12E-16
2.5E+04	2.20E+08	5.35E-16
3.0E+04	2.41E+08	3.89E-16
4.0E+04	2.78E+08	2.13E-16
5.0E+04	3.11E+08	1.19E-16
6.0E+04	3.40E+08	7.18E-17
8.0E+04	3.93E+08	3.12E-17
1.0E+05	4.39E+08	1.36E-17
1.5E+05	5.38E+08	3.32E-18
2.0E+05	6.21E+08	1.24E-18
2.5E+05	6.94E+08	6.06E-19
3.0E+05	7.61E+08	3.48E-19
4.0E+05	8.78E+08	1.81E-19
5.7E+05	1.05E+09	1.01E-19

References: 169, 170, 172, 173, 174, 175, 176

Accuracy: $1 < E < 1X10^4$ eV/amu - 20%

Note: (1) References 172 and 173 (Ciric et al.) are the only experimental results. Other data are theoretical data which in the energy region of overlap agree well with experiment.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 3.0E+02 eV/amu, E_{max} = 5.7E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-76.4723	-2.98918	-3.20835	.0657861	.450590	.217258	.0371010	-.0738139	-.0123768

The fit represents the above cross sections with an rms deviation of 2.8%.
The maximum deviation is 5.3% at 9.0E+03 eV/amu.
See appendix for Chebyshev fit details.

$He^{2+} + H \rightarrow He^{+}(2p) + H^{+}$



Electron Capture Rate Coefficients for
$He^{2+} + H \rightarrow He^{+}(2p) + H^{+}$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	Equal Temp.	H Temp. (eV) 100.	500.	1000.	5000.	10000.	15000.	20000.
1.0E+02	2.98E-10	2.98E-10	7.39E-09	2.06E-08	8.70E-08	1.03E-07	9.73E-08	8.77E-08
2.0E+02	1.72E-09	4.98E-10	8.00E-09	2.12E-08	8.72E-08	1.03E-07	9.72E-08	8.77E-08
4.0E+02	6.78E-09	1.03E-09	9.26E-09	2.26E-08	8.76E-08	1.03E-07	9.71E-08	8.76E-08
7.0E+02	1.65E-08	2.11E-09	1.12E-08	2.46E-08	8.82E-08	1.03E-07	9.70E-08	8.74E-08
1.0E+03	2.65E-08	3.48E-09	1.32E-08	2.65E-08	8.87E-08	1.03E-07	9.69E-08	8.73E-08
2.0E+03	5.46E-08	9.26E-09	1.99E-08	3.29E-08	9.06E-08	1.03E-07	9.64E-08	8.68E-08
4.0E+03	8.68E-08	2.26E-08	3.29E-08	4.45E-08	9.36E-08	1.02E-07	9.55E-08	8.58E-08
7.0E+03	1.02E-07	4.12E-08	4.98E-08	5.92E-08	9.71E-08	1.02E-07	9.41E-08	8.44E-08
1.0E+04	1.01E-07	5.65E-08	6.33E-08	7.08E-08	9.96E-08	1.01E-07	9.27E-08	8.30E-08
2.0E+04	7.83E-08	8.76E-08	9.06E-08	9.36E-08	1.03E-07	9.73E-08	8.78E-08	7.83E-08

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+02 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

H Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
100.	-3.80891E+01	3.04269E+00	-5.03692E-02	-2.14921E-01	7.02489E-03	1.33936E-02	-3.51323E-03
500.	-3.53407E+01	1.33422E+00	2.37587E-01	-8.16883E-02	-4.11875E-02	4.97820E-04	3.60655E-03
1000.	-3.42278E+01	7.87638E-01	2.02270E-01	-2.32859E-02	-3.10588E-02	-7.16158E-03	1.05223E-03
5000.	-3.23931E+01	8.58442E-02	2.79249E-02	-7.64042E-04	-5.28180E-03	-2.73625E-03	-8.72863E-04
10000.	-3.22047E+01	-1.79017E-02	-1.27932E-02	-7.35687E-03	-3.35962E-03	-1.20009E-03	-3.44007E-04
15000.	-3.23427E+01	-4.11290E-02	-2.23406E-02	-9.19084E-03	-3.13037E-03	-9.49262E-04	-2.08801E-04
20000.	-3.25558E+01	-4.63175E-02	-2.45398E-02	-9.65926E-03	-2.98432E-03	-7.43123E-04	-1.57791E-04
Equal Temp.	-3.61487E+01	2.77246E+00	-1.05020E+00	7.78393E-03	-2.49781E-02	4.65115E-03	2.17778E-03

See appendix for Chebyshev fit details.

$He^{2+} + H \rightarrow He^{+}(2p) + H^{+}$

Maxwellian – Maxwellian



Electron Capture Rate Coefficients for
$H + He^{2+} \rightarrow He^{+}(2p) + H^{+}$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	H Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	1.28E-07	1.40E-07	5.91E-08	1.69E-08	5.97E-09	7.71E-10**	1.23E-10**
2.0E+00	1.28E-07	1.40E-07	5.91E-08	1.69E-08	5.98E-09	7.70E-10**	1.23E-10**
4.0E+00	1.28E-07	1.40E-07	5.91E-08	1.69E-08	5.98E-09	7.70E-10**	1.23E-10**
7.0E+00	1.28E-07	1.39E-07	5.91E-08	1.69E-08	5.98E-09	7.71E-10**	1.23E-10**
1.0E+01	1.29E-07	1.39E-07	5.90E-08	1.69E-08	5.98E-09	7.71E-10**	1.23E-10**
2.0E+01	1.29E-07	1.39E-07	5.90E-08	1.69E-08	5.98E-09	7.71E-10**	1.23E-10**
4.0E+01	1.29E-07	1.39E-07	5.89E-08	1.69E-08	5.99E-09	7.71E-10**	1.23E-10**
7.0E+01	1.29E-07	1.39E-07	5.89E-08	1.69E-08	5.99E-09	7.72E-10*	1.23E-10**
1.0E+02	1.29E-07	1.38E-07	5.88E-08	1.69E-08	6.00E-09	7.72E-10*	1.23E-10**
2.0E+02	1.29E-07	1.38E-07	5.88E-08	1.69E-08	6.02E-09	7.73E-10*	1.23E-10**
4.0E+02	1.28E-07	1.37E-07	5.87E-08	1.70E-08	6.05E-09	7.75E-10*	1.23E-10**
7.0E+02	1.28E-07	1.36E-07	5.87E-08	1.71E-08	6.09E-09	7.77E-10*	1.23E-10**
1.0E+03	1.28E-07	1.34E-07	5.87E-08	1.71E-08	6.14E-09	7.80E-10*	1.23E-10**
2.0E+03	1.28E-07	1.31E-07	5.89E-08	1.74E-08	6.28E-09	7.88E-10*	1.23E-10**
4.0E+03	1.26E-07	1.25E-07	5.93E-08	1.78E-08	6.53E-09	8.03E-10*	1.21E-10**
7.0E+03	1.22E-07	1.18E-07	5.95E-08	1.85E-08	6.88E-09	8.27E-10*	1.18E-10**
1.0E+04	1.19E-07	1.13E-07	5.93E-08	1.92E-08	7.22E-09	8.52E-10*	1.14E-10**
2.0E+04	1.10E-07	9.85E-08	5.73E-08	2.12E-08	8.38E-09	9.47E-10*	1.06E-10**

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

H Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-3.17882E+01	-4.92276E-02	-3.82539E-02	-2.43393E-02	-1.22728E-02	-4.54785E-03	-7.79239E-04
20000.	-3.17232E+01	-1.30716E-01	-8.13104E-02	-4.04778E-02	-1.57809E-02	-3.98767E-03	3.36890E-05
40000.	-3.32955E+01	-5.40047E-03	-1.99166E-03	-4.45358E-03	-6.27891E-03	-5.14199E-03	-2.92410E-03
70000.	-3.57028E+01	8.02848E-02	5.54250E-02	3.03851E-02	1.33090E-02	4.49405E-03	7.72971E-04
100000.	-3.77338E+01	1.19799E-01	7.98553E-02	4.24318E-02	1.81018E-02	6.30836E-03	2.00635E-03
200000.	-4.18916E+01	6.78830E-02	4.78977E-02	2.81571E-02	1.45668E-02	6.26534E-03	2.69138E-03
500000.	-4.56871E+01	-4.63255E-02	-3.72726E-02	-2.45372E-02	-1.26363E-02	-4.18959E-03	7.18096E-04

See appendix for Chebyshev fit details.

$$H + He^{2+} \rightarrow He^{+}(2p) + H^{+}$$

Beam – Maxwellian



Line Emission Cross Sections of Electron Capture Collisions for

$$He^{2+} + H \rightarrow He^{+}(Ly\alpha, Ly\beta, Ly\gamma, H\alpha) + H^{+}$$

(Ly_α)		(Ly_β)		(Ly_γ)		(H_α)	
Energy eV/amu	Cross Section cm^2	Energy eV/amu	Cross Section cm^2	Energy eV/amu	Cross Section cm^2	Energy eV/amu	Cross Section cm^2
1.2E+03	2.68E-16	1.6E+03	6.37E-18	4.8E+03	2.39E-18	2.8E+03	1.99E-17
1.5E+03	3.21E-16	2.2E+03	1.30E-17	5.0E+03	2.45E-18	2.9E+03	2.03E-17
2.2E+03	4.90E-16	2.9E+03	1.93E-17	5.7E+03	2.75E-18	3.6E+03	2.36E-17
2.9E+03	6.35E-16	3.6E+03	2.18E-17	6.4E+03	3.08E-18	4.3E+03	2.73E-17
3.6E+03	7.49E-16	4.3E+03	2.32E-17	7.2E+03	3.48E-18	5.0E+03	3.17E-17
4.3E+03	8.30E-16	5.0E+03	2.39E-17	7.9E+03	3.96E-18	5.7E+03	3.70E-17
5.0E+03	8.82E-16	5.7E+03	2.47E-17	8.6E+03	4.47E-18	6.5E+03	4.22E-17
5.7E+03	9.11E-16	6.5E+03	2.54E-17	9.1E+03	4.89E-18	7.2E+03	4.70E-17
6.5E+03	9.31E-16	7.2E+03	2.64E-17	9.3E+03	5.08E-18	7.9E+03	5.03E-17
7.2E+03	9.50E-16	7.9E+03	2.75E-17	1.0E+04	5.84E-18	8.6E+03	5.24E-17
7.9E+03	9.59E-16	8.6E+03	2.87E-17			9.3E+03	5.35E-17
8.6E+03	9.73E-16	9.3E+03	3.04E-17			1.0E+04	5.40E-17
9.3E+03	9.87E-16	1.0E+04	3.24E-17				
1.0E+04	9.91E-16						

References: 173

Accuracy: 30%

Notes: (1) Transition wavelengths are: Lyman-α - (2p->1s) 30.4 nm; Lyman-β - (3p->1s) 25.6 nm; Lyman-γ - (4p->1s) 24.3 nm; Balmer-α - (n=3->2) 164 nm.
(2) The Balmer-β emission cannot be measured since the transition has the same wave length as the Lyman-α radiation.

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

(Ly_α) E_{min} = 1.2E+03 eV/amu, E_{max} = 1.0E+04 eV/amu
(Ly_β) E_{min} = 1.6E+03 eV/amu, E_{max} = 1.0E+04 eV/amu
(Ly_γ) E_{min} = 4.8E+03 eV/amu, E_{max} = 1.0E+04 eV/amu
(H_α) E_{min} = 2.8E+03 eV/amu, E_{max} = 1.0E+04 eV/amu

Chebyshev Fitting Parameters for Cross Sections

	A0	A1	A2	A3	A4	A5	A6	A7	A8
(Ly_α)	-70.0765	.658545	-.176878	-.00575199	.0163374	.00346973	-.00221833	-.00217295	-4.61984E-04
(Ly_β)	-77.0977	.692364	-.245444	.142376	-.00794442	-.0119412	.0163687	-.00929968	.00446139
(Ly_γ)	-80,3639	.441485	.0512890	.00405517	9.45294E-04	.00161819	.00182462	-4.78184E-04	-6.34056E-04
(H_α)	-75.8681	.546644	-.0142448	-.0516937	-.0137147	.00553121	.00525928	-.00130955	9.09970E-05

The fit represents the (Ly_α) cross section with an rms deviation of 0.1%.
The maximum deviation is 0.2% at 7.9E+03 eV/amu.
The fit represents the (Ly_β) cross section with an rms deviation of 0.2%.
The maximum deviation is 0.3% at 8.6E+03 eV/amu.
The fit represents the (Ly_γ) cross section with an rms deviation of 0.1%.
The maximum deviation is 0.2% at 8.6E+03 eV/amu.
The fit represents the (H_α) cross section with an rms deviation of 0.1%.
The maximum deviation is 0.2% at 6.4E+03 eV/amu.

See appendix for Chebyshev fit details.

$$He^{2+} + H \rightarrow He^{+}(Ly_{\alpha}, Ly_{\beta}, Ly_{\gamma}, H_{\alpha}) + H^{+}$$



Electron Capture Cross Sections for

$He^{2+} + H \rightarrow He^{+}(3p,3s+3d,4p) + H^{+}$

(3p) Energy eV/amu	(3p) Cross Section cm^2	(3s+3d) Energy eV/amu	(3s+3d) Cross Section cm^2	(4p) Energy eV/amu	(4p) Cross Section cm^2
1.6E+03	7.71E-18	2.5E+03	1.78E-17	5.0E+03	2.49E-18
2.0E+03	1.46E-17	3.0E+03	1.87E-17	5.5E+03	2.73E-18
2.5E+03	2.09E-17	3.5E+03	2.01E-17	6.0E+03	2.99E-18
3.0E+03	2.40E-17	4.0E+03	2.23E-17	7.0E+03	3.56E-18
3.5E+03	2.63E-17	4.5E+03	2.51E-17	8.0E+03	4.22E-18
4.0E+03	2.83E-17	5.0E+03	2.81E-17	9.0E+03	4.96E-18
4.5E+03	2.97E-17	6.0E+03	3.49E-17	1.0E+04	6.03E-18
5.5E+03	3.20E-17	7.0E+03	4.15E-17		
6.0E+03	3.29E-17	8.0E+03	4.63E-17		
7.0E+03	3.43E-17	9.0E+03	4.89E-17		
8.0E+03	3.57E-17	1.0E+04	4.97E-17		
9.0E+03	3.67E-17				
1.0E+04	3.74E-17				

References: 172, 173

Accuracy: 30%

Notes: (1) The 3s and 3d states could not be determined separately due to the degeneracy of states in He^{+}.
(2) Capture into the ground state is negligible at low energies for a H target.
(3) The cross section (σ_i) for capture into state i is determined by measuring the emission cross section (σ_{ij}) for a transition from state i to j, correcting for any branching from state i and correcting for cascading from upper states.

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

(3p) E_{min} = 1.6E+03 eV/amu, E_{max} = 1.0E+04 eV/amu
(3s+3d) E_{min} = 2.5E+03 eV/amu, E_{max} = 1.0E+04 eV/amu
(4p) E_{min} = 5.0E+03 eV/amu, E_{max} = 1.0E+04 eV/amu

Chebyshev Fitting Parameters for Cross Sections

	A0	A1	A2	A3	A4	A5	A6	A7	A8
(3p)	-76.6256	.667773	-.257128	.115045	-.0496129	.0142615	9.03999E-04	-.00746439	.00421196
(3s+3d)	-76.1346	.575824	.0291040	-.0715718	-.0133196	.00539953	-.00104695	.00377347	-.00137059
(4p)	-80.2658	.432003	.0343796	.00725815	.00469173	.00296970	.00183023		

The fit represents the (3p) cross section with an rms deviation of 0.1%.
The maximum deviation is 0.3% at 4.5E+03 eV/amu.
The fit represents the (3s+3d) cross section with an rms deviation of 0.1%.
The maximum deviation is 0.1% at 4.0E+03 eV/amu.
The fit represents the (4p) cross section with an rms deviation of 0.0%.
The maximum deviation is 0.0%

See appendix for Chebyshev fit details.

$He^{2+} + H \rightarrow He^{+}(3p,3s+3d,4p) + H^{+}$



Electron Capture Cross Sections for

$He^{2+} + H_2 \rightarrow He^{+}(2s,2p)$

$He^{+}(2p)$ Energy eV/amu	Cross Section cm^2
1.2E+03	4.90E-17
1.5E+03	5.94E-17
2.0E+03	8.22E-17
3.0E+03	1.35E-16
4.0E+03	2.15E-16
5.0E+03	3.16E-16
7.0E+03	4.79E-16
9.0E+03	6.18E-16
1.0E+04	7.08E-16

$He^{+}(2s)$ Energy eV/amu	Cross Section cm^2
2.0E+03	7.99E-18
3.0E+03	1.97E-17
4.0E+03	3.33E-17
5.0E+03	4.56E-17
5.5E+03	5.12E-17
6.0E+03	5.76E-17
7.0E+03	7.46E-17
7.5E+03	8.86E-17
8.0E+03	9.88E-17
9.0E+03	1.17E-16
1.0E+04	1.28E-16
1.5E+04	1.37E-16

References: $He^{+}(2s)$ - 56, 145, 177; $He^{+}(2p)$ - 172, 173

Accuracy: $He^{+}(2s)$ - 60%; $He^{+}(2p)$ - 30%

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

$He^{+}(2p)$ E_{min} = 1.2E+03 eV/amu, E_{max} = 1.0E+04 eV/amu
$He^{+}(2s)$ E_{min} = 2.0E+03 eV/amu, E_{max} = 2.0E+04 eV/amu

Chebyshev Fitting Parameters for Cross Sections

	A0	A1	A2	A3	A4	A5	A6	A7	A8
$He^{+}(2p)$	-72.5057	1.39004	.0296375	-.0736032	-.0100517	.0209333	.0217835	-.00206200	-.00793171
$He^{+}(2s)$	-75.1849	1.42417	-.425662	-.0917408	-.0265286	.0447348	.0526597		

The fit represents the $He^{+}(2p)$ cross section with an rms deviation of 0.0%.
The maximum deviation is 0.0%

The fit represents the $He^{+}(2s)$ cross section with an rms deviation of 2.4%.
The maximum deviation is 4.6% at 1.3E+04 eV/amu.

See appendix for Chebyshev fit details.

$He^{2+} + H_2 \rightarrow He^{+}(2s, 2p)$



Line Emission Cross Sections of Electron Capture Collisions for

$$He^{2+} + H_2 \rightarrow He^{+}(Ly_\alpha, Ly_\beta, Ly_\gamma, H_\alpha)$$

Ly_α Energy eV/amu	Ly_α Cross Section cm^2	H_α Energy eV/amu	H_α Cross Section cm^2	Ly_β Energy eV/amu	Ly_β Cross Section cm^2	Ly_γ Energy eV/amu	Ly_γ Cross Section cm^2
1.3E+03	4.82E-17	2.5E+03	6.44E-18	1.7E+03	1.77E-18	5.0E+03	4.93E-19
1.5E+03	5.62E-17	2.9E+03	7.44E-18	2.2E+03	2.52E-18	5.7E+03	7.56E-19
2.2E+03	9.21E-17	3.6E+03	9.63E-18	2.9E+03	3.75E-18	6.5E+03	1.15E-18
2.9E+03	1.37E-16	4.3E+03	1.18E-17	3.6E+03	5.18E-18	7.2E+03	1.77E-18
3.6E+03	1.90E-16	5.0E+03	1.44E-17	4.3E+03	6.94E-18	7.9E+03	2.32E-18
4.3E+03	2.58E-16	5.7E+03	1.75E-17	5.0E+03	8.79E-18	8.6E+03	2.68E-18
5.0E+03	3.25E-16	6.4E+03	2.10E-17	5.7E+03	1.06E-17	9.3E+03	2.94E-18
5.7E+03	3.94E-16	7.2E+03	2.48E-17	6.4E+03	1.24E-17	1.0E+04	3.15E-18
6.5E+03	4.59E-16	7.9E+03	2.89E-17	7.2E+03	1.42E-17		
7.2E+03	5.24E-16	8.6E+03	3.23E-17	7.9E+03	1.51E-17		
7.9E+03	5.88E-16	9.3E+03	3.49E-17	8.6E+03	1.55E-17		
8.6E+03	6.56E-16	1.0E+04	3.65E-17	9.3E+03	1.56E-17		
9.3E+03	7.36E-16			1.0E+04	1.52E-17		
1.0E+04	8.06E-16						

References: 173

Accuracy: 30%

Notes: (1) Transition wavelengths are: Lyman-α (2p->1s) 30.4 nm; Lyman-β (3p->1s) 25.6 nm; Lyman-γ (4p->1s) 24.3 nm; Balmer-α (H_α, n=3->2) 164 nm.
(2) The Balmer-β radiation cannot be measured since the transition has the same wave length as the Lyman-α radiation.

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

(Ly_α) E_{min} = 1.3E+03 eV/amu, E_{max} = 1.0E+04 eV/amu
(H_α) E_{min} = 2.5E+03 eV/amu, E_{max} = 1.0E+04 eV/amu
(Ly_β) E_{min} = 1.7E+03 eV/amu, E_{max} = 1.0E+04 eV/amu
(Ly_γ) E_{min} = 5.0E+03 eV/amu, E_{max} = 1.0E+04 eV/amu

Chebyshev Fitting Parameters for Cross Sections

	A0	A1	A2	A3	A4	A5	A6	A7	A8
(Ly_α)	-72.3892	1.45574	.0140540	-.0435817	.00709587	.00345598	.0136342	-.00698786	-.00272240
(H_α)	-77.4506	.904442	.0279735	-.0253175	-.0268277	-.00920018	.00430881	-.00254202	.00325844
(Ly_β)	-79.3508	1.17675	-.116050	-.0780599	-.0348000	-.0102977	.0150173	-.0123859	.0116818
(Ly_γ)	-82.1736	.953689	-.126724	-.0689492	.0127445	.0372528	-.0256612	.00533416	

The fit represents the Ly_α cross section with an rms deviation of 0.3%.
The maximum deviation is 0.6% at 3.6E+03 eV/amu.

The fit represents the H_α cross section with an rms deviation of 0.1%.
The maximum deviation is 0.2% at 7.1E+03 eV/amu.

The fit represents the Ly_β cross section with an rms deviation of 0.4%.
The maximum deviation is 0.7% at 8.6E+03 eV/amu.

The fit represents the Ly_γ cross section with an rms deviation of 0.0%.
The maximum deviation is 0.0%.
See appendix for Chebyshev fit details.

$$He^{2+} + H_2 \rightarrow He^{+}(Ly_{\alpha}, Ly_{\beta}, Ly_{\gamma}, H_{\alpha})$$

Cross Section vs. Energy



Electron Capture Cross Sections for

$He^{2+} + H_2 \rightarrow He^{+}(3p,3s+3d,4p)$

(3p)		(3s+3d)		(4p)	
Energy eV/amu	Cross Section cm^2	Energy eV/amu	Cross Section cm^2	Energy eV/amu	Cross Section cm^2
2.5E+03	6.70E-18	1.9E+03	1.96E-18	5.0E+03	5.78E-19
3.0E+03	7.05E-18	2.3E+03	2.64E-18	5.8E+03	9.10E-19
3.5E+03	9.24E-18	3.0E+03	4.59E-18	6.5E+03	1.39E-18
4.3E+03	1.17E-17	3.5E+03	6.18E-18	7.3E+03	2.10E-18
5.0E+03	1.45E-17	4.3E+03	7.72E-18	8.0E+03	2.82E-18
5.8E+03	1.75E-17	5.0E+03	9.58E-18	8.5E+03	3.30E-18
6.5E+03	2.05E-17	5.8E+03	1.17E-17	9.3E+03	3.64E-18
7.3E+03	2.35E-17	6.5E+03	1.38E-17	1.0E+04	3.84E-18
8.0E+03	2.65E-17	7.3E+03	1.56E-17		
8.5E+03	2.96E-17	8.0E+03	1.68E-17		
9.3E+03	3.25E-17	8.5E+03	1.75E-17		
1.0E+04	3.54E-17	9.3E+03	1.78E-17		
		1.0E+04	1.71E-17		

References: 173

Accuracy: 30%

Notes: (1) The 3s and 3d states could not be determined separately due to the degeneracy of states in He^{+}.
(2) The cross section for capture into state i is determined by measuring the emission cross section (σ_{ij}) for a transition from state i to j, correcting for any branching from state i and correcting for cascading from upper states.

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

(3p) E_{min} = 2.5E+03 eV/amu, E_{max} = 1.0E+04 eV/amu
(3s+3d) E_{min} = 1.9E+03 eV/amu, E_{max} = 1.0E+04 eV/amu
(4p) E_{min} = 5.0E+03 eV/amu, E_{max} = 1.0E+04 eV/amu

Chebyshev Fitting Parameters for Cross Sections

	A0	A1	A2	A3	A4	A5	A6	A7	A8
(3p)	-77.5134	.877384	.0236181	-.0330936	.0119138	-.00648625	.00667714	-.00553227	.00234918
(3s+3d)	-78.9402	1.10256	-.166612	-.0123653	-.0567272	-.00713119	.00939783	4.30617E-04	-.00609836
(4p)	-81.8167	.998271	-.145948	-.0608582	.0111762	.0174494	-.00474599	-.00803524	

The fit represents the 3p cross section with an rms deviation of 0.1%.
The maximum deviation is 0.1% at 7.9E+03 eV/amu.

The fit represents the 3s+3d cross section with an rms deviation of 0.2%.
The maximum deviation is 0.4% at 8.6E+03 eV/amu.

The fit represents the 4p cross section with an rms deviation of 0.0%.
The maximum deviation is 0.0%.

See appendix for Chebyshev fit details.

$He^{2+} + H_2 \rightarrow He^{+}(3p, 3s+3d, 4p)$



Electron Capture Cross Sections for
$He^{2+} + He \rightarrow He^{+}(2s) + He^{+}$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.0E+03	6.21E+07	2.79E-18
3.0E+03	7.61E+07	3.30E-18
4.0E+03	8.79E+07	4.04E-18
5.0E+03	9.82E+07	5.32E-18
7.0E+03	1.16E+08	9.18E-18
9.0E+03	1.32E+08	1.37E-17
1.0E+04	1.39E+08	1.62E-17
1.5E+04	1.70E+08	3.04E-17
2.0E+04	1.96E+08	3.75E-17
2.5E+04	2.20E+08	3.90E-17
3.0E+04	2.41E+08	3.84E-17
4.0E+04	2.78E+08	3.36E-17
6.0E+04	3.40E+08	2.21E-17
8.0E+04	3.93E+08	1.40E-17
1.0E+05	4.39E+08	8.65E-18
1.5E+05	5.38E+08	2.83E-18
2.0E+05	6.21E+08	1.26E-18
2.5E+05	6.94E+08	6.87E-19
3.0E+05	7.61E+08	4.10E-19
4.0E+05	8.78E+08	1.75E-19
4.7E+05	9.52E+08	1.27E-19

References: 56, 145, 177, 178, 179, 180, 181, 182

Accuracy: $E < 2x10^4$ eV/amu - 50%; $E > 2x10^4$ eV/amu - unknown

Notes: (1) Experimental data (ref. 56, 145, 177) extends only to $2x10^4$ eV/amu. In the energy region of overlap ($7x10^3$ eV/amu-$2x10^4$ eV/amu) between experimental data and theoretical data (ref. 178, 180, 181) the theoretical data is greater than experimental by 1.5-2.5 orders of magnitude. Data above $2x10^4$ eV/amu should be used with caution.
(2) No experimental data exists for $He^{2+} + He \rightarrow He^{+}(2p)$.

For a Chebyshev fit of the above cross sections it is necessary to use the following parameters.
E_{min} = 2.0E+03 eV/amu, E_{max} = 4.7E+05 eV/amu

Chebyshev Fitting Parameters for Cross Sections

A0	A1	A2	A3	A4	A5	A6	A7	A8
-80.4752	-1.31324	-2.07504	-.281865	.383911	5.87536E-03	-.0220039	.0340012	-.0262370

The fit represents the above cross sections with an rms deviation of 2.7%.
The maximum deviation is 5.6% at 4.0E+05 eV/amu.
See appendix for Chebyshev fit details.

$He^{2+} + He \rightarrow He^{+}(2s) + He^{+}$

Cross Section vs. Energy



Electron Capture Rate Coefficients for
$He^{2+} + He \rightarrow He^{+}(2s) + He^{+}$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	Equal Temp.	He Temp. (eV) 500.	1000.	2000.	4000.	7000.	10000.	20000.
7.0E+02	1.96E-12	7.88E-13	5.12E-12	2.78E-11	1.11E-10	3.08E-10	5.88E-10	1.84E-09
1.0E+03	1.01E-11	2.81E-12	1.01E-11	3.76E-11	1.27E-10	3.32E-10	6.20E-10	1.88E-09
2.0E+03	7.73E-11	2.20E-11	3.76E-11	7.73E-11	1.86E-10	4.20E-10	7.31E-10	2.02E-09
4.0E+03	3.32E-10	1.01E-10	1.27E-10	1.86E-10	3.32E-10	6.20E-10	9.70E-10	2.28E-09
7.0E+03	9.70E-10	2.92E-10	3.32E-10	4.20E-10	6.20E-10	9.70E-10	1.35E-09	2.65E-09
1.0E+04	1.75E-09	5.67E-10	6.20E-10	7.31E-10	9.70E-10	1.35E-09	1.75E-09	3.01E-09
2.0E+04	4.01E-09	1.82E-09	1.88E-09	2.02E-09	2.28E-09	2.65E-09	3.01E-09	4.01E-09

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 7.0E+02 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
500.	-4.71093E+01	3.78408E+00	-4.25226E-01	1.09485E-01	-1.88308E-02	-2.18471E-02	1.64914E-03
1000.	-4.59288E+01	2.96506E+00	-6.94375E-02	4.01519E-03	-9.10887E-03	-1.53339E-02	-4.97257E-04
2000.	-4.46311E+01	2.18675E+00	1.67647E-01	-3.86105E-02	-1.57224E-02	-6.51960E-03	-3.71278E-04
4000.	-4.32631E+01	1.54495E+00	2.39004E-01	-3.13418E-02	-1.96927E-02	-2.77831E-03	6.46638E-04
7000.	-4.20414E+01	1.09232E+00	2.10178E-01	-1.36333E-02	-1.49539E-02	-2.16792E-03	5.39745E-04
10000.	-4.12037E+01	8.21778E-01	1.74596E-01	-3.48824E-03	-1.10046E-02	-2.36544E-03	-2.58516E-04
20000.	-3.96301E+01	3.84302E-01	9.69796E-02	5.67825E-03	-3.92022E-03	-1.53584E-03	-5.16122E-04
Equal Temp.	-4.49085E+01	3.65780E+00	-6.02220E-01	1.33794E-01	-9.85265E-02	2.12708E-02	8.09123E-03

See appendix for Chebyshev fit details.

$He^{2+} + He \rightarrow He^{+}(2s) + He^{+}$

Maxwellian — Maxwellian



Electron Capture Rate Coefficients for
$He + He^{2+} \rightarrow He^{+}(2s) + He^{+}$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	60000.	80000.	100000.	200000.
1.0E+00	2.25E-09	7.35E-09	9.33E-09	7.51E-09	5.50E-09*	3.80E-09*	7.83E-10**
2.0E+00	2.25E-09	7.36E-09	9.32E-09	7.51E-09	5.50E-09*	3.80E-09*	7.83E-10**
4.0E+00	2.25E-09	7.35E-09	9.32E-09	7.51E-09	5.49E-09*	3.80E-09*	7.83E-10**
7.0E+00	2.25E-09	7.35E-09	9.31E-09	7.51E-09	5.49E-09	3.79E-09*	7.83E-10**
1.0E+01	2.25E-09	7.34E-09	9.31E-09	7.50E-09	5.49E-09	3.79E-09*	7.83E-10**
2.0E+01	2.26E-09	7.33E-09	9.30E-09	7.50E-09	5.49E-09	3.79E-09*	7.83E-10**
4.0E+01	2.26E-09	7.32E-09	9.29E-09	7.49E-09	5.48E-09	3.79E-09*	7.83E-10**
7.0E+01	2.27E-09	7.31E-09	9.27E-09	7.48E-09	5.48E-09	3.78E-09*	7.83E-10**
1.0E+02	2.28E-09	7.30E-09	9.26E-09	7.47E-09	5.47E-09	3.78E-09	7.83E-10*
2.0E+02	2.31E-09	7.28E-09	9.23E-09	7.45E-09	5.46E-09	3.78E-09	7.84E-10*
4.0E+02	2.37E-09	7.26E-09	9.19E-09	7.43E-09	5.45E-09	3.77E-09	7.85E-10*
7.0E+02	2.45E-09	7.23E-09	9.14E-09	7.40E-09	5.43E-09	3.76E-09	7.87E-10*
1.0E+03	2.54E-09	7.22E-09	9.10E-09	7.38E-09	5.42E-09	3.76E-09	7.88E-10*
2.0E+03	2.79E-09	7.17E-09	8.99E-09	7.32E-09	5.40E-09	3.76E-09	7.93E-10*
4.0E+03	3.20E-09	7.11E-09	8.81E-09	7.23E-09	5.36E-09	3.76E-09	8.02E-10*
7.0E+03	3.68E-09	7.05E-09	8.57E-09	7.11E-09	5.31E-09	3.77E-09	8.15E-10*
1.0E+04	4.06E-09	7.01E-09	8.35E-09	6.98E-09	5.25E-09	3.77E-09	8.29E-10*
2.0E+04	4.94E-09	6.91E-09	7.72E-09	6.58E-09	5.08E-09	3.75E-09	8.76E-10*

Accuracy: * - Possible Error Greater Than 10%
** - Possible Error Greater Than 100%

Notes: For Chebyshev fits of the above rate coefficients it is necessary to use the following parameters.
E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Chebyshev Fitting Parameters for Rate Coefficients

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-3.94248E+01	3.34599E-01	1.94015E-01	6.86806E-02	4.95694E-03	-9.99564E-03	-6.10649E-03
20000.	-3.74937E+01	-2.78645E-02	-1.11501E-02	-3.09862E-03	-9.33585E-04	8.71220E-06	1.09550E-04
40000.	-3.70648E+01	-6.95474E-02	-4.17914E-02	-2.13044E-02	-9.81107E-03	-3.95679E-03	-1.20270E-03
60000.	-3.74705E+01	-4.74430E-02	-2.81261E-02	-1.48208E-02	-7.60963E-03	-3.68669E-03	-1.56303E-03
80000.	-3.80745E+01	-2.90089E-02	-1.58763E-02	-8.00256E-03	-4.24777E-03	-2.17637E-03	-8.77601E-04
100000.	-3.87879E+01	-5.76796E-03	1.27600E-04	8.95128E-04	-2.39349E-04	-7.73294E-04	-5.69452E-04
200000.	-4.18939E+01	3.77781E-02	2.64380E-02	1.53165E-02	7.52659E-03	3.09706E-03	1.14409E-03

See appendix for Chebyshev fit details.

$He + He^{2+} \rightarrow He^{+}(2s) + He^{+}$



Electron Capture Cross Sections for

$He^{2+} + Li \rightarrow He^{+}(2p,3p,4p,3s+3d,4s+4d) + Li^{+}$

(2p)		(3p)		(4p)		(3s+3d)		(4s+4d)	
Energy eV/amu	Cross Section cm^2	Energy eV/amu	Cross Section cm^2	Energy eV/amu	Cross Section cm^2	Energy eV/amu	Cross Section cm^2	Energy eV/amu	Cross Section cm^2
1.1E+04	3.70E-16	5.6E+02	4.96E-15	1.3E+03	2.99E-16	3.8E+03	6.02E-15	3.8E+03	9.77E-16
1.5E+04	2.71E-16	7.1E+02	4.56E-15	1.5E+03	3.54E-16	5.0E+03	5.31E-15	5.0E+03	1.39E-15
2.0E+04	1.97E-16	9.1E+02	4.20E-15	2.0E+03	4.29E-16	7.0E+03	4.17E-15	6.0E+03	1.51E-15
3.0E+04	1.15E-16	1.5E+03	3.55E-15	3.0E+03	5.30E-16	1.0E+04	2.64E-15	7.0E+03	1.50E-15
3.8E+04	5.90E-17	2.0E+03	3.23E-15	4.0E+03	6.08E-16	1.5E+04	9.31E-16	1.0E+04	1.28E-15
		3.0E+03	2.82E-15	6.0E+03	7.51E-16	2.0E+04	3.60E-16	1.5E+04	6.27E-16
		5.0E+03	2.24E-15	7.0E+03	8.00E-16	2.5E+04	1.50E-16	2.0E+04	3.38E-16
		7.0E+03	1.68E-15	8.6E+03	8.36E-16	3.0E+04	7.39E-17	2.5E+04	1.89E-16
		1.0E+04	9.50E-16	1.0E+04	7.94E-16	3.8E+04	2.19E-17	3.0E+04	1.18E-16
		1.5E+04	3.53E-16	1.5E+04	4.72E-16			3.7E+04	6.50E-17
		2.0E+04	1.45E-16	2.0E+04	2.34E-16				
		2.5E+04	5.49E-17	2.5E+04	9.89E-17				
		3.0E+04	1.77E-17	3.0E+04	4.39E-17				
		3.2E+04	1.08E-17	3.8E+04	1.44E-17				

References: 240, 243, 245, 259, 262, 263, 264

Accuracy: unknown

Notes: (1) Cross sections into excited states are determined by measuring the emission from the state and subtracting cascade transitions from higher states and making corrections for the branching ratios.
(2) It is believed that the data are accurate to within 60%.
(3) References 263 and 264 contain theoretical data that were not included in compiling the present data.

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

(2p) E_{min} = 1.1E+04 eV/amu, E_{max} = 3.8E+04 eV/amu
(3p) E_{min} = 5.6E+02 eV/amu, E_{max} = 3.2E+04 eV/amu
(4p) E_{min} = 1.3E+03 eV/amu, E_{max} = 3.8E+04 eV/amu
(3s+3d) E_{min} = 3.8E+03 eV/amu, E_{max} = 3.8E+04 eV/amu
(4s+4d) E_{min} = 3.8E+03 eV/amu, E_{max} = 3.7E+04 eV/amu

Chebyshev Fitting Parameters for Cross Sections

	A0	A1	A2	A3	A4	A5	A6	A7	A8
(2p)	-72.5775	-.854328	-.132073	-.0636576	-.0301663				
(3p)	-69.3960	-2.53366	-1.15152	-.477709	-.109491	-.0224733	-.0359280	-.0364469	-.0124596
(4p)	-71.9553	-1.18550	-1.26810	-.417040	-.0333091	.0822485	.0142456	.00155260	.00317940
(3s+3d)	-69.5364	-2.77071	-.724373	-.0299068	.0383153	-.0274998	-.0511074	.0199350	-.0465182
(4s+4d)	-70.4843	-1.46194	-.666629	.0919477	.0166653	-.00399733	-.0199710	.0194456	-.00484478

The fit represents the above cross sections with an rms deviation of 0.0%.
The maximum deviation is 0.0%.
The fit represents the above cross sections with an rms deviation of 0.8%.
The maximum deviation is 1.5% at 3.0E+04 eV/amu.
The fit represents the above cross sections with an rms deviation of 1.2%.
The maximum deviation is 2.7% at 1.5E+04 eV/amu.
The fit represents the above cross sections with an rms deviation of 0.0%.
The maximum deviation is 0.0%.
The fit represents the above cross sections with an rms deviation of 1.1%.
The maximum deviation is 1.7% at 7.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$He^{2+} + Li \rightarrow He^{+}(2p, 3p, 4p, 3s+3d, 4s+4d) + Li^{+}$



Line Emission Cross Sections of Electron Capture Collisions for

$$He^{2+} + Li \rightarrow He^{+}(Ly_{\alpha}, Ly_{\beta}, Ly_{\gamma}) + Li^{+}$$

(Ly_{α}) Energy eV/amu	Cross Section cm^2	(Ly_{β}) Energy eV/amu	Cross Section cm^2	(Ly_{γ}) Energy eV/amu	Cross Section cm^2
8.1E-01	4.72E-16	8.3E-01	3.71E-16	1.3E+03	2.55E-16
1.0E+00	4.83E-16	1.0E+00	4.42E-16	1.5E+03	3.01E-16
1.5E+00	5.53E-16	1.5E+00	6.23E-16	2.0E+03	3.77E-16
2.0E+00	7.30E-16	2.0E+00	7.70E-16	3.0E+03	4.69E-16
2.5E+00	9.40E-16	3.0E+00	8.83E-16	4.0E+03	5.31E-16
3.0E+00	1.12E-15	4.0E+00	8.42E-16	6.0E+03	6.25E-16
4.0E+00	1.26E-15	5.0E+00	7.65E-16	8.0E+03	6.88E-16
6.0E+00	1.28E-15	6.0E+00	7.02E-16	9.1E+03	6.94E-16
8.0E+00	1.21E-15	8.0E+00	5.99E-16	1.0E+04	6.74E-16
1.0E+01	1.13E-15	1.0E+01	5.32E-16	1.5E+04	4.06E-16
1.3E+01	1.04E-15	1.4E+01	4.43E-16	2.0E+04	2.02E-16
5.4E+02	7.23E-15	5.4E+02	4.68E-15	2.5E+04	8.81E-17
8.0E+02	7.35E-15	7.0E+02	4.04E-15	3.0E+04	4.00E-17
1.0E+03	7.30E-15	9.0E+02	3.59E-15	3.5E+04	1.98E-17
1.5E+03	7.31E-15	9.9E+02	3.42E-15		
2.0E+03	7.27E-15	1.5E+03	2.93E-15		
3.0E+03	7.07E-15	2.0E+03	2.75E-15		
5.0E+03	6.77E-15	3.0E+03	2.67E-15		
6.9E+03	5.67E-15	4.0E+03	2.63E-15		
1.0E+04	4.10E-15	5.0E+03	2.51E-15		
1.5E+04	2.04E-15	7.0E+03	2.03E-15		
2.0E+04	1.00E-15	9.0E+03	1.49E-15		
3.0E+04	3.39E-16	1.0E+04	1.29E-15		
3.8E+04	1.79E-16	1.5E+04	5.97E-16		
		2.0E+04	3.02E-16		
		3.0E+04	6.52E-17		
		3.8E+04	2.20E-17		

References: 240, 243, 245, 259, 262

Accuracy: unknown

Notes: (1) On purpose, no attempt has been made to extrapolate the data between the low and high energy regions.
(2) The emission cross section is defined as the photon emission from state nl with no correction for cascading from higher states.

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

(Ly_{α}) E_{min} = 8.1E-01 eV/amu, E_{max} = 3.8E+04 eV/amu
(Ly_{β}) E_{min} = 8.3E-01 eV/amu, E_{max} = 3.8E+04 eV/amu
(Ly_{γ}) E_{min} = 1.3E+03 eV/amu, E_{max} = 3.8E+04 eV/amu

Chebyshev Fitting Parameters for Cross Sections

	A0	A1	A2	A3	A4	A5	A6	A7	A8
(Ly_{α})	-68.3492	.342390	-1.29878	-.802770	-.377264	.0373709	-.0758118	-.0906659	.182862
(Ly_{β})	-69.8158	-.378103	-1.48217	-1.03893	-.380773	.0694862	-.330382	-.0604033	.195831
(Ly_{γ})	-72.2366	-1.16996	-1.25436	-.404927	-.0551144	.0691596	.0107803	-.0171457	-.0103497

The fit represents the (Ly_{α}) cross section with an rms deviation of 3.3%.
The maximum deviation is 6.3% at 2.0E+00 eV/amu.
The fit represents the (Ly_{β}) cross section with an rms deviation of 4.6%.
The maximum deviation is 7.3% at 1.4E+01 eV/amu.
The fit represents the (Ly_{γ}) cross section with an rms deviation of 1.4%.
The maximum deviation is 2.7% at 1.5E+04 eV/amu.

See appendix for Chebyshev fit details.

$$He^{2+} + Li \rightarrow He^{+}(Ly_{\alpha}, Ly_{\beta}, Ly_{\gamma}) + Li^{+}$$

Cross Section vs. Energy



Line Emission Cross Sections of Electron Capture Collisions for

$$He^{2+} + Li \rightarrow He^{+}(H_\alpha, H_\beta, H_\gamma, H_\delta) + Li^{+}$$

(H_α)		(H_β)		(H_γ)		(H_δ)	
Energy eV/amu	Cross Section cm^2	Energy eV/amu	Cross Section cm^2	Energy eV/amu	Cross Section cm^2	Energy eV/amu	Cross Section cm^2
3.9E+03	9.19E-15	3.8E+03	7.87E-16	5.0E+03	2.26E-16	5.0E+03	1.19E-16
5.1E+03	8.03E-15	5.1E+03	1.05E-15	6.0E+03	3.19E-16	6.0E+03	1.24E-16
6.1E+03	7.23E-15	6.0E+03	1.10E-15	7.1E+03	3.60E-16	8.1E+03	1.32E-16
7.1E+03	6.55E-15	7.1E+03	1.11E-15	8.0E+03	3.66E-16	1.0E+04	1.38E-16
8.1E+03	5.73E-15	1.0E+04	9.79E-16	1.0E+04	3.41E-16	1.5E+04	1.29E-16
1.0E+04	4.07E-15	1.5E+04	5.20E-16	1.5E+04	2.33E-16	2.0E+04	8.23E-17
1.5E+04	1.49E-15	2.0E+04	2.71E-16	2.0E+04	1.46E-16	2.5E+04	3.58E-17
2.0E+04	6.02E-16	3.0E+04	9.35E-17	2.5E+04	7.35E-17	3.0E+04	2.30E-17
2.5E+04	2.60E-16	3.8E+04	5.04E-17	3.0E+04	4.61E-17	3.8E+04	1.59E-17
3.0E+04	1.03E-16			3.8E+04	2.99E-17		
3.7E+04	4.36E-17						

References: 240, 243, 245, 262

Accuracy: 60%

Notes: (1) The emission cross section is defined as the photon emission from state nl with no corrections for cascading i.e. Balmer-α emission cross section is the emission of photons in the 3 ->2 transition at λ =164 nm which does not include any other transitions.

(2) The probable accuracy of the data is estimated at 60%.

For Chebyshev fits of the above cross sections it is necessary to use the following parameters.

(H_α) E_{min} = 3.9E+03 eV/amu, E_{max} = 3.7E+04 eV/amu
(H_β) E_{min} = 3.8E+03 eV/amu, E_{max} = 3.8E+04 eV/amu
(H_γ) E_{min} = 5.0E+03 eV/amu, E_{max} = 3.8E+04 eV/amu
(H_δ) E_{min} = 5.0E+03 eV/amu, E_{max} = 3.8E+04 eV/amu

Chebyshev Fitting Parameters for Cross Sections

	A0	A1	A2	A3	A4	A5	A6	A7	A8
(H_α)	-68.6366	-2.68265	-.765389	-.0308549	.0354272	.00127287	.0359553	.0365120	.0161142
(H_β)	-71.0392	-1.46157	-.677621	.0638841	.0375648	.00465953	-.00918230	.0189071	.0164169
(H_γ)	-72.9366	-1.14183	-.554068	.105201	.0358399	.0591287	-.00855619	-.0336544	-.0423521
(H_δ)	-74.4125	-.995923	-.511481	-.0119520	.159667	.0670561	-.0432569	-.0655825	-.0724875

The fit represents the (H_α) cross section with an rms deviation of 0.6%.
The maximum deviation is 1.0% at 2.0E+04 eV/amu.
The fit represents the (H_β) cross section with an rms deviation of 0.0%.
The maximum deviation is 0.0%.
The fit represents the (H_γ) cross section with an rms deviation of 0.5%.
The maximum deviation is 1.1% at 8.0E+03 eV/amu.
The fit represents the (H_δ) cross section with an rms deviation of 0.0%.
The maximum deviation is 0.0%.

See appendix for Chebyshev fit details.

$$He^{2+} + Li \rightarrow He^{+}(H_{\alpha}, H_{\beta}, H_{\gamma}, H_{\delta}) + Li^{+}$$



C. Cross Sections for Excitation and Spectral Line Emission

(* denotes rate coefficient data also)

	Page
H + H -> H(2s)	C-2
-> H(2p)	C-2
-> H(3s)	C-4
-> H(3p)	C-4
-> H(3d)	C-4
-> H(Balmer-α)	C-4
$H + H_2$ -> H(2s) [projectile]	C-6
-> H(2p) [projectile]	C-6
-> (2s) [target]	C-6
-> (2p) [target]	C-6
-> (Lyman-α)	C-8
-> H(3s)	C-10
-> H(3p+3d)	C-10
-> H(Balmer-α)	C-12
-> H(4s)	C-14
-> H(4p+4d)	C-14
-> H(Balmer-β)	C-14
$H(2s) + H_2$ -> H(Lyman-α)	C-16
-> H(3s)	C-18
-> H(Balmer-α)	C-18
H(2s) + He -> H(Lyman-α)	C-20
-> H(3s)	C-22
-> H(Balmer-α)	C-22
H + He -> H(Lyman-α)	C-24*
-> H(2s)	C-30
-> H(2p)	C-30
-> H(3s)	C-32
-> H(3p)	C-32
-> H(3d)	C-32
-> H(3p+3d)	C-32
-> H(Balmer-α)	C-34
-> H(Balmer-β)	C-34
-> H(4s)	C-36
-> H(4p+4d)	C-36
-> He(3^1P)	C-38
-> He(3^3P)	C-38
-> He(3^3D)	C-38
-> He(4^1S)	C-40
-> He(4^1P)	C-40
-> He(4^1D)	C-40
-> He(4^3S)	C-42
-> He(4^3P)	C-42
-> He(4^3D)	C-42

C. (Cont'd)

$H^+ + H$ -> H(2s) C-44
-> H(2p) C-44*
-> H(n=2) C-48
-> H(n=3) C-48
-> H(n=4) C-48

$H^+ + H_2$ -> $H^+ + H(2s) + H$ C-50
-> $H^+ + H(2p) + H$ C-50
-> $H^+ + H(3p) + H$ C-52
-> H(Lyman-α) C-54
-> H(Balmer-α) C-56
-> H(Balmer-β) C-56
-> H(Balmer-γ) C-56
-> H_2(Lyman-Band) C-58
-> H_2(Lyman-Band) C-58

$H^+ + He$ -> He(2^1S) C-60*
-> He(2^1P) C-60*
-> He($2^1S + 2^1P$) C-60
-> He(3^1S) C-66
-> He(3^1P) C-66
-> He(3^1D) C-66
-> He(4^1S) C-68
-> He(4^1P) C-68
-> He(4^1D) C-68
-> v.u.v.(20-140 nm) C-70

$He^+ + H$ -> H(n=2) C-72
-> H(2s) C-74
-> H(2p) C-74*

$He^+ + H_2$ -> H(Lyman-α) C-78
-> H(Lyman-β) C-78
-> H(Balmer-α) C-80
-> H(Balmer-β) C-80
-> H(2s) C-82
-> H(2p) C-82
-> H(3s) C-84
-> H(3p) C-84
-> He^+(2s) C-86

$He^+ + He$ -> He(2^1S) [target] C-88
-> He(2^1P) [target] C-88
-> He(2^3S) [target] C-90
-> He(2^3P) [target] C-90
-> He(3^1S) [target] C-92
-> He(3^1P) [target] C-92
-> He(3^1D) [target] C-92
-> He(3^3S) [target] C-94
-> He(3^3P) [target] C-94
-> He(3^3D) [target] C-94

C. (Cont'd)

He^+ + He -> He(4^1S) [target] C-96
-> He(4^1P) [target] C-96
-> He(4^1D) [target] C-96
-> He(4^3S) [target] C-98
-> He(4^3P) [target] C-98
-> He(4^3D) [target] C-98
-> He^+(n=2) [projectile] C-100
-> He^+(n=3) [projectile] C-102

He + H_2 -> H(Balmer-α) C-104
-> H(Balmer-β) C-104

He + He -> He(2^1S) C-106
-> He(2^1P) C-106
-> He(3^1P) C-108
-> He(3^1D) C-108
-> He(3^3P) C-108
-> He(3^3D) C-108
-> He(4^1S) C-110
-> He(4^1P) C-110
-> He(4^3S) C-112
-> He(4^3P) C-112
-> He($2s^2$ ^1S) C-114
-> He($2s2p^3P$) C-114
-> He($2p^2$ ^1D) C-114
-> He($2s2p^1P$) C-114
-> He^-($1s2s^2$ ^2S) C-114

He^{2+} + H -> H(2s) C-116
-> H(2p) C-116

He(2^3S) + He(1^1S) -> He(3^1P) + He C-118
-> He(3^3P) + He C-120
-> He(4^3S) + He C-122
-> He(4^3D) + He C-122
-> He + He(4^3S) C-124
-> He + He(4^3D) C-124

H^+ + Li(2s) -> Li(2p) C-126
-> Li(670.8 nm) C-126

He^+ + Li(2s) -> Li(670.8 nm) C-128

He^{2+} + Li(2s) -> Li(2p) C-130
-> Li(670.8 nm) C-130

Excitation Cross Sections for

H + H -> H(2s,2p) + H(Σ)

H(2s)		H(2p)	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
2.0E+03	8.03E-18	4.0E+03	1.62E-17
3.0E+03	9.80E-18	5.0E+03	2.29E-17
4.0E+03	1.15E-17	6.0E+03	2.44E-17
5.0E+03	1.29E-17	7.0E+03	2.47E-17
6.0E+03	1.43E-17	8.0E+03	2.41E-17
7.0E+03	1.51E-17	9.0E+03	2.35E-17
8.0E+03	1.56E-17	1.0E+04	2.28E-17
9.0E+03	1.59E-17	1.5E+04	1.92E-17
1.0E+04	1.60E-17	2.0E+04	1.58E-17
1.5E+04	1.37E-17	3.0E+04	1.17E-17
2.0E+04	1.04E-17		
3.0E+04	6.64E-18		
4.0E+04	4.65E-18		
5.0E+04	3.48E-18		
6.0E+04	2.74E-18		
7.0E+04	2.28E-18		
8.0E+04	2.02E-18		
9.0E+04	1.84E-18		

References: 15, 592, 593, 594, 595, 596, 597, 598

Accuracy: Unknown

Notes: (1) The symbol H(Σ) denotes that the target atom may be left in any [excited] state after the collision. (2) All theoretical treatments of the H + H excitation cross sections are in serious disagreement with experiment and with each other, both in absolute magnitude and energy dependence. For various theoretical treatments see Flannery (Ref. 594 - four state impact parameter method), McLaughlin and Bell (Ref. 595 - close coupling exchange), Shingal et al. (Ref. 596 - four state semiclassical impact parameter), and Khurana et al. (Ref. 597 - 2nd Born approximation and also distorted wave Born).

Chebyshev Fitting Parameters for Cross Sections

H(2s) E_{min} = 2.0E+03, E_{max} = 9.0E+04
H(2p) E_{min} = 4.0E+03, E_{max} = 3.0E+04

	A0	A1	A2	A3	A4	A5	A06
H(2s)	-79.1098	-.784062	-.651723	.0462865	.114234	-.00487654	-.0114391
H(2p)	-77.1082	-.256046	-.232311	.0701782	-.0315673	.0229762	.00612383

The fit represents the H(2s) cross section with an rms deviation of 1.2%. The maximum deviation is 2.8% at 2.0E+04 eV/amu.

The fit represents the H(2p) cross section with an rms deviation of 0.3%. The maximum deviation is 0.6% at 6.0E+03 eV/amu.

See appendix for Chebyshev fit details.

H + H -> H(2s,2p)


Cross Section vs. Energy
10^{-16}
10^{-17}
10^{-18}
Cross Section (cm^2)
10^3
10^4
10^5
Energy (eV/amu)
Δ = H(2p)
× = H(2s)
Recommended Data
Chebyshev Fit

Excitation and Spectral Line Emission Cross Sections for

H + H -> H(3s,3p,3d) + H(Σ)
-> Balmer-α emission

H(3s)		H(3p)		H(3d)		Hα	
Energy (eV/amu)	σ (cm²)	Energy (eV/amu)	σ (cm²)	Energy (eV/amu)	σ (cm²)	Energy (eV/amu)	σ (cm²)
1.0E+03	1.68E-18	1.0E+03	7.24E-19	1.0E+03	2.53E-20	1.0E+03	1.87E-18
2.0E+03	6.73E-18	2.0E+03	5.76E-18	2.0E+03	3.22E-19	2.0E+03	8.27E-18
4.0E+03	1.22E-17	4.0E+03	1.45E-17	4.0E+03	1.14E-18	4.0E+03	1.45E-17
7.0E+03	8.59E-18	6.0E+03	1.63E-17	6.0E+03	1.39E-18	7.0E+03	1.20E-17
1.0E+04	6.47E-18	7.0E+03	1.59E-17	7.0E+03	1.38E-18	1.0E+04	9.34E-18
2.0E+04	3.48E-18	1.0E+04	1.34E-17	1.0E+04	1.18E-18	2.0E+04	5.05E-18
4.0E+04	1.76E-18	2.0E+04	7.42E-18	2.0E+04	6.78E-19	4.0E+04	2.68E-18
7.0E+04	1.07E-18	4.0E+04	4.27E-18	4.0E+04	4.11E-19	7.0E+04	1.71E-18
1.0E+05	7.69E-19	7.0E+04	3.11E-18	7.0E+04	2.99E-19	1.0E+05	1.32E-18
2.0E+05	4.08E-19	1.0E+05	2.55E-18	1.0E+05	2.44E-19	2.0E+05	7.60E-19
4.0E+05	2.11E-19	2.0E+05	1.61E-18	2.0E+05	1.50E-19	4.0E+05	3.97E-19
7.0E+05	1.20E-19	4.0E+05	9.29E-19	4.0E+05	8.23E-20	7.0E+05	2.32E-19
1.0E+06	8.72E-20	7.0E+05	5.57E-19	7.0E+05	4.91E-20	1.0E+06	1.65E-19
		1.0E+06	4.00E-19	1.0E+06	3.56E-20		

Reference: 598

Accuracy: Unknown

Notes: (1) We are not aware of any experimental data for the excitation of the n = 3 levels of H by H impact. (2) The data shown are obtained from McLaughlin and Bell (Ref. 598) using first order exchange theory. They also calculated these excitation cross sections using the first Born approximation. Below 10^5 eV/amu the Born calculations are as much as a factor of 5 below the data obtained from the exchange theory. These data are placed here to provide an insight as to the magnitude and energy dependence of the n = 3 excitation cross sections. (3) The Hα emission cross section is given by Hα = σ(3s) + 0.118 σ(3p) + σ(3d). (4) The symbol Σ denotes that the target atom may be left in any state.

Chebyshev Fitting Parameters for Cross Sections

H(3s)	E_{min}	=	1.0E+03	eV/amu,	E_{max}	=	1.0E+06 eV/amu
H(3p)	E_{min}	=	1.0E+03	eV/amu,	E_{max}	=	1.0E+06 eV/amu
H(3d)	E_{min}	=	1.0E+03	eV/amu,	E_{max}	=	1.0E+06 eV/amu
Hα	E_{min}	=	1.0E+03	eV/amu,	E_{max}	=	1.0E+06 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H(3s)	-82.5799	-2.05302	-.908967	.534101	-.249917	.0520636	.0418468
H(3p)	-81.2005	-.954175	-1.20328	.638574	-.328341	.0263311	.0719202
H(3d)	-86.3820	-.612848	-1.47849	.754596	-.362929	.0428931	.0876543
Hα	-81.7586	-1.76650	-.933877	.519497	.264797	.0406777	.0464266

The fit represents the H(3s) cross section with an rms deviation of 4.2%.
The maximum deviation is 6.2% at 2.0E+03 eV/amu.
The fit represents the H(3p) cross section with an rms deviation of 2.0%.
The maximum deviation is 2.9% at 2.0E+05 eV/amu.
The fit represents the H(3d) cross section with an rms deviation of 2.6%.
The maximum deviation is 4.6% at 2.0E+04 eV/amu.
The fit represents the Hα cross section with an rms deviation of 1.6%.
The maximum deviation is 2.2% at 2.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$H + H \rightarrow H(3s,3p,3d,H_\alpha)$



Projectile Excitation and Target Dissociative Excitation Cross Sections for

$$H + H_2 \rightarrow H(2s,2p) + \Sigma H_2$$
$$\rightarrow \Sigma H + H(2s,2p) + H$$

H(2s)[proj.]		H(2p)[proj.]		H(2s)[targ.]		H(2p)[targ.]	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
2.0E+03	5.56E-18	5.0E+03	4.13E-17	5.0E+03	2.78E-18	5.0E+03	6.90E-18
3.0E+03	6.69E-18	6.0E+03	3.53E-17	6.0E+03	3.26E-18	6.0E+03	8.60E-18
4.0E+03	7.62E-18	7.0E+03	3.11E-17	7.0E+03	3.57E-18	7.0E+03	9.97E-18
5.0E+03	8.47E-18	8.0E+03	2.79E-17	8.0E+03	3.75E-18	8.0E+03	1.12E-17
6.0E+03	9.19E-18	9.0E+03	2.61E-17	9.0E+03	3.81E-18	9.0E+03	1.22E-17
7.0E+03	9.90E-18	1.0E+04	2.49E-17	1.0E+04	3.86E-18	1.0E+04	1.32E-17
8.0E+03	1.05E-17	1.5E+04	2.40E-17	1.3E+04	3.88E-18	1.3E+04	1.39E-17
9.0E+03	1.11E-17	2.0E+04	2.08E-17	1.5E+04	3.83E-18	1.5E+04	1.37E-17
1.0E+04	1.16E-17	2.5E+04	1.78E-17	2.0E+04	3.68E-18	2.0E+04	1.16E-17
1.5E+04	1.32E-17			2.5E+04	3.53E-18	2.5E+04	9.03E-18
2.0E+04	1.20E-17						
2.5E+04	9.84E-18						

References: 124, 592

Accuracy: Unknown

Notes: (1) H is considered the projectile and H_2 the target. (2) The symbol Σ denotes all states in which the products are left. (3) The n=2->1 transition is at wavelength 121.57nm.

Chebyshev Fitting Parameters for Cross Sections

H(2s)	proj.	E_{min}	=	2.0E+03	eV/amu,	E_{max}	=	2.5E+04 eV/amu
H(2p)	proj.	E_{min}	=	5.0E+03	eV/amu,	E_{max}	=	2.5E+04 eV/amu
H(2s)	targ.	E_{min}	=	5.0E+03	eV/amu,	E_{max}	=	2.5E+04 eV/amu
H(2p)	targ.	E_{min}	=	5.0E+03	eV/amu,	E_{max}	=	2.5E+04 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H(2s) proj.	-78.5298	.381193	-.153241	-.0881960	-.0354361	-.00839689	.00756242
H(2p) proj.	-76.3640	-.378165	.0409330	-.0624299	-.0180012	.0197844	.0125324
H(2s) targ.	-80.3831	.0903819	-.109415	.0282363	-.00676568	.000903034	.00303522
H(2p) targ.	-78.2178	.158165	-.269341	-.0251991	.00457227	.00145158	-.00685275

The fit represents the H(2s) proj. cross section with an rms deviation of 0.5%. The maximum deviation is 0.8% at 1.0E+04 eV/amu.

The fit represents the H(2p) proj. cross section with an rms deviation of 0.2%. The maximum deviation is 0.3% at 8.0E+03 eV/amu.

The fit represents the H(2s) targ. cross section with an rms deviation of 0.2%. The maximum deviation is 0.3% at 9.0E+03 eV/amu.

The fit represents the H(2p) targ. cross section with an rms deviation of 0.4%. The maximum deviation is 0.7% at 1.3E+04 eV/amu.

See appendix for Chebyshev fit details.

$$H + H_2 \rightarrow H(2s,2p) + \Sigma H_2$$
$$H + H_2 \rightarrow \Sigma H + H(2s,2p) + H$$



Total Spectral Line Emission Cross Sections for

$H + H_2 \rightarrow H(\text{Lyman-}\alpha)$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.5E+03	5.38E+07	7.17E-17
2.0E+03	6.21E+07	6.67E-17
3.0E+03	7.61E+07	5.98E-17
4.0E+03	8.79E+07	5.38E-17
5.0E+03	9.82E+07	4.91E-17
6.0E+03	1.08E+08	4.50E-17
7.0E+03	1.16E+08	4.22E-17
8.0E+03	1.24E+08	4.04E-17
9.0E+03	1.32E+08	3.92E-17
1.0E+04	1.39E+08	3.80E-17
1.5E+04	1.70E+08	3.48E-17
2.0E+04	1.96E+08	3.20E-17
2.5E+04	2.20E+08	2.97E-17
3.0E+04	2.41E+08	2.81E-17
4.0E+04	2.78E+08	2.52E-17
5.0E+04	3.11E+08	2.28E-17
6.0E+04	3.40E+08	2.15E-17
7.0E+04	3.68E+08	2.06E-17
8.0E+04	3.93E+08	1.98E-17
9.0E+04	4.17E+08	1.92E-17
1.0E+05	4.39E+08	1.88E-17
1.3E+05	5.01E+08	1.77E-17

References: 124, 194, 333

Accuracy: Unknown

Note: The total spectral line emission includes radiation from excitation of the H projectile and the dissociative excitation of the H_2 target.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.5E+03 eV/amu, E_{max} = 1.3E+05 eV/amu

A0	A1	A2	A3	A4	A5	A6
-75.7615	-.720784	.0118290	.00540110	-.00257496	.0252003	-.00232715

The fit represents the above cross section with an rms deviation of 1.2% The maximum deviation is 2.4% at 2.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$H + H_2 \rightarrow \Sigma Ly_\alpha$



Excitation Cross Sections for

$$H + H_2 \rightarrow H(3s) + H_2$$
$$\rightarrow H(3p+3d) + H_2$$

H(3s)		H(3p+3d)	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section* (cm^2)
2.0E+02	1.84E-18	2.0E+02	5.72E-18
4.0E+02	4.61E-18	4.0E+02	1.01E-17
6.4E+02	5.66E-18	7.0E+02	1.25E-17
7.0E+02	5.61E-18	1.0E+03	1.35E-17
1.0E+03	4.95E-18	1.5E+03	1.28E-17
1.5E+03	3.53E-18	2.0E+03	1.17E-17
2.0E+03	2.96E-18	3.0E+03	8.66E-18
4.0E+03	2.43E-18	4.0E+03	6.34E-18
5.0E+03	2.36E-18	7.0E+03	3.91E-18
7.0E+03	2.46E-18	1.0E+04	3.64E-18
1.0E+04	3.21E-18	1.5E+04	3.13E-18
1.5E+04	4.07E-18	2.0E+04	2.13E-18
2.0E+04	4.41E-18	4.0E+04	1.29E-18
2.3E+04	4.50E-18	7.0E+04	1.17E-18
2.5E+04	4.46E-18	1.0E+05	1.16E-18
4.0E+04	3.59E-18		
7.0E+04	2.86E-18	* see notes (3) and (4)	
1.0E+05	2.73E-18		

References: 133, 500, 601

Accuracy: Unknown

Notes: (1) Because of its much longer lifetime, the 3s excitation is resolvable in the Balmer-α emission. (2) The 3s cross section represents that for excitation of the H projectile only. (3) The cross section labelled H(3p + 3d) is determined by subtracting σ(3s) from the total Balmer-α emission cross section, and therefore represents σ(3d) + 0.12 σ(3p) because of branching ratios. (4) For energies above 2 keV/amu, the H(3p + 3d) cross section represents excitation of the projectile only. At lower energies, contributions from dissociative excitation of H_2 target molecules are included. Comparison of the data suggests that these contributions are small.

Chebyshev Fitting Parameters for Cross Sections

H(3s) E_{min} = 2.0E+02 eV/amu, E_{max} = 1.0E+05 eV/amu
H(3p+3d) E_{min} = 2.0E+02 eV/amu, E_{max} = 1.0E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H(3s)	-80.5534	-.0150679	-.145036	.109627	-.341180	.113781	.131159
H(3p+3d)	-80.1203	-1.13376	-.496856	.359704	.00520846	-.0168457	.0653151

The fit represents the H(3s) cross section with an rms deviation of 3.2%. The maximum deviation is 7.0% at 4.0E+04 eV/amu.

The fit represents the H(3p+3d) cross section with an rms deviation of 7.1%. The maximum deviation is 14.5% at 7.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$H + H_2 \rightarrow H(3s) + H_2$
$H + H_2 \rightarrow H(3p+3d) + H_2$



Total Spectral Line Emission Cross Sections for

$H + H_2 \rightarrow$ (Balmer-α)

Total ($H + H_2$)		H_2 (targ.)		H (proj.)	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
5.0E+01	1.98E-19	2.0E+03	1.90E-18	2.0E+03	1.58E-17
7.0E+01	6.35E-19	3.0E+03	1.87E-18	3.0E+03	1.14E-17
1.0E+02	1.67E-18	4.0E+03	1.90E-18	4.0E+03	9.06E-18
1.5E+02	4.93E-18	5.0E+03	1.95E-18	5.0E+03	7.61E-18
2.0E+02	7.69E-18	6.0E+03	2.02E-18	6.0E+03	6.83E-18
4.0E+02	1.47E-17	7.0E+03	2.11E-18	7.0E+03	6.54E-18
7.0E+02	1.80E-17	8.0E+03	2.24E-18	7.5E+03	6.52E-18
9.0E+02	1.84E-17	9.0E+03	2.37E-18	8.0E+03	6.62E-18
1.0E+03	1.82E-17	1.0E+04	2.54E-18	9.0E+03	6.88E-18
1.5E+03	1.75E-17	1.5E+04	2.94E-18	1.0E+04	7.04E-18
2.0E+03	1.56E-17	2.0E+04	3.15E-18	1.5E+04	7.11E-18
4.0E+03	1.10E-17	2.3E+04	3.18E-18	2.0E+04	6.69E-18
7.0E+03	8.76E-18	3.0E+04	2.89E-18	3.0E+04	5.39E-18
8.0E+03	8.69E-18	4.0E+04	2.30E-18	4.0E+04	4.73E-18
1.0E+04	9.54E-18	5.0E+04	1.93E-18	5.0E+04	4.43E-18
1.4E+04	1.01E-17	6.0E+04	1.63E-18	6.0E+04	4.25E-18
2.0E+04	9.84E-18	7.0E+04	1.41E-18	7.0E+04	4.20E-18
4.0E+04	7.14E-18	8.0E+04	1.24E-18	8.0E+04	4.18E-18
7.0E+04	5.26E-18	9.0E+04	1.11E-18	9.0E+04	4.17E-18
1.0E+05	4.61E-18	1.0E+05	1.01E-18	1.0E+05	4.17E-18

<u>References:</u> 133, 500, 601

<u>Accuracy:</u> Unknown

<u>Notes:</u> (1) The total Balmer-α emission cross section $H\alpha$ is equal to the sum of the emission from the projectile ($H_{p\alpha}$) and the target ($H_{T\alpha}$). (2) Balmer-α radiation is from the n=3->n=2 transition in H with λ = 656.28nm.

Chebyshev Fitting Parameters for Cross Sections

Total	E_{min}	=	5.0E+01	eV/amu,	E_{max}	=	1.0E+05 eV/amu
H_2	E_{min}	=	2.0E+03	eV/amu,	E_{max}	=	1.0E+05 eV/amu
H	E_{min}	=	2.0E+03	eV/amu,	E_{max}	=	1.0E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
Total	-79.7992	.953923	-1.39409	.663856	-.262806	-.0749405	.0516520
H_2	-81.5995	-.183064	-.361113	-.185120	.0559114	.0591802	-.0130234
H	-79.0077	-.627859	.128139	-.0950362	.0772477	.0565053	-.0596124

The fit represents the Total cross section with an rms deviation of 7.0%. The maximum deviation is 13.6% at 7.0E+03 eV/amu.

The fit represents the H_2 cross section with an rms deviation of 1.4%. The maximum deviation is 3.0% at 4.0E+04 eV/amu.

The fit represents the H cross section with an rms deviation of 1.9%. The maximum deviation is 3.4% at 3.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$H + H_2 \rightarrow H_\alpha$



Total Excitation and Spectral Line Emission Cross Sections for

$H + H_2 \rightarrow H(4s,4p+4d)$
$\rightarrow H(Balmer-\beta)$

H(4s)		H(4p+4d)		H(Balmer-β)	
Energy (eV/amu)	σ (cm²)	Energy (eV/amu)	σ (cm²)	Energy (eV/amu)	σ (cm²)
2.0E+02	2.69E-19	2.0E+02	9.25E-19	6.0E+01	3.13E-20
3.0E+02	5.20E-19	3.0E+02	1.50E-18	7.0E+01	5.98E-20
4.0E+02	8.30E-19	4.0E+02	1.91E-18	1.0E+02	2.19E-19
5.0E+02	1.01E-18	5.0E+02	2.20E-18	1.5E+02	7.38E-19
6.0E+02	1.11E-18	6.0E+02	2.40E-18	2.0E+02	1.24E-18
7.0E+02	1.12E-18	7.0E+02	2.54E-18	4.0E+02	2.67E-18
8.0E+02	1.10E-18	8.0E+02	2.66E-18	7.0E+02	3.66E-18
9.0E+02	1.06E-18	9.0E+02	2.76E-18	1.0E+03	3.90E-18
1.0E+03	1.02E-18	1.0E+03	2.86E-18	1.5E+03	3.95E-18
1.5E+03	8.58E-19	1.3E+03	2.99E-18	2.0E+03	3.69E-18
2.0E+03	7.48E-19	1.5E+03	2.98E-18	2.5E+03	3.37E-18
2.5E+03	6.67E-19	2.0E+03	2.84E-18		
		2.5E+03	2.64E-18		

References: 600, 601

Accuracy: Unknown

Notes: (1) The cross sections include excitation of the projectile H, dissociative excitation of the target H_2, and H Balmer-β line emission from both projectile and target. (2) The Balmer-β radiation has a wavelength of 486.13nm, and corresponds to the n=4->n=2 transition in H.

Chebyshev Fitting Parameters for cross Sections

H(4s)	E_{min}	=	2.0E+02	eV/amu,	E_{max}	=	2.5E+03 eV/amu
H(4p+4d)	E_{min}	=	2.0E+02	eV/amu,	E_{max}	=	2.5E+03 eV/amu
H(Balmer-β)	E_{min}	=	6.0E+01	eV/amu,	E_{max}	=	2.5E+03 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H(4s)	-83.7360	.403599	-.495244	.0926298	.0476553	-.0421657	.00996875
H(4p+4d)	-81.4782	.513746	-.253426	.0120434	-.0154708	-.00146007	.00795248
H(Balmer-β)	-82.9678	2.14325	-1.06498	.222304	-.0396703	-.0277369	.0217573

The fit represents the H(4s) cross section with an rms deviation of 0.4%. The maximum deviation is 1.1% at 5.0E+02 eV/amu.

The fit represents the H(4p+4d) cross section with an rms deviation of 0.3%. The maximum deviation is 0.4% at 4.0E+02 eV/amu.

The fit represents the H(Balmer-β) cross section with an rms deviation of 1.6%. The maximum deviation is 2.5% at 1.0E+02 eV/amu.

See appendix for Chebyshev fit details.

$H + H_2 \rightarrow H(4s, 4p+4d)$
$H + H_2 \rightarrow H_\beta$



Total Spectral Line Emission Cross Sections for

$H(2s) + H_2 \rightarrow H(\text{Lyman-}\alpha)$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
7.0E+03	1.16E+08	1.46E-16
8.0E+03	1.24E+08	1.49E-16
9.0E+03	1.32E+08	1.50E-16
1.0E+04	1.39E+08	1.49E-16
1.1E+04	1.46E+08	1.48E-16
1.2E+04	1.52E+08	1.46E-16
1.3E+04	1.58E+08	1.43E-16
1.4E+04	1.64E+08	1.41E-16
1.5E+04	1.70E+08	1.38E-16
1.6E+04	1.76E+08	1.34E-16
1.7E+04	1.81E+08	1.31E-16
1.8E+04	1.86E+08	1.27E-16
1.9E+04	1.91E+08	1.23E-16
2.0E+04	1.96E+08	1.20E-16
2.1E+04	2.01E+08	1.16E-16
2.2E+04	2.06E+08	1.13E-16
2.3E+04	2.11E+08	1.10E-16
2.4E+04	2.15E+08	1.07E-16
2.5E+04	2.20E+08	1.04E-16
2.6E+04	2.24E+08	1.02E-16

References: 333, 692

Accuracy: 60%

Notes: (1) These data include emission from excitation of the projectile H and from the dissociative excitation of H_2. (2) Lyman-α emission occurs at a wavelength of 121.57nm.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 7.0E+03 eV/amu, E_{max} = 2.6E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-73.1370	-.183791	-.0767839	.00130611	.00262379	.00284123	-8.70436E-05

The fit represents the above cross section with an rms deviation of 0.2%. The maximum deviation is 0.4% at 1.3E+04 eV/amu.

See appendix for Chebyshev fit details.

$H(2s) + H_2 \rightarrow Ly_\alpha$



Excitation and Spectral Line Emission Cross Sections for

$$H(2s) + H_2 \rightarrow H(3s)$$
$$\rightarrow H(\text{Balmer-}\alpha)$$

H(Balmer-α)		H(3s)	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
5.0E+03	2.27E-17	5.0E+03	1.58E-17
6.0E+03	2.29E-17	6.0E+03	1.60E-17
8.0E+03	2.31E-17	8.0E+03	1.64E-17
1.0E+04	2.32E-17	9.0E+03	1.65E-17
1.2E+04	2.33E-17	1.0E+04	1.64E-17
1.4E+04	2.31E-17	1.2E+04	1.60E-17
1.6E+04	2.22E-17	1.4E+04	1.53E-17
1.8E+04	2.06E-17	1.6E+04	1.45E-17
2.0E+04	1.88E-17	1.8E+04	1.36E-17
2.2E+04	1.75E-17	2.0E+04	1.28E-17
2.4E+04	1.66E-17	2.2E+04	1.21E-17
2.6E+04	1.58E-17	2.4E+04	1.15E-17
		2.6E+04	1.10E-17

Reference: 691

Accuracy: 50%

Notes: (1) The Balmer-α cross sections are determined by measuring the emission at λ = 656.28nm from the n = 3 states of H and are related to the level excitation cross sections as follows: $\sigma(H\alpha) = \sigma(3s) + 0.118\ \sigma(3p) + \sigma(3d)$. The factor 0.118 arises from the branching ratio for decay of the 3p state. (2) The $\sigma(3s)$ have been corrected for cascading.

Chebyshev Fitting Parameters for Cross Sections

H(Balmer-α) E_{min} = 5.0E+03 eV/amu, E_{max} = 2.6E+04 eV/amu
H(3s) E_{min} = 5.0E+03 eV/amu, E_{max} = 2.6E+04 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H(Balmer-α)	-76.8150	-.162721	-.109854	-.0325036	.0085463	.0165529	.00525437
H(3s)	-77.5541	-.175436	-.101810	-.00744771	.0115273	.00202386	-7.45889E-05

The fit represents the H(Balmer-α) cross section with an rms deviation of 0.5%. The maximum deviation is 1.0% at 2.0E+04 eV/amu.

The fit represents the H(3s) cross section with an rms deviation of 0.1%. The maximum deviation is 0.1% at 1.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$H(2s) + H_2 \rightarrow H(3s)$
$H(2s) + H_2 \rightarrow H_\alpha$



Spectral Line Emission Cross Sections for

H(2s) + He -> H(Lyman-α)

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.0E+04	1.39E+08	1.86E-17
1.1E+04	1.46E+08	1.75E-17
1.2E+04	1.52E+08	1.65E-17
1.3E+04	1.58E+08	1.56E-17
1.4E+04	1.64E+08	1.48E-17
1.5E+04	1.70E+08	1.46E-17
1.6E+04	1.76E+08	1.43E-17
1.7E+04	1.81E+08	1.41E-17
1.8E+04	1.86E+08	1.37E-17
1.9E+04	1.91E+08	1.32E-17
2.0E+04	1.96E+08	1.28E-17
2.1E+04	2.01E+08	1.24E-17
2.2E+04	2.06E+08	1.22E-17
2.3E+04	2.11E+08	1.20E-17
2.4E+04	2.15E+08	1.19E-17
2.5E+04	2.20E+08	1.19E-17
2.6E+04	2.24E+08	1.18E-17

Reference: 333

Accuracy: 80%

Note: The H Lyman-α radiation is at a wavelength of 121.57nm.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.0E+04 eV/amu, E_{max} = 2.6E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-77.5542	-.232829	.0249886	-.00284785	.0873114	.0106037	-.00679600

The fit represents the above cross section with an rms deviation of 0.5%. The maximum deviation is 1.2% at 1.4E+04 eV/amu.

See appendix for Chebyshev fit details.

$H(2s) + He \rightarrow H(Ly_\alpha)$



Excitation and Spectral Line Emission Cross Sections For

H(2s) + He -> H(3s)
-> H(Balmer-α)

H(3s)		H(Balmer-α)	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
5.0E+03	7.05E-18	5.0E+03	1.15E-17
6.0E+03	6.62E-18	6.0E+03	1.11E-17
8.0E+03	5.95E-18	8.0E+03	1.04E-17
1.0E+04	5.53E-18	1.0E+04	9.86E-18
1.2E+04	5.21E-18	1.2E+04	9.35E-18
1.4E+04	5.05E-18	1.4E+04	8.97E-18
1.6E+04	4.95E-18	1.6E+04	8.66E-18
1.8E+04	4.83E-18	1.8E+04	8.37E-18
2.0E+04	4.67E-18	2.0E+04	8.13E-18
2.2E+04	4.44E-18	2.2E+04	7.87E-18
2.4E+04	4.23E-18	2.4E+04	7.64E-18
2.6E+04	4.07E-18	2.6E+04	7.42E-18

Reference: 691

Accuracy: 50%

Notes: (1) The Balmer-α emission cross section is given by $\sigma(3s) + 0.118\ \sigma(3p) + \sigma(3d)$. The factor 0.118 is to correct for the branching ratio of the 3p state. (2) The $\sigma(3s)$ has been corrected for cascading. (3) The Balmer-α radiation is at a wavelength of 656.28nm.

Chebyshev fitting Parameters for Cross Sections

H(3s) E_{min} = 5.0E+03 eV/amu, E_{max} = 2.6E+04 eV/amu
H(Balmer-α) E_{min} = 5.0E+03 eV/amu, E_{max} = 2.6E+04 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H(3s)	-79.5230	-.258972	7.15363E-04	-.0165922	-.0141266	-2.93267E-04	.00596856
H(Balmer-α)	-78.4151	-.218179	-.0140659	2.10927E-05	-.00210701	-.00118313	3.35892E-04

The fit represents the H(3s) cross section with an rms deviation of 0.3%. The maximum deviation is 0.5% at 8.0E+03 eV/amu.

The fit represents the H(Balmer-α) cross section with an rms deviation of 0.1%. The maximum deviation is 0.1% at 8.0E+03 eV/amu.

See appendix for Chebyshev fit details.

H(2s) + He -> H_α
H(2s) + He -> H(3s)



Spectral Line Emission Cross Sections for

H + He -> H(Lyman-α)

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
3.0E+01	7.61E+06	1.01E-17
4.0E+01	8.79E+06	2.41E-17
7.0E+01	1.16E+07	4.35E-17
1.0E+02	1.39E+07	5.23E-17
2.0E+02	1.96E+07	6.65E-17
4.0E+02	2.78E+07	6.96E-17
7.0E+02	3.68E+07	6.90E-17
1.0E+03	4.39E+07	6.58E-17
2.0E+03	6.21E+07	5.61E-17
7.0E+03	1.16E+08	2.63E-17
1.0E+04	1.39E+08	1.95E-17
2.0E+04	1.96E+08	1.00E-17
4.0E+04	2.78E+08	4.40E-18
7.0E+04	3.68E+08	1.90E-18
9.0E+04	4.17E+08	1.26E-18

References: 46, 639, 643, 644, 645, 651

Accuracy: 80%

Note: The Lyman-α emission at 121.6nm includes small contributions due to cascade from the n = 3 and higher levels.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 3.0E+01 eV/amu, E_{max} = 9.0E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-77.2423	-1.27626	-1.33104	.105602	-.131109	.0901017	-.0744630

The fit represents the above cross sections with an rms deviation of 3.8%. The maximum deviation is 5.8% at 3.0E+01 eV/amu.

See appendix for Chebyshev fit details.

H + He -> H(Lyman$_\alpha$)



Spectral Line Emission Rate Coefficients for

H + He -> H(Lyman-α)

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H Temp. (eV)	Equal Temp.	He Temp. (eV) 10.	100.	500.	1000.	5000.	10000.	20000.
1.0E+01	4.37E-11*	4.37E-11*	3.03E-10	1.10E-09	1.64E-09	2.86E-09	3.02E-09	2.85E-09
2.0E+01	1.86E-10*	1.56E-10*	4.13E-10	1.15E-09	1.67E-09	2.87E-09	3.02E-09	2.85E-09
4.0E+01	4.64E-10	3.86E-10	6.06E-10	1.26E-09	1.74E-09	2.87E-09	3.02E-09	2.85E-09
6.0E+01	6.92E-10	5.84E-10	7.71E-10	1.35E-09	1.80E-09	2.88E-09	3.02E-09	2.85E-09
8.0E+01	8.80E-10	7.52E-10	9.14E-10	1.44E-09	1.85E-09	2.89E-09	3.02E-09	2.84E-09
1.0E+02	1.04E-09	8.97E-10	1.04E-09	1.51E-09	1.91E-09	2.89E-09	3.02E-09	2.84E-09
2.0E+02	1.61E-09	1.43E-09	1.51E-09	1.84E-09	2.13E-09	2.92E-09	3.01E-09	2.83E-09
4.0E+02	2.22E-09	2.03E-09	2.08E-09	2.26E-09	2.44E-09	2.96E-09	3.01E-09	2.81E-09
6.0E+02	2.55E-09	2.38E-09	2.41E-09	2.53E-09	2.64E-09	2.99E-09	3.00E-09	2.80E-09
8.0E+02	2.74E-09	2.60E-09	2.62E-09	2.69E-09	2.77E-09	3.00E-09	2.99E-09	2.78E-09
1.0E+03	2.86E-09	2.74E-09	2.75E-09	2.81E-09	2.86E-09	3.01E-09	2.98E-09	2.76E-09
2.0E+03	3.02E-09	3.00E-09	3.00E-09	3.01E-09	3.01E-09	2.99E-09	2.90E-09	2.67E-09
4.0E+03	2.85E-09	2.94E-09	2.94E-09	2.93E-09	2.92E-09	2.83E-09	2.71E-09	2.50E-09
6.0E+03	2.62E-09	2.76E-09	2.76E-09	2.75E-09	2.74E-09	2.64E-09	2.54E-09	2.34E-09
8.0E+03	2.42E-09	2.58E-09	2.58E-09	2.57E-09	2.56E-09	2.47E-09	2.38E-09	2.20E-09
1.0E+04	2.24E-09	2.42E-09	2.41E-09	2.41E-09	2.40E-09	2.32E-09	2.24E-09	2.08E-09
1.5E+04	1.88E-09	2.08E-09	2.08E-09	2.07E-09	2.07E-09	2.01E-09	1.94E-09	1.82E-09
2.0E+04	1.61E-09	1.82E-09	1.82E-09	1.81E-09	1.81E-09	1.76E-09	1.71E-09	1.61E-09

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+01 eV, E_{max} = 2.0E+04 eV

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
10.	-41.8408	1.73612	-1.00696	.107279	-.0576945	.0226348	-.00478217
100.	-41.0379	1.03224	-.535100	-.134058	.0352264	-.00215004	-.00222457
500.	-40.2599	.410395	-.247167	-.176225	.00337909	.0177532	-.00518248
1000.	-39.9551	.185465	-.185562	-.151177	-.0159001	.0155346	-1.77904E-04
5000.	-39.4938	-.156752	-.142409	-.0847522	-.0277793	-.00101284	.00283315
10000.	-39.4782	-.210389	-.141402	-.0704827	-.0240167	-.00386296	8.94493E-04
20000.	-39.6096	-.218997	-.135741	-.0627840	-.0213979	-.00486653	-4.40643E-04
Equal Temp.	-41.7630	1.61688	-1.06969	.147665	-.0861736	.0380393	-.0145355

See appendix for Chebyshev fit details.

H + He -> H(Lyman$_\alpha$)

Maxwellian – Maxwellian



Spectral Line Emission Rate Coefficients for

He + H -> He + H(Lyman-α)

Beam - Maxwellian Rate Coefficients (cm^3/s)

H Temp. (eV)	He Energy (eV/amu) 10000.	20000.	30000.	40000.	50000.	60000.	70000.
1.0E+00	2.68E-09	1.96E-09	1.49E-09	1.22E-09	9.78E-10	8.15E-10	6.98E-10
2.0E+00	2.68E-09	1.96E-09	1.49E-09	1.22E-09	9.78E-10	8.15E-10	6.98E-10
4.0E+00	2.68E-09	1.96E-09	1.49E-09	1.22E-09	9.78E-10	8.15E-10	6.98E-10
7.0E+00	2.68E-09	1.96E-09	1.49E-09	1.22E-09	9.78E-10	8.15E-10	6.97E-10
1.0E+01	2.68E-09	1.96E-09	1.49E-09	1.22E-09	9.78E-10	8.15E-10	6.97E-10
2.0E+01	2.68E-09	1.96E-09	1.49E-09	1.22E-09	9.78E-10	8.15E-10	6.97E-10
4.0E+01	2.68E-09	1.95E-09	1.49E-09	1.22E-09	9.78E-10	8.15E-10	6.97E-10
7.0E+01	2.67E-09	1.95E-09	1.49E-09	1.21E-09	9.78E-10	8.15E-10	6.96E-10
1.0E+02	2.67E-09	1.95E-09	1.49E-09	1.21E-09	9.79E-10	8.15E-10	6.96E-10
2.0E+02	2.66E-09	1.94E-09	1.49E-09	1.21E-09	9.80E-10	8.16E-10	6.95E-10
4.0E+02	2.64E-09	1.93E-09	1.49E-09	1.20E-09	9.81E-10	8.17E-10	6.90E-10
7.0E+02	2.61E-09	1.92E-09	1.49E-09	1.20E-09	9.81E-10	8.17E-10	6.74E-10
1.0E+03	2.58E-09	1.91E-09	1.48E-09	1.20E-09	9.80E-10	8.13E-10	6.58E-10
2.0E+03	2.49E-09	1.88E-09	1.47E-09	1.19E-09	9.73E-10	7.92E-10	6.18E-10*
4.0E+03	2.32E-09	1.81E-09	1.44E-09	1.16E-09	9.43E-10	7.53E-10*	5.81E-10*
7.0E+03	2.12E-09	1.70E-09	1.37E-09	1.11E-09	8.98E-10	7.15E-10*	5.59E-10*
1.0E+04	1.96E-09	1.59E-09	1.30E-09	1.06E-09	8.60E-10*	6.90E-10*	5.47E-10*
2.0E+04	1.53E-09	1.29E-09	1.08E-09*	9.04E-10*	7.54E-10*	6.26E-10*	5.18E-10*

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-39.6987	-.197802	-.134319	-.0723203	-.0304433	-.00974030	-.00251684
20000.	-40.2569	-.138429	-.0956246	-.0581643	-.0310565	-.0136698	-.00449523
30000.	-40.7530	-.0951522	-.0755129	-.0512811	-.0288118	-.0130297	-.00477600
40000.	-41.1570	-.0957055	-.0666225	-.0426726	-.0248045	-.0119141	-.00390059
50000.	-41.5800	-.0821883	-.0654040	-.0420679	-.0203976	-.00620683	2.45851E-06
60000.	-41.9647	-.0972904	-.0701222	-.0364437	-.0102495	.00215119	.00335803
70000.	-42.3333	-.139674	-.0746309	-.0176390	.00996598	.0106575	-8.59901E-05

See appendix for Chebyshev fit details.

H + He -> H(Lyman$_\alpha$)



Excitation Cross Sections for

H + He -> H(2s,2p)

H(2s) Energy (eV/amu)	H(2s) Cross Section (cm^2)	H(2p) Energy (eV/amu)	H(2p) Cross Section (cm^2)
6.0E+01	1.43E-18	2.0E+01	1.87E-18
7.0E+01	1.81E-18	4.0E+01	2.36E-17
1.0E+02	3.04E-18	7.0E+01	4.33E-17
2.0E+02	6.70E-18	1.0E+02	5.22E-17
4.0E+02	1.10E-17	2.0E+02	6.08E-17
7.0E+02	1.29E-17	4.0E+02	6.30E-17
8.6E+02	1.31E-17	7.0E+02	6.11E-17
1.0E+03	1.30E-17	1.0E+03	5.81E-17
2.0E+03	1.10E-17	2.0E+03	5.16E-17
4.0E+03	8.41E-18	4.0E+03	3.73E-17
7.0E+03	6.54E-18	7.0E+03	2.58E-17
1.0E+04	5.51E-18	1.0E+04	1.95E-17
2.0E+04	4.15E-18	2.0E+04	1.01E-17
4.0E+04	3.64E-18	4.0E+04	5.16E-18
7.0E+04	2.67E-18	7.0E+04	4.52E-18
1.0E+05	2.04E-18	1.0E+05	4.41E-18
1.4E+05	1.59E-18		

<u>References:</u> 46, 639, 640, 641, 642, 643, 644, 645, 646, 648, 650

<u>Accuracy:</u> 50%

<u>Notes:</u> (1) Excitation of an excited state is found by measuring the emission from that state and correcting for cascades into that state. (2) The 2s state cross sections are found from the electric-field-induced transition (2s -> 2p -> 1s). (3) If cascading is neglected the Lyman-α emission cross sections are equal to the 2p excitation cross sections.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

H(2s) E_{min} = 6.0E+01 eV/amu, E_{max} = 1.4E+05 eV/amu
H(2p) E_{min} = 2.0E+01 eV/amu, E_{max} = 1.0E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H(2s)	-80.0462	-.212388	-.933176	.333291	-.0922211	-.0813910	.00473762
H(2p)	-77.4567	-.447338	-1.40980	.577607	-.164863	.291318	-.0758830

The fit represents the H(2s) cross section with an rms deviation of 2.2%. The maximum deviation is 4.3% at 1.0E+05 eV/amu.

The fit represents the H(2p) cross section with an rms deviation of 3.8%. The maximum deviation is 6.9% at 4.0E+04 eV/amu.

See appendix for Chebyshev fit details.

H + He -> H(2s,2p)


Cross Section vs. Energy
Cross Section (cm^2)
10^{-16}
10^{-17}
10^{-18}
10^1
10^2
10^3
10^4
10^5
10^6
Energy (eV/amu)
Δ = H(2p)
× = H(2s)
Recommended Data
Chebyshev Fit

Excitation Cross Sections for

H + He -> H(3s,3p,3d)
-> H(3p+3d)

H(3s)		H(3p)		H(3d)		H(3p+3d)	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
1.5E+02	1.00E-18	5.0E+03	2.41E-18	1.0E+04	1.20E-18	1.5E+02	2.54E-18
2.0E+02	1.84E-18	5.5E+03	2.33E-18	1.3E+04	1.05E-18	2.0E+02	3.28E-18
4.0E+02	4.18E-18	6.0E+03	2.25E-18	1.5E+04	8.64E-19	3.0E+02	4.24E-18
6.0E+02	4.64E-18	7.0E+03	2.08E-18	2.0E+04	5.96E-19	4.0E+02	4.41E-18
7.0E+02	4.57E-18	8.0E+03	1.94E-18	2.5E+04	4.88E-19	5.0E+02	4.33E-18
1.0E+03	3.99E-18	9.0E+03	1.82E-18	3.0E+04	4.48E-19	6.0E+02	4.26E-18
2.0E+03	2.74E-18	1.5E+04	1.38E-18	3.5E+04	4.41E-19	7.0E+02	4.17E-18
4.0E+03	1.90E-18	2.0E+04	1.15E-18			8.0E+02	4.08E-18
7.0E+03	1.46E-18	2.5E+04	1.02E-18			9.0E+02	4.02E-18
1.0E+04	1.26E-18	3.0E+04	9.31E-19			1.0E+03	3.95E-18
2.0E+04	9.39E-19	3.5E+04	8.75E-19			1.5E+03	3.70E-18
3.5E+04	7.98E-19	4.0E+04	8.40E-19			2.0E+03	3.50E-18
						2.5E+03	3.36E-18

References: 500, 601, 645

Accuracy: See notes

Notes: (1) The excitation cross section is found by measuring the emission cross section from that state and correcting for cascade from measurements or using known transition probabilities. (2) Excitation cross sections are often inaccurate due to problems with the absolute calibration of the photon detector.

Chebyshev Fitting Parameters for Cross Sections

H(3s)	E_{min}	=	1.5E+02	eV/amu,	E_{max}	=	3.5E+04 eV/amu
H(3p)	E_{min}	=	5.0E+03	eV/amu,	E_{max}	=	4.0E+04 eV/amu
H(3d)	E_{min}	=	1.0E+04	eV/amu,	E_{max}	=	3.5E+04 eV/amu
H(3p+3d)	E_{min}	=	1.5E+02	eV/amu,	E_{max}	=	2.5E+03 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H(3s)	-81.8143	-.503310	-.548175	.379377	-.131005	.0123345	.0265299
H(3p)	-82.1981	-.551194	.00275721	.0217903	.00543228	.00224727	-.00356334
H(3d)	-83.6281	-.558730	.0544644	.0648205	-.0102811	-.00660745	.00511818
H(3p+3d)	-80.3437	.0712330	-.198360	.0790347	-.0161614	-.00946588	.0121131

The fit represents the H(3s) cross section with an rms deviation of 0.5%. The maximum deviation is 1.1% at 1.0E+03 eV/amu.

The fit represents the H(3p) cross section with an rms deviation of 0.2%. The maximum deviation is 0.5% at 2.0E+04 eV/amu.

The fit represents the H(3d) cross section with an rms deviation of 0.0%. The maximum deviation is 0.0% at 2.0E+04 eV/amu.

The fit represents the H(3p+3d) cross section with an rms deviation of 0.3%. The maximum deviation is 0.7% at 5.0E+02 eV/amu.

See appendix for Chebyshev fit details.

H + He –> H(3s,3d,3p)
H + He –> H(3p+3d)



Spectral Line Emission Cross Sections for

H + He -> H(Balmer-α,Balmer-β)

H(Balmer-α) Energy (eV/amu)	Cross Section (cm^2)	H(Balmer-β) Energy (eV/amu)	Cross Section (cm^2)
3.0E+01	5.35E-20	4.0E+01	2.19E-20
4.0E+01	2.73E-19	7.0E+01	8.68E-20
7.0E+01	9.03E-19	1.0E+02	1.51E-19
1.0E+02	1.36E-18	1.5E+02	4.00E-19
1.5E+02	3.47E-18	2.0E+02	7.19E-19
2.0E+02	5.11E-18	4.0E+02	1.88E-18
4.0E+02	8.99E-18	7.0E+02	2.28E-18
5.0E+02	9.14E-18	1.0E+03	2.17E-18
7.0E+02	8.77E-18	2.0E+03	1.73E-18
1.0E+03	7.86E-18	2.5E+03	1.60E-18
2.0E+03	6.22E-18		
4.0E+03	4.55E-18		
7.0E+03	3.35E-18		
1.0E+04	2.72E-18		
2.0E+04	1.64E-18		
3.0E+04	1.39E-18		
3.5E+04	1.35E-18		

<u>References:</u> 500, 601

<u>Accuracy:</u> 50%

<u>Notes:</u> (1) The Balmer-α wavelength is 656.3nm and the Balmer-β wavelength is 486.1nm. (2) These cross sections are for the total line emission including cascade contributions and branching corrections. Balmer-α cross sections are the sum of σ(3s) + 0.118 σ(3p) + σ(4D) where the factor 0.118 is the fraction of the 3p state decaying by Balmer-α radiation. Balmer-β emission cross sections are for the n=4->2 transitions with the branching ratio of the 4s to 3p transition being 0.584.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

H(Balmer-α) E_{min} = 3.0E+01 eV/amu, E_{max} = 3.5E+04 eV/amu
H(Balmer-β) E_{min} = 4.0E+01 eV/amu, E_{max} = 2.5E+03 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H(Balmer-α)	-81.8706	.929497	-1.66225	.554407	-.0518605	.00374127	-.0418064
H(Balmer-β)	-84.3982	2.15229	-.976804	-.0709114	.119448	.0571307	-.0675516

The fit represents the H(Balmer-α) cross section with an rms deviation of 13.9%. The maximum deviation is 35.9% at 1.0E+02 eV/amu.

The fit represents the H(Balmer-β) cross section with an rms deviation of 3.6%. The maximum deviation is 8.7% at 1.0E+02 eV/amu.

See appendix for Chebyshev fit details.

$H + He \rightarrow H_\alpha, H_\beta$



Excitation Cross Sections for

H + He -> H(4s,4p+4d)

H(4s)		H(4p+4d)	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
1.5E+02	1.30E-19	1.5E+02	3.07E-19
2.0E+02	2.60E-19	2.0E+02	5.35E-19
3.0E+02	4.77E-19	3.0E+02	1.06E-18
4.0E+02	6.75E-19	4.0E+02	1.30E-18
5.0E+02	8.10E-19	5.0E+02	1.37E-18
6.0E+02	8.78E-19	6.0E+02	1.38E-18
6.4E+02	8.79E-19	7.0E+02	1.37E-18
7.0E+02	8.72E-19	8.0E+02	1.35E-18
8.0E+02	8.40E-19	9.0E+02	1.33E-18
9.0E+02	7.99E-19	1.0E+03	1.31E-18
1.0E+03	7.56E-19	1.5E+03	1.23E-18
1.5E+03	6.06E-19	2.0E+03	1.19E-18
2.0E+03	5.21E-19	2.5E+03	1.16E-18
2.5E+03	4.62E-19		

Reference: 601

Accuracy: 50%

Notes: (1) As far as we are aware only one measurement has been made of the n = 4 excited state formation. Thus, the stated accuracy only includes accountable errors. (2) These data have ignored any cascade contributions.

Chebyshev Fitting Parameters for Cross Sections

H(4s) E_{min} = 1.5E+02 eV/amu, E_{max} = 2.5E+03 eV/amu
H(4p+4d) E_{min} = 1.5E+02 eV/amu, E_{max} = 2.5E+03 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H(4s)	-84.5175	.519837	-.602123	.107056	.0399676	.00471154	-.0330016
H(4p+4d)	-83.0777	.537485	-.440970	.159743	-.00409905	-.0305454	.0217549

The fit represents the H(4s) cross section with an rms deviation of 0.7%. The maximum deviation is 1.3% at 2.0E+03 eV/amu.

The fit represents the H(4p+4d) cross section with an rms deviation of 0.6%. The maximum deviation is 1.1% at 1.5E+03 eV/amu.

See appendix for Chebyshev fit details.

H + He -> H(4s,4p+4d)


Cross Section vs. Energy
10^{-17}
10^{-18}
10^{-19}
Cross Section (cm^2)
10^2
10^3
10^4
Energy (eV/amu)
Δ = H(4p+4d)
× = H(4s)
Recommended Data
Chebyshev Fit

Excitation Cross Sections for

H + He -> He(3^1P,3^3P,3^3D)

He(3^1P)		He(3^3P)		He(3^3D)	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
1.0E+04	5.60E-20	1.0E+04	9.67E-19	1.0E+04	1.60E-19
1.3E+04	1.16E-19	1.3E+04	1.16E-18	1.3E+04	2.19E-19
1.5E+04	1.79E-19	1.5E+04	1.18E-18	1.4E+04	2.37E-19
1.8E+04	2.19E-19	1.8E+04	1.04E-18	1.5E+04	2.32E-19
2.0E+04	2.53E-19	2.0E+04	8.64E-19	1.8E+04	1.95E-19
2.3E+04	2.83E-19	2.3E+04	7.19E-19	2.0E+04	1.62E-19
2.5E+04	3.14E-19	2.5E+04	5.93E-19	2.3E+04	1.37E-19
2.8E+04	3.46E-19	2.8E+04	4.92E-19	2.5E+04	1.17E-19
3.0E+04	3.75E-19	3.0E+04	4.07E-19	2.8E+04	1.00E-19
3.3E+04	4.01E-19	3.3E+04	3.33E-19	3.0E+04	8.62E-20
3.5E+04	4.29E-19	3.5E+04	2.72E-19	3.3E+04	7.57E-20
				3.5E+04	6.86E-20

References: 166, 678, 683, 684

Accuracy: Unknown

Notes: (1) Only one absolute measurement of cross sections for excitation of these states has been made. See: Van Eck et al. (Ref. 678). These data have been corrected in the review papers of de Heer (Ref. 166) and Thomas (Ref. 684). The corrected values are probably too low by a factor of 2-3.

Chebyshev Fitting Parameters for Cross Sections

He(3^1P) E_{min} = 1.0E+04 eV/amu, E_{max} = 3.5E+04 eV/amu
He(3^3P) E_{min} = 1.0E+04 eV/amu, E_{max} = 3.5E+04 eV/amu
He(3^3D) E_{min} = 1.0E+04 eV/amu, E_{max} = 3.5E+04 eV/amu

	A0	A1	A2	A3	A4	A5	A6
He(3^1P)	-86.1894	.957985	-.222479	.0796768	-.00575983	-.0194864	.0121076
He(3^3P)	-83.6070	-.654163	-.317139	.0375222	-3.81314E-04	-.0176396	.00667369
He(3^3D)	-86.8654	-.503446	-.290764	.103755	-.00276954	-.0224516	.0247251

The fit represents the He(3^1P) cross section with an rms deviation of 0.3%. The maximum deviation is 0.6% at 1.8E+04 eV/amu.

The fit represents the He(3^3P) cross section with an rms deviation of 0.2%. The maximum deviation is 0.3% at 2.0E+04 eV/amu.

The fit represents the He(3^3D) cross section with an rms deviation of 0.4%. The maximum deviation is 0.8% at 1.8E+04 eV/amu.

See appendix for Chebyshev fit details.

H + He –> He(3^1P,3^3P,3^3D)



Excitation Cross Sections for

H + He -> He(4^1S,4^1P,4^1D)

He(4^1S) Energy (eV/amu)	He(4^1S) σ (cm^2)	He(4^1P) Energy (eV/amu)	He(4^1P) σ (cm^2)	He(4^1D) Energy (eV/amu)	He(4^1D) σ (cm^2)
1.0E+04	2.87E-19	1.0E+04	6.19E-20	1.0E+04	5.34E-20
1.5E+04	3.64E-19	1.3E+04	1.30E-19	1.5E+04	8.41E-20
2.0E+04	4.09E-19	1.5E+04	1.92E-19	2.0E+04	9.12E-20
2.5E+04	4.34E-19	2.0E+04	3.03E-19	2.5E+04	8.01E-20
3.0E+04	4.34E-19	2.5E+04	3.80E-19	3.0E+04	7.68E-20
3.5E+04	4.27E-19	3.0E+04	4.48E-19	3.0E+04	7.72E-20
4.0E+04	4.08E-19	3.5E+04	5.13E-19	3.7E+04	8.87E-20
4.5E+04	3.88E-19			4.0E+04	8.79E-20
5.0E+04	3.68E-19			4.5E+04	8.51E-20
5.5E+04	3.51E-19			5.0E+04	8.48E-20
6.0E+04	3.33E-19			5.5E+04	8.24E-20
6.5E+04	3.10E-19			6.0E+04	7.83E-20
7.0E+04	2.81E-19			6.5E+04	7.27E-20
7.5E+04	2.48E-19			7.0E+04	6.66E-20
8.0E+04	2.20E-19			7.5E+04	6.28E-20
8.5E+04	1.97E-19			8.0E+04	6.00E-20
9.0E+04	1.80E-19			8.5E+04	5.81E-20
9.5E+04	1.69E-19			9.0E+04	5.62E-20
1.0E+05	1.61E-19			9.5E+04	5.47E-20
				1.0E+05	5.34E-20

References: 166, 677, 678, 679, 684

Accuracy: E > $2x10^3$ eV/amu - 40%
E < $2x10^3$ eV/amu - Unknown

Notes: (1) The data have been obtained by measuring the line emissions for the following transitions: (a) 4^1S -> 2^2P, 504.7nm, (b) 4^2D -> 2^2P, 492.2nm, (c) 4^2P -> 2^2S, 396.5nm. (2) The cross section data have been corrected for cascades using known transition probabilities. (3) The corrected data of Van Eck et al. (Ref. 678) - (see corrections of de Heer Ref. 166 and Thomas Ref. 684) have been normalized to the data of Blair et al. (Ref. 677) at $2x10^3$ eV/amu for the 4^2S and 4^2D states.

Chebyshev Fitting Parameters for Cross Sections

He(4^1S) E_{min} = 1.0E+04 eV/amu, E_{max} = 1.0E+05 eV/amu
He(4^1P) E_{min} = 1.0E+04 eV/amu, E_{max} = 3.5E+04 eV/amu
He(4^1D) E_{min} = 1.0E+04 eV/amu, E_{max} = 1.0E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
He(4^1S)	-85.2953	-.236370	-.375570	-.0526312	1.43777E-04	-.00898368	.0314058
He(4^1P)	-85.8813	.991657	-.212955	.0558510	-.00900116	.00987058	-.00878102
He(4^1D)	-88.1942	-.0703862	-.233023	.0274405	-.0616908	.0587687	.0287005

The fit represents the He(4^1S) cross section with an rms deviation of 1.7%.
The maximum deviation is 2.7% at 5.0E+04 eV/amu.
The fit represents the He(4^1P) cross section with an rms deviation of 0.1%.
The maximum deviation is 5.0% at 1.2E+04 eV/amu.
The fit represents the He(4^2D) cross section with an rms deviation of 3.1%.
The maximum deviation is 5.3% at 2.9E+04 eV/amu.

See appendix for Chebyshev fit details.

H + He -> He(4 ¹S,4 ¹P,4 ¹D)



Excitation Cross Sections for

H + He -> He(4^3S,4^3P,4^3D)

He(4^3S) Energy (eV/amu)	σ (cm^2)	He(4^3P) Energy (eV/amu)	σ (cm^2)	He(4^3D) Energy (eV/amu)	σ (cm^2)
1.0E+04	3.69E-19	1.0E+04	8.78E-19	1.0E+04	1.59E-19
1.5E+04	8.38E-19	1.2E+04	1.08E-18	1.5E+04	2.06E-19
2.0E+04	1.00E-18	1.4E+04	1.15E-18	2.0E+04	1.71E-19
3.0E+04	8.70E-19	1.5E+04	1.15E-18	2.5E+04	1.33E-19
4.0E+04	5.43E-19	2.0E+04	8.82E-19	3.0E+04	9.92E-20
5.0E+04	3.56E-19	2.5E+04	6.05E-19	3.5E+04	8.03E-20
6.0E+04	2.48E-19	3.0E+04	4.11E-19	4.0E+04	6.71E-20
7.0E+04	1.79E-19	3.5E+04	3.00E-19	5.0E+04	5.21E-20
8.0E+04	1.25E-19			6.0E+04	4.36E-20
9.0E+04	9.92E-20			7.0E+04	3.80E-20
1.0E+05	8.50E-20			8.0E+04	3.38E-20
				9.0E+04	3.05E-20
				1.0E+05	2.80E-20

References: 166, 677, 678, 684

Accuracy: $E > 2x10^3$ eV/amu - 40%
$E < 2x10^3$ eV/amu - Unknown

Notes: (1) The data have been obtained by measuring the line emissions for the following transitions: (a) 4^3S -> 2^3P, 471.3nm, (b) 4^3P -> 2^3S, 318.8nm, (c) 4^3D -> 2^3P, 447.1nm. (2) The cross section data have been corrected for cascading using known transition probabilities. (3) The corrected data of Van Eck et al. (Ref. 678) - (see corrections of de Heer Ref. 166 and Thomas, Ref. 684) have been normalized to the data of Blair et al. (Ref. 677) at $2x10^3$ eV/amu for the 4^1S and 4^1D states.

Chebyshev Fitting Parameters for Cross Sections

He(4^3S) E_{min} = 1.0E+04 eV/amu, E_{max} = 1.0E+05 eV/amu
He(4^3P) E_{min} = 1.0E+04 eV/amu, E_{max} = 3.5E+04 eV/amu
He(4^3D) E_{min} = 1.0E+04 eV/amu, E_{max} = 1.0E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
He(4^3S)	-84.9236	-.910048	-.757778	.164455	.0260068	-7.50878E-04	.00468518
He(4^3P)	-83.6191	-.587215	-.319615	.0516334	.0108396	-.00145124	.00476885
He(4^3D)	-87.8723	-1.03617	-.177049	.184100	-.0530501	-.0167810	.0122418

The fit represents the He(4^3S) cross section with an rms deviation of 2.2%. The maximum deviation is 2.6% at 8.0E+04 eV/amu.

The fit represents the He(4^3P) cross section with an rms deviation of 0.2%. The maximum deviation is 0.5% at 1.4E+04 eV/amu.

The fit represents the He(4^3D) cross section with an rms deviation of 0.6%. The maximum deviation is 1.1% at 2.0E+04 eV/amu.

See appendix for Chebyshev fit details.

H + He –> He(4^3S,4^3P,4^3D)



Excitation Cross Sections for

$H^+ + H \rightarrow H^+ + H(2s,2p)$

H(2s) Energy (eV/amu)	H(2s) Cross Section (cm^2)	H(2p) Energy (eV/amu)	H(2p) Cross Section (cm^2)
5.0E+03	5.42E-18	5.0E+02	2.43E-18
6.0E+03	5.38E-18	6.0E+02	8.90E-18
7.0E+03	5.36E-18	7.0E+02	1.64E-17
8.0E+03	5.40E-18	8.0E+02	2.03E-17
9.0E+03	5.43E-18	9.0E+02	2.25E-17
1.0E+04	5.54E-18	1.0E+03	2.41E-17
1.2E+04	5.87E-18	1.5E+03	2.73E-17
1.5E+04	8.62E-18	2.0E+03	2.80E-17
2.0E+04	1.04E-17	3.0E+03	2.91E-17
2.5E+04	1.06E-17	4.0E+03	2.97E-17
		5.0E+03	3.02E-17
		6.0E+03	3.06E-17
		7.0E+03	2.98E-17
		8.0E+03	2.78E-17
		9.0E+03	2.56E-17
		1.0E+04	2.40E-17
		1.1E+04	2.35E-17
		1.5E+04	2.81E-17
		2.0E+04	3.65E-17
		3.0E+04	5.50E-17

<u>References:</u> H(2s) - 12, 14
H(2p) - 14, 17, 712

<u>Accuracy:</u> H(2s) - Unknown
H(2p) - 40%

<u>Notes:</u> (1) Since the cascading into the 2p state is less than 5%, the cross section for excitation of the 2p state is essentially the same as Lyman-α emission cross section. The 2p data has not been corrected for cascading. (2) The two sets of data for the 2s state differ by a factor of 2 to 3.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

H(2s) E_{min} = 5.0E+03 eV/amu, E_{max} = 2.5E+04 eV/amu
H(2p) E_{min} = 5.0E+02 eV/amu, E_{max} = 3.0E+04 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H(2s)	-78.9900	.401365	.0930849	-.0759638	-.0649979	.00791412	.0468211
H(2p)	-76.6674	.854989	-.357230	.529756	-.161903	.168930	-.136550

The fit represents the H(2s) cross section with an rms deviation of 2.3%. The maximum deviation is 4.8% at 1.2E+04 eV/amu.

The fit represents the H(2p) cross section with an rms deviation of 3.7%. The maximum deviation is 6.3% at 1.1E+04 eV/amu.

See appendix for Chebyshev fit details.

$H^+ + H \rightarrow H^+ + H(2s,2p)$



Excitation Rate Coefficients for

$H^+ + H \rightarrow H^+ + H(2p)$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H Temp. (eV)	Equal Temp.	H Temp. (eV) 500.	750.	1000.	1250.	5000.	10000.	20000.
5.0E+02	1.16E-09	1.16E-09	1.39E-09	1.58E-09	1.75E-09	3.33E-09	4.11E-09*	3.43E-09#
7.5E+02	1.58E-09	1.39E-09	1.58E-09	1.75E-09	1.90E-09	3.41E-09	4.11E-09*	3.41E-09#
1.0E+03	1.90E-09	1.58E-09	1.75E-09	1.90E-09	2.03E-09	3.48E-09	4.11E-09*	3.39E-09#
1.3E+03	2.16E-09	1.75E-09	1.90E-09	2.03E-09	2.16E-09	3.55E-09	4.11E-09*	3.36E-09#
1.5E+03	2.38E-09	1.90E-09	2.03E-09	2.16E-09	2.27E-09	3.61E-09*	4.11E-09*	3.34E-09#
1.8E+03	2.59E-09	2.03E-09	2.16E-09	2.27E-09	2.38E-09	3.67E-09*	4.10E-09*	3.31E-09#
2.0E+03	2.80E-09	2.16E-09	2.27E-09	2.38E-09	2.49E-09	3.73E-09*	4.10E-09*	3.29E-09#
3.0E+03	3.48E-09	2.59E-09	2.70E-09	2.80E-09	2.89E-09	3.91E-09*	4.06E-09*	3.20E-09#
4.0E+03	3.91E-09*	2.99E-09	3.08E-09	3.17E-09	3.25E-09	4.03E-09*	4.00E-09*	3.11E-09#
5.0E+03	4.09E-09*	3.33E-09	3.41E-09	3.48E-09	3.55E-09	4.09E-09*	3.93E-09#	3.02E-09#
6.0E+03	4.10E-09*	3.61E-09*	3.67E-09*	3.73E-09*	3.78E-09*	4.11E-09*	3.85E-09#	2.94E-09#
7.0E+03	4.00E-09*	3.83E-09*	3.87E-09*	3.91E-09*	3.94E-09*	4.10E-09*	3.76E-09#	2.85E-09#
8.0E+03	3.85E-09#	3.98E-09*	4.00E-09*	4.03E-09*	4.05E-09*	4.06E-09*	3.67E-09#	2.77E-09#
9.0E+03	3.67E-09#	4.07E-09*	4.08E-09*	4.09E-09*	4.10E-09*	4.00E-09*	3.58E-09#	2.70E-09#
1.0E+04	3.48E-09#	4.11E-09*	4.11E-09*	4.11E-09*	4.11E-09*	3.93E-09#	3.48E-09#	2.62E-09#
1.2E+04	3.11E-09#	4.08E-09*	4.07E-09*	4.06E-09*	4.05E-09*	3.76E-09#	3.29E-09#	2.48E-09#
1.5E+04	2.62E-09#	3.89E-09#	3.87E-09#	3.85E-09#	3.83E-09#	3.48E-09#	3.02E-09#	2.29E-09#
2.0E+04	2.02E-09#	3.43E-09#	3.41E-09#	3.39E-09#	3.36E-09#	3.02E-09#	2.62E-09#	2.02E-09#

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 5.0E+02 eV, E_{max} = 2.0E+04 eV

H Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
500.	-39.7100	.631450	-.153568	-.0909606	-.0319929	2.01118E-04	.00981582
750.	-39.5923	.543848	-.128283	-.0971754	-.0299354	.00227565	.00859152
1000.	-39.4997	.476362	-.113723	-.0995434	-.0281721	.00377301	.00746751
1250.	-39.4237	.421755	-.105533	-.0997854	-.0264000	.00472594	.00647274
5000.	-38.9001	.0101105	-.108849	-.0609619	-.0111639	.00213821	.00143014
10000.	-38.8452	-.187122	-.105864	-.0371107	-.00678512	2.66271E-04	5.36963E-04
20000.	-39.3124	-.240533	-.0956162	-.0251663	-.00414891	-1.62169E-04	1.56753E-04
Equal Temp.	-39.7457	.353158	-.421145	-.103566	-3.86751E-04	.272186	-.00215387

See appendix for Chebyshev fit details.

$H^+ + H \rightarrow H^+ + H(2p)$

Maxwellian – Maxwellian



Excitation Cross Sections for

$H^+ + H \rightarrow H^+ + H(n=2,3,4)$

H(n=2)		H(n=3)		H(n=4)	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
1.5E+04	3.43E-17	1.5E+04	1.09E-17	2.5E+04	7.69E-18
2.0E+04	5.22E-17	2.0E+04	1.29E-17	3.0E+04	8.68E-18
2.5E+04	6.64E-17	2.5E+04	1.53E-17	4.0E+04	1.02E-17
3.0E+04	7.89E-17	3.0E+04	1.80E-17	5.0E+04	1.12E-17
4.0E+04	9.59E-17	4.0E+04	2.33E-17	6.0E+04	1.18E-17
5.0E+04	1.05E-16	5.0E+04	2.54E-17	7.0E+04	1.19E-17
5.5E+04	1.07E-16	6.0E+04	2.51E-17	8.0E+04	1.18E-17
6.0E+04	1.07E-16	7.0E+04	2.42E-17	9.0E+04	1.14E-17
7.0E+04	1.03E-16	8.0E+04	2.32E-17	1.0E+05	1.10E-17
8.0E+04	9.72E-17	9.0E+04	2.21E-17	1.5E+05	9.06E-18
9.0E+04	9.28E-17	1.0E+05	2.14E-17	2.0E+05	7.69E-18
1.0E+05	8.90E-17	1.5E+05	1.70E-17		
1.5E+05	7.48E-17	2.0E+05	1.36E-17		
2.0E+05	6.53E-17				

<u>References:</u> 709, 710, 711

<u>Accuracy:</u> 40%

<u>Notes:</u> (1) These cross sections were determined by measuring the energy loss of the incident proton beam after passing through an H target. The energy resolution was not sufficient to separate the nl levels. (2) To obtain absolute cross sections the results were normalized to the Born approximation of Bates and Griffing (Ref. 711) at 200 keV.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

H(n=2)	E_{min}	=	1.5E+04	eV/amu,	E_{max}	=	2.0E+05 eV/amu
H(n=3)	E_{min}	=	1.5E+04	eV/amu,	E_{max}	=	2.0E+05 eV/amu
H(n=4)	E_{min}	=	2.5E+04	eV/amu,	E_{max}	=	2.0E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H(n=2)	-74.4276	.262934	-.386115	.0616937	.0293129	-.00255599	-.0190829
H(n=3)	-77.2728	.149236	-.351549	-.0157777	.0571627	-.0247442	-.0161701
H(n=4)	-78.4118	-.00702352	-.213691	.00801652	.0182541	-8.70179E-04	-.00538651

The fit represents the H(n=2) cross section with an rms deviation of 0.5%. The maximum deviation is 0.9% at 8.0E+04 eV/amu.

The fit represents the H(n=3) cross section with an rms deviation of 1.2%. The maximum deviation is 1.5% at 7.0E+04 eV/amu.

The fit represents the H(n=4) cross section with an rms deviation of 0.2%. The maximum deviation is 0.2% at 5.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$H^+ + H \rightarrow H(n=2,3,4)$ Target



Dissociative Excitation Cross Sections for

$H^+ + H_2 \rightarrow H^+ + H(2s,2p) + H$

H(2s)		H(2p)	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
5.0E+03	1.32E-18	4.0E+03	6.36E-18
7.0E+03	3.59E-18	7.0E+03	1.84E-17
1.0E+04	4.70E-18	1.0E+04	2.76E-17
1.5E+04	5.73E-18	1.5E+04	3.12E-17
2.0E+04	6.38E-18	2.0E+04	2.96E-17
4.0E+04	7.69E-18	4.0E+04	2.14E-17
7.0E+04	8.00E-18	7.0E+04	1.48E-17
1.0E+05	7.88E-18	1.0E+05	1.21E-17
1.5E+05	7.04E-18	1.5E+05	9.23E-18
2.0E+05	6.23E-18	2.0E+05	7.42E-18
4.0E+05	4.18E-18	4.0E+05	4.34E-18
7.0E+05	2.75E-18	7.0E+05	2.81E-18
1.0E+06	2.00E-18	1.0E+06	2.14E-18

References: 122, 124, 128, 649

Accuracy: 50%

Notes: (1) The 2s cross section has been measured by field-induced emission of Lyman-α radiation (2s -> 2p -> 1s). If cascade is neglected, this is equal to the cross section for H(2s) formation. (2) The 2p cross section was determined by measuring the Lyman-α emission. If cascade is neglected, this is equal to the cross section for H(2p) formation.

Chebyshev Fitting Parameters for Cross Sections

H(2s) E_{min} = 5.0E+03 eV/amu, E_{max} = 1.0E+06 eV/amu
H(2p) E_{min} = 4.0E+03 eV/amu, E_{max} = 1.0E+06 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H(2s)	-80.1116	-.0216909	-.735933	.117161	-.0952706	.0960395	-.0678792
H(2p)	-78.5492	-.910686	-.758592	.347330	-.131263	.0249389	.0249642

The fit represents the H(2s) cross section with an rms deviation of 3.1%. The maximum deviation is 5.9% at 1.0E+04 eV/amu.

The fit represents the H(2p) cross section with an rms deviation of 1.6%. The maximum deviation is 2.5% at 7.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$H^+ + H_2 \rightarrow H^+ + H(2s,2p)$



Dissociative Excitation Cross Sections for

$H^+ + H_2 \rightarrow H^+ + H(3p) + H$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.5E+04	1.70E+08	1.99E-18
1.6E+04	1.76E+08	2.14E-18
1.7E+04	1.81E+08	2.25E-18
1.8E+04	1.86E+08	2.34E-18
1.9E+04	1.91E+08	2.38E-18
2.0E+04	1.96E+08	2.39E-18
2.1E+04	2.01E+08	2.36E-18
2.2E+04	2.06E+08	2.31E-18
2.3E+04	2.11E+08	2.24E-18
2.4E+04	2.15E+08	2.16E-18
2.5E+04	2.20E+08	2.07E-18
2.6E+04	2.24E+08	1.95E-18
2.7E+04	2.28E+08	1.81E-18
2.8E+04	2.32E+08	1.65E-18
2.9E+04	2.37E+08	1.45E-18
3.0E+04	2.41E+08	1.20E-18

Reference: 122

Accuracy: 30%

Note: The cross sections for H(3p) formation deduced from emission measurements on the (unsubstantiated) assumption that cascade may be neglected.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.5E+04 eV/amu, E_{max} = 3.0E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-81.5559	-.198605	-.205491	-.0404366	-.0225806	-.0130785	-.00450361

The fit represents the above cross section with an rms deviation of 0.2%. The maximum deviation is 0.4% at 2.7E+04 eV/amu.

See appendix for Chebyshev fit details.

$H^+ + H_2 \rightarrow H(3p)$ Target



Spectral Line Emission Cross Sections for

$H^+ + H_2 \rightarrow H(\text{Lyman-}\alpha)$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
5.0E+02	3.11E+07	6.56E-18
7.0E+02	3.68E+07	9.30E-18
1.0E+03	4.39E+07	1.31E-17
1.5E+03	5.38E+07	1.86E-17
2.0E+03	6.21E+07	2.35E-17
4.0E+03	8.79E+07	3.74E-17
7.0E+03	1.16E+08	4.98E-17
1.0E+04	1.39E+08	5.68E-17
1.5E+04	1.70E+08	6.17E-17
2.0E+04	1.96E+08	5.50E-17
4.0E+04	2.78E+08	2.78E-17
7.0E+04	3.68E+08	1.77E-17
1.0E+05	4.39E+08	1.40E-17
1.4E+05	5.20E+08	1.20E-17

References: 188, 194, 195, 676

Accuracy: 50%

Note: Cross sections correspond to 121.57nm emission resulting from dissociative excitation of H_2 by proton impact.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 5.0E+02 eV/amu, E_{max} = 1.4E+05 eV/amu

A0	A1	A2	A3	A4	A5	A6
-76.9474	.268573	-.911806	-.0361054	.132508	.0899171	-.00698679

The fit represents the above cross section with an rms deviation of 4.5%. The maximum deviation is 10.4% at 4.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$H^+ + H_2 \rightarrow Ly_\alpha$ Target



Spectral Line Emission Cross Sections for

$H^+ + H_2 \rightarrow H(\text{Balmer-}\alpha, \text{Balmer-}\beta, \text{Balmer-}\gamma)$

H_α		H_β		H_γ	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
1.5E+03	2.09E-19	5.0E+03	1.75E-19	1.0E+04	1.00E-19
2.0E+03	2.70E-19	6.0E+03	2.37E-19	1.4E+04	1.09E-19
4.0E+03	5.80E-19	7.0E+03	2.98E-19	1.5E+04	1.08E-19
7.0E+03	1.51E-18	8.0E+03	3.53E-19	2.0E+04	9.91E-20
1.0E+04	2.24E-18	9.0E+03	3.95E-19	3.0E+04	7.64E-20
1.5E+04	2.62E-18	1.0E+04	4.28E-19	4.0E+04	6.00E-20
2.0E+04	2.44E-18	1.5E+04	4.70E-19	5.0E+04	4.93E-20
4.0E+04	1.55E-18	2.0E+04	4.22E-19	6.0E+04	4.21E-20
7.0E+04	9.06E-19	3.0E+04	3.00E-19	7.0E+04	3.69E-20
1.0E+05	6.50E-19	4.0E+04	2.29E-19	8.0E+04	3.29E-20
1.5E+05	4.57E-19	5.0E+04	1.88E-19	9.0E+04	2.96E-20
2.0E+05	3.57E-19	6.0E+04	1.59E-19	1.0E+05	2.70E-20
4.0E+05	1.96E-19	7.0E+04	1.37E-19	1.5E+05	1.87E-20
7.0E+05	1.20E-19	8.0E+04	1.21E-19	2.0E+05	1.46E-20
1.0E+06	8.90E-20	9.0E+04	1.08E-19	3.0E+05	1.02E-20
		1.0E+05	9.78E-20	4.0E+05	7.97E-21
		1.5E+05	6.72E-20	5.0E+05	6.52E-21
		2.0E+05	5.18E-20	6.0E+05	5.57E-21
		3.0E+05	3.60E-20	7.0E+05	4.88E-21
		4.0E+05	2.75E-20	8.0E+05	4.38E-21
		5.0E+05	2.25E-20	9.0E+05	3.98E-21
		6.0E+05	1.90E-20	1.0E+06	3.65E-21
		7.0E+05	1.66E-20		
		8.0E+05	1.46E-20		
		9.0E+05	1.32E-20		
		1.0E+06	1.19E-20		

<u>References:</u> 133, 139, 655

<u>Accuracy:</u> 50%

<u>Notes:</u> (1) The data correspond to Balmer-series emission resulting from dissociative excitation of H_2, and have been normalized in the high energy region to data of Edwards and Thomas (Ref. 655). (2) Differences in magnitudes of the reported cross sections for each transition are due to differences in calibration between the experimental systems. (3) The wavelengths are as follows: H_α= 656.28nm; H_β=486.13nm; H_γ=434.05nm.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

H_α E_{min} = 1.5E+03 eV/amu, E_{max} = 1.0E+06 eV/amu
H_β E_{min} = 6.0E+03 eV/amu, E_{max} = 1.0E+06 eV/amu
H_γ E_{min} = 1.0E+04 eV/amu, E_{max} = 1.0E+06 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H_α	-84.6647	-.627902	-1.28581	.366073	.150564	-.204615	.0536705
H_β	-87.6626	-1.78903	-.434179	.278457	-.129747	.0301375	.0113784
H_γ	-90.3588	-1.79519	-.168084	.119053	-.0545231	.0225899	-.00228634

The fit represents the H_α cross section with an rms deviation of 5.4%.
The maximum deviation is 12.3% at 4.0E+03 eV/amu.
The fit represents the H_β cross section with an rms deviation of 1.4%.
The maximum deviation is 3.1% at 4.0E+04 eV/amu.
The fit represents the H_γ cross section with an rms deviation of 0.4%.
The maximum deviation is 0.8% at 5.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$H^+ + H_2 \rightarrow H_\alpha, H_\beta, H_\gamma$



Spectral Band Emission Cross Sections for

$$H^+ + H_2 \rightarrow H^+ + H_2^*$$

160.6nm Band		418.0nm Band	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
7.0E+03	2.17E-18	1.5E+05	8.06E-21
8.0E+03	2.31E-18	2.0E+05	6.25E-21
9.0E+03	2.44E-18	2.5E+05	5.06E-21
1.0E+04	2.56E-18	3.0E+05	4.21E-21
1.5E+04	3.00E-18	3.5E+05	3.62E-21
2.0E+04	3.30E-18	4.0E+05	3.17E-21
3.0E+04	3.69E-18	5.0E+05	2.54E-21
4.0E+04	3.84E-18	6.0E+05	2.11E-21
5.0E+04	3.87E-18	7.0E+05	1.81E-21
6.0E+04	3.83E-18	8.0E+05	1.59E-21
7.0E+04	3.76E-18	9.0E+05	1.39E-21
8.0E+04	3.67E-18		
9.0E+04	3.55E-18		
1.0E+05	3.42E-18		
1.3E+05	3.01E-18		

<u>References:</u> 194, 654

<u>Accuracy:</u> 50%

<u>Note:</u> The 160.6nm Band is the Lyman band of H_2 [B $^1\Sigma_u^+$ -> X $^1\Sigma_g^+$, (4,11), (5,12) and (6,13)]. The 418.0nm Band of H_2 is for the transitions [3d $^1\pi_g$ -> 2p $^1\Sigma_u^+$ (1,0)].

Chebyshev Fitting Parameters for Cross Sections

160.6 nm E_{min} = 7.0E+03 eV/amu, E_{max} = 1.3E+05 eV/amu
418.0 nm E_{min} = 1.5E+05 eV/amu, E_{max} = 9.0E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
160.6 nm	-80.6464	.194014	-.181339	-.0301739	-.00212964	3.13808E-04	-.00210888
418.0 nm	-94.2526	-.881302	-.0131335	.00371515	-.00433338	-6.45602E-04	-.00195489

The fit represents the 160.6nm emission cross section with an rms deviation of 0.2%.
The maximum deviation is 0.3% at 2.0E+04 eV/amu.

The fit represents the 418.0nm emission cross section with an rms deviation of 0.2%.
The maximum deviation is 0.3% at 7.0E+05 eV/amu.

See appendix for Chebyshev fit details.

$H^+ + H_2 \rightarrow 160.6, 418.0$ nm



Excitation Cross Sections for

$H^+ + He \rightarrow H^+ + He^*(2^1S, 2^1P)$
$H^+ + He^*(2^1S + 2^1P)$

$He(2^1S)$ Energy (eV/amu)	σ (cm^2)	$He(2^1P)$ Energy (eV/amu)	σ (cm^2)	$He(2^1S + 2^1P)$ Energy (eV/amu)	σ (cm^2)
2.5E+04	3.79E-18	2.3E+04	2.00E-18	2.3E+04	5.38E-18
3.0E+04	4.52E-18	2.5E+04	2.44E-18	3.0E+04	7.52E-18
3.5E+04	5.28E-18	3.0E+04	3.44E-18	4.0E+04	1.12E-17
4.0E+04	6.05E-18	3.5E+04	4.61E-18	5.0E+04	1.47E-17
4.5E+04	6.68E-18	4.0E+04	5.76E-18	6.0E+04	1.50E-17
5.0E+04	6.87E-18	4.5E+04	6.91E-18	7.0E+04	1.48E-17
5.5E+04	6.59E-18	5.0E+04	7.65E-18	8.0E+04	1.55E-17
6.0E+04	6.22E-18	5.5E+04	7.96E-18	9.0E+04	1.71E-17
7.0E+04	5.33E-18	6.0E+04	8.39E-18	1.0E+05	1.79E-17
8.0E+04	4.94E-18	7.0E+04	9.39E-18	1.3E+05	1.62E-17
9.0E+04	5.16E-18	8.0E+04	1.08E-17		
1.0E+05	5.43E-18	9.0E+04	1.23E-17		
		1.0E+05	1.36E-17		
		1.2E+05	1.43E-17		
		1.5E+05	1.35E-17		
		2.0E+05	1.21E-17		
		2.5E+05	1.10E-17		
		3.0E+05	1.01E-17		
		3.5E+05	9.32E-18		
		4.0E+05	8.71E-18		
		5.0E+05	7.80E-18		
		6.0E+05	6.91E-18		
		7.0E+05	6.31E-18		
		8.0E+05	5.88E-18		
		9.0E+05	5.52E-18		
		1.0E+06	5.23E-18		

<u>References:</u> 665, 680, 681, 682

<u>Accuracy:</u> $E < 1.5 \times 10^5$ eV/amu - 50%
$E > 1.5 \times 10^5$ eV/amu - 30%

<u>Note:</u> For angular distributions see Park et al. (Ref. 681), and Kvale et al. (Ref. 665).

Chebyshev Fitting Parameters for Cross Sections

$He^*(2^1S)$	E_{min} = 2.5E+04 eV/amu,	E_{max} = 1.0E+05 eV/amu
$He^*(2^1P)$	E_{min} = 2.3E+04 eV/amu,	E_{max} = 1.0E+06 eV/amu
$He^*(2^1S + 2^1P)$	E_{min} = 2.3E+04 eV/amu,	E_{max} = 1.3E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
$He^*(2^1S)$	-79.6082	.117213	-.178720	.0616470	.0764221	.00195815	-.0273944
$He^*(2^1P)$	-79.1761	.318296	-.694295	.161212	.0240083	-.00395403	-.0285411
$He^*(2^1S + 2^1P)$	-77.9705	.598490	-.190056	.0136746	.00624072	-.0619807	-.0440015

The fit represents the $He^*(2^1S)$ cross section with an rms deviation of 0.5%.
The maximum deviation is 0.7% at 8.0E+04 eV/amu.
The fit represents the $He^*(2^1P)$ cross section with an rms deviation of 3.7%.
The maximum deviation is 8.4% at 7.0E+04 eV/amu.
The fit represents the $He^*(2^1S + 2^1P)$ cross section with an rms deviation of 1.6%.
The maximum deviation is 2.0% at 7.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$H^+ + He \rightarrow He(2^1S, 2^1P, 2^1S + 2^1P)$



Excitation Rate Coefficients for

$He + H^+ \rightarrow H^+ + He^*(2^1S)$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H Temp. (eV)	He Energy (eV/amu) 30000.	40000.	50000.	60000.	70000.	80000.	100000.
1.0E+00	1.09E-09	1.68E-09	2.13E-09	2.11E-09	1.96E-09	1.94E-09	1.19E-09#
2.0E+00	1.09E-09	1.68E-09	2.13E-09	2.11E-09	1.96E-09	1.95E-09	1.18E-09#
4.0E+00	1.09E-09	1.68E-09	2.13E-09	2.11E-09	1.96E-09	1.95E-09	1.18E-09#
7.0E+00	1.09E-09	1.68E-09	2.12E-09	2.11E-09	1.96E-09	1.95E-09	1.18E-09#
1.0E+01	1.09E-09	1.68E-09	2.12E-09	2.11E-09	1.96E-09	1.95E-09	1.17E-09#
2.0E+01	1.09E-09	1.68E-09	2.12E-09	2.11E-09	1.97E-09	1.96E-09	1.16E-09#
4.0E+01	1.09E-09	1.68E-09	2.11E-09	2.11E-09	1.97E-09	1.96E-09	1.15E-09#
7.0E+01	1.09E-09	1.68E-09	2.10E-09	2.10E-09	1.97E-09	1.97E-09	1.14E-09#
1.0E+02	1.09E-09	1.68E-09	2.09E-09	2.10E-09	1.98E-09	1.98E-09	1.13E-09#
2.0E+02	1.07E-09	1.68E-09	2.08E-09	2.09E-09	1.99E-09	2.00E-09	1.11E-09#
4.0E+02	1.04E-09	1.68E-09	2.04E-09	2.07E-09	2.00E-09	2.00E-09	1.07E-09#
7.0E+02	1.04E-09*	1.67E-09	2.00E-09	2.06E-09	2.01E-09	1.95E-09	1.04E-09#
1.0E+03	1.05E-09*	1.66E-09	1.97E-09	2.04E-09	2.00E-09	1.89E-09*	1.02E-09#
2.0E+03	1.10E-09*	1.61E-09	1.89E-09	1.97E-09	1.91E-09	1.71E-09*	9.77E-10#
4.0E+03	1.17E-09*	1.55E-09	1.75E-09	1.79E-09*	1.69E-09*	1.48E-09*	9.29E-10#
7.0E+03	1.23E-09*	1.47E-09	1.57E-09*	1.56E-09*	1.45E-09*	1.28E-09*	8.75E-10#
1.0E+04	1.23E-09*	1.38E-09*	1.43E-09*	1.39E-09*	1.29E-09*	1.15E-09*	8.27E-10#
2.0E+04	1.15E-09*	1.15E-09*	1.12E-09*	1.06E-09*	9.83E-10#	8.92E-10#	7.02E-10#

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
30000.	-41.2314	.0438795	.0359901	.0113183	-.01831772	-.0298761	-.0164179
40000.	-40.5478	-.126080	-.0918486	-.0530702	-.0226281	-.00730499	-.00342970
50000.	-40.2209	-.240437	-.148187	-.0719185	-.0283797	-.00898769	-8.65703E-04
60000.	-40.2314	-.246864	-.171734	-.0935161	-.0372433	-.00687580	.00536462
70000.	-40.3601	-.241292	-.189416	-.108326	-.0372244	.00170975	.0115451
80000.	-40.4485	-.303064	-.213746	-.0974470	-.0150843	.0124774	.00596267
100000.	-41.3985	-.220781	-.0969877	-.0320398	-.0136188	-.00923349	-.00545964

See appendix for Chebyshev fit details.

$He + H^+ \rightarrow H^+ + He(2^1S)$



Excitation Rate Coefficients for

$He + H^+ \rightarrow H^+ + He^*(2^1P)$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H Temp. (eV)	He Energy (eV/amu) 30000.	40000.	60000.	80000.	100000.	200000.	400000.
1.0E+00	8.28E-10	1.60E-09	2.86E-09	4.24E-09	5.97E-09	7.52E-09	7.65E-09
2.0E+00	8.28E-10	1.60E-09	2.86E-09	4.24E-09	5.96E-09	7.52E-09	7.65E-09
4.0E+00	8.29E-10	1.60E-09	2.86E-09	4.24E-09	5.96E-09	7.51E-09	7.65E-09
7.0E+00	8.29E-10	1.60E-09	2.86E-09	4.25E-09	5.96E-09	7.51E-09	7.65E-09
1.0E+01	8.30E-10	1.60E-09	2.86E-09	4.25E-09	5.95E-09	7.51E-09	7.65E-09
2.0E+01	8.32E-10	1.60E-09	2.86E-09	4.25E-09	5.94E-09	7.51E-09	7.65E-09
4.0E+01	8.36E-10	1.61E-09	2.86E-09	4.25E-09	5.93E-09	7.51E-09	7.65E-09
7.0E+01	8.41E-10	1.61E-09	2.87E-09	4.26E-09	5.92E-09	7.51E-09	7.65E-09
1.0E+02	8.47E-10	1.61E-09	2.87E-09	4.27E-09	5.91E-09	7.51E-09	7.65E-09
2.0E+02	8.60E-10	1.63E-09	2.89E-09	4.28E-09	5.89E-09	7.51E-09	7.65E-09
4.0E+02	8.80E-10	1.65E-09	2.92E-09	4.32E-09	5.87E-09	7.50E-09	7.65E-09
7.0E+02	9.14E-10	1.67E-09	2.96E-09	4.36E-09	5.84E-09	7.51E-09	7.65E-09
1.0E+03	9.51E-10	1.69E-09	2.99E-09	4.41E-09	5.82E-09	7.50E-09	7.65E-09
2.0E+03	1.07E-09	1.78E-09	3.12E-09	4.51E-09	5.77E-09	7.50E-09	7.65E-09
4.0E+03	1.30E-09	1.97E-09	3.33E-09	4.63E-09	5.73E-09	7.48E-09	7.65E-09
7.0E+03	1.63E-09	2.28E-09	3.58E-09	4.75E-09	5.70E-09	7.45E-09	7.64E-09
1.0E+04	1.93E-09	2.57E-09	3.78E-09	4.85E-09	5.69E-09	7.41E-09	7.63E-09
2.0E+04	2.79E-09	3.32E-09	4.29E-09	5.09E-09	5.74E-09	7.27E-09	7.60E-09

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
30000.	-41.2504	.486127	.299576	.128300	.0302620	-.00574350	-.00993827
40000.	-40.1898	.273759	.181714	.0905572	.0322161	.00429196	-.00603203
60000.	-39.1561	.162678	.0999502	.0424831	.00981070	-.00141938	-.00247916
80000.	-38.4598	.0783748	.0416546	.0129989	.00102165	-5.19012E-04	6.56879E-04
100000.	-37.9111	-.0245928	-.00529346	.00275039	.00343274	.00202306	8.64146E-04
200000.	-37.4242	-.0102227	-.00738113	-.00482593	-.00285436	-.00141545	-4.70357E-04
400000.	-37.3788	-.00169868	-.00156122	-.00125546	-8.18521E-04	-4.46724E-04	-1.48377E-04

See appendix for Chebyshev fit details.

$He + H^+ \rightarrow H^+ + He(2^1P)$



Excitation Cross Sections for

$H^+ + He \rightarrow H^+ + He^*(3^1S, 3^1P, 3^1D)$

$He^*(3^1S)$		$He^*(3^1P)$		$He^*(3^1D)$	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
5.0E+03	6.81E-20	2.0E+03	7.41E-21	3.5E+03	4.51E-20
6.0E+03	1.08E-19	4.0E+03	4.38E-20	4.0E+03	6.18E-20
7.0E+03	1.65E-19	7.0E+03	1.90E-19	7.0E+03	3.11E-19
8.0E+03	2.26E-19	1.0E+04	5.04E-19	1.0E+04	3.90E-19
9.0E+03	2.96E-19	1.5E+04	1.04E-18	1.5E+04	4.30E-19
1.0E+04	3.65E-19	2.0E+04	1.25E-18	2.0E+04	3.74E-19
1.5E+04	6.83E-19	2.7E+04	1.21E-18	2.8E+04	2.83E-19
2.0E+04	9.53E-19	4.0E+04	1.56E-18	4.0E+04	3.57E-19
3.0E+04	1.30E-18	7.0E+04	2.50E-18	5.5E+04	3.82E-19
4.0E+04	1.43E-18	1.0E+05	3.05E-18	7.0E+04	3.47E-19
5.0E+04	1.39E-18	1.3E+05	3.27E-18	1.0E+05	2.68E-19
6.0E+04	1.29E-18	2.0E+05	3.13E-18	1.5E+05	2.02E-19
7.0E+04	1.21E-18	4.0E+05	2.35E-18		
8.0E+04	1.12E-18	7.0E+05	1.67E-18		
9.0E+04	1.03E-18	1.0E+06	1.30E-18		
1.0E+05	9.63E-19				
1.5E+05	6.92E-19				
2.0E+05	5.07E-19				
2.5E+05	3.91E-19				

<u>References:</u> 680, 685, 686, 687, 688, 689, 690

<u>Accuracy:</u> 40%

<u>Notes:</u> None

Chebyshev Fitting Parameters for Cross Sections

$He^*(3^1S)$	E_{min}	=	5.0E+03	eV/amu,	E_{max}	=	2.5E+05 eV/amu
$He^*(3^1P)$	E_{min}	=	2.0E+03	eV/amu,	E_{max}	=	1.0E+06 eV/amu
$He^*(3^1D)$	E_{min}	=	3.5E+03	eV/amu,	E_{max}	=	1.5E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
$He^*(3^1S)$	-84.4433	.775561	-1.05948	.110960	.0219972	-.00305073	-.00996704
$He^*(3^1P)$	-84.5124	2.50529	-1.63211	.183239	-.0277337	-.0862483	.168352
$He^*(3^1D)$	-85.9726	.503209	-.748814	.283442	-.187874	-.0352298	.123942

The fit represents the $He^*(3^1S)$ cross section with an rms deviation of 1.6%. The maximum deviation is 3.9% at 6.0E+03 eV/amu.

The fit represents the $He^*(3^1P)$ cross section with an rms deviation of 9.8%. The maximum deviation is 18.7% at 2.7E+04 eV/amu.

The fit represents the $He^*(3^1D)$ cross section with an rms deviation of 6.2%. The maximum deviation is 13.9% at 2.8E+04 eV/amu.

See appendix for Chebyshev fit details.

$H^+ + He \rightarrow He(3^1S, 3^1P, 3^1D)$



Excitation Cross Sections for

$H^+ + He \rightarrow H^+ + He^*(4^1S, 4^1P, 4^1D)$

He*(4¹S)		He*(4¹P)		He*(4¹D)	
Energy (eV/amu)	σ (cm²)	Energy (eV/amu)	σ (cm²)	Energy (eV/amu)	σ (cm²)
1.0E+03	2.09E-22	1.0E+04	1.19E-19	2.0E+03	2.21E-21
2.0E+03	1.64E-21	1.5E+04	2.63E-19	4.0E+03	7.14E-21
3.0E+03	2.98E-21	2.0E+04	3.19E-19	7.0E+03	7.63E-20
4.0E+03	2.02E-21	3.0E+04	3.61E-19	1.0E+04	1.52E-19
4.5E+03	1.70E-21	4.0E+04	4.49E-19	1.3E+04	1.79E-19
7.0E+03	3.94E-20	5.0E+04	5.72E-19	2.0E+04	1.18E-19
1.0E+04	8.88E-20	6.0E+04	6.92E-19	2.6E+04	1.05E-19
1.5E+04	1.34E-19	7.0E+04	7.73E-19	4.0E+04	1.30E-19
2.0E+04	2.15E-19	8.0E+04	8.43E-19	5.0E+04	1.38E-19
4.0E+04	4.94E-19	9.0E+04	9.14E-19	7.0E+04	1.31E-19
4.4E+04	5.15E-19	1.0E+05	9.72E-19	1.0E+05	1.07E-19
7.0E+04	4.01E-19	1.3E+05	1.06E-18	1.5E+05	7.17E-20
1.0E+05	2.87E-19	1.5E+05	1.05E-18	2.0E+05	5.12E-20
1.5E+05	1.93E-19	2.0E+05	1.00E-18	4.0E+05	2.42E-20
2.0E+05	1.41E-19	3.0E+05	8.79E-19	4.0E+05	2.42E-20
4.0E+05	6.74E-20	4.0E+05	7.84E-19	7.0E+05	1.34E-20
7.0E+05	3.94E-20	5.0E+05	6.81E-19	1.0E+06	9.00E-21
1.0E+06	2.69E-20	6.0E+05	6.07E-19		
		7.0E+05	5.46E-19		
		8.0E+05	5.00E-19		
		9.0E+05	4.50E-19		
		1.0E+06	4.00E-19		

<u>References:</u> 680, 685, 686, 687, 688, 689, 690

<u>Accuracy:</u> 40%

<u>Notes:</u> None

<u>Chebyshev Fitting Parameters for Cross Sections</u>

He*(4¹S)	E_{min} =	1.0E+03	eV/amu,	E_{max} =	1.0E+06	eV/amu
He*(4¹P)	E_{min} =	1.0E+04	eV/amu,	E_{max} =	1.0E+06	eV/amu
He*(4¹D)	E_{min} =	2.0E+03	eV/amu,	E_{max} =	1.0E+06	eV/amu

	A0	A1	A2	A3	A4	A5	A6
He*(4¹S)	-90.7744	2.47036	-2.37315	-.112038	.549272	-.0385738	-.264616
He*(4¹P)	-84.4673	.568166	-.658688	-.0606377	.0615161	.0857392	-.0897306
He*(4¹D)	-90.3972	.468672	-1.92730	.452357	7.32888E-04	-.125659	.290353

The fit represents the He*(4¹S) cross section with an rms deviation of 60.6%. The maximum deviation is 217.9% at 4.5E+03 eV/amu.

The fit represents the He*(4¹P) cross section with an rms deviation of 2.4%. The maximum deviation is 4.9% at 3.0E+04 eV/amu.

The fit represents the He*(4¹D) cross section with an rms deviation of 25.2%. The maximum deviation is 50.7% at 2.6E+04 eV/amu.

See appendix for Chebyshev fit details.

$H^+ + He \rightarrow H^+ + He(4^1S, 4^1P, 4^1D)$



Total Vacuum Ultraviolet Emission Cross Sections for

$H^+ + He \rightarrow$ v.u.v. (λ = 20-140nm)

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
4.2E+03	9.00E+07	1.05E-18
5.0E+03	9.82E+07	1.25E-18
6.0E+03	1.08E+08	1.49E-18
7.0E+03	1.16E+08	1.70E-18
8.0E+03	1.24E+08	1.89E-18
9.0E+03	1.32E+08	2.05E-18
1.0E+04	1.39E+08	2.20E-18
1.5E+04	1.70E+08	2.73E-18
2.0E+04	1.96E+08	3.00E-18
3.0E+04	2.41E+08	3.21E-18
4.0E+04	2.78E+08	3.30E-18
5.0E+04	3.11E+08	3.33E-18
6.0E+04	3.40E+08	3.34E-18
7.0E+04	3.68E+08	3.32E-18
8.0E+04	3.93E+08	3.31E-18
9.0E+04	4.17E+08	3.28E-18
1.0E+05	4.39E+08	3.25E-18
1.5E+05	5.38E+08	3.14E-18

Reference: 654

Accuracy: 40%

Note: Values shown in the graph correspond to values of measured cross sections for v.u.v. emission which have been corrected for the photoelectric yield integrated over the wavelength range of sensitivity of the detector (approx. 20-140nm).

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 4.2E+03 eV/amu, E_{max} = 1.5E+05 eV/amu

A0	A1	A2	A3	A4	A5	A6
-81.1574	.502834	-.275390	.0508166	.00293296	-.00661360	.00122541

The fit represents the above cross section with an rms deviation of 0.3%. The maximum deviation is 0.5% at 3.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$H^+ + He \rightarrow$ v.u.v. (20–140 nm)



Excitation Cross Sections for

$He^+ + H \rightarrow H(n=2)$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
3.8E+03	8.56E+07	5.98E-17
4.0E+03	8.79E+07	6.01E-17
5.0E+03	9.82E+07	6.07E-17
6.0E+03	1.08E+08	6.13E-17
7.0E+03	1.16E+08	6.16E-17
8.0E+03	1.24E+08	6.19E-17
9.0E+03	1.32E+08	6.18E-17
1.0E+04	1.39E+08	6.17E-17
1.5E+04	1.70E+08	6.17E-17
2.0E+04	1.96E+08	6.27E-17
3.0E+04	2.41E+08	6.59E-17
4.0E+04	2.78E+08	7.52E-17
5.0E+04	3.11E+08	8.96E-17
5.2E+04	3.17E+08	9.30E-17

Reference: 716

Accuracy: 50%

Note: These data were obtained by measuring the energy loss of the He^+ ion after collision. The resolution of the energy-loss spectrometer was not sufficient to resolve the 2s and 2p states.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 3.8E+03 eV/amu, E_{max} = 5.2E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-74.4909	.160714	.0943278	.0580551	.0178172	.00276895	-4.72682E-04

The fit represents the above cross section with an rms deviation of 0.3%. The maximum deviation is 0.5% at 3.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + H \rightarrow He^+ + H(n=2)$



Excitation Cross Sections for

$He^+ + H \rightarrow He^+ + H(2s,2p)$

H(2s) Energy (eV/amu)	H(2s) Cross Section (cm^2)	H(2p) Energy (eV/amu)	H(2p) Cross Section (cm^2)
1.5E+03	1.04E-17	1.3E+02	9.03E-18
2.0E+03	6.59E-18	1.5E+02	1.36E-17
2.8E+03	4.60E-18	2.0E+02	2.18E-17
3.0E+03	4.61E-18	3.0E+02	2.95E-17
4.0E+03	7.95E-18	4.0E+02	3.22E-17
4.7E+03	1.12E-17	5.0E+02	3.39E-17
5.0E+03	1.11E-17	6.0E+02	3.51E-17
6.0E+03	5.82E-18	7.0E+02	3.60E-17
6.6E+03	4.51E-18	8.0E+02	3.66E-17
		9.0E+02	3.72E-17
		1.0E+03	3.75E-17
		1.5E+03	3.85E-17
		2.0E+03	3.90E-17
		3.0E+03	3.96E-17
		4.0E+03	3.98E-17
		5.0E+03	4.02E-17
		6.0E+03	4.04E-17
		7.0E+03	4.07E-17
		8.0E+03	4.07E-17
		9.0E+03	4.08E-17
		9.6E+03	4.08E-17

References: 701, 712, 715

Accuracy: 30%

Note: The data for the excitation of the 2p state are (within the experimental error) the same as the Lyman-α emission cross section.

Chebyshev Fitting Parameters for Cross Sections

H(2s) E_{min} = 1.5E+03 eV/amu, E_{max} = 6.6E+03 eV/amu
H(2p) E_{min} = 1.3E+02 eV/amu, E_{max} = 9.6E+03 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H(2s)	-78.9847	-.124567	.114965	-.358545	-.199857		
H(2p)	-76.1170	.522520	-.318078	.179576	-.0921414	.0455819	-.0220305

The fit represents the H(2s) cross section with an rms deviation of 10.1%. The maximum deviation is 25.2% at 6.0E+03 eV/amu.

The fit represents the H(2p) cross section with an rms deviation of 0.5%. The maximum deviation is 0.8% at 1.5E+03 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + H \rightarrow He^+ + H(2s,2p)$



Excitation Rate Coefficients for

$He^+ + H \rightarrow He^+ + H(2p)$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

He Temp. (eV)	Equal Temp.	H Temp. (eV) 100.	200.	500.	1000.	5000.	10000.	20000.
1.0E+02	3.27E-10*	3.27E-10*	6.34E-10	1.24E-09	1.88E-09	2.52E-09*	1.57E-09#	7.49E-10#
2.0E+02	6.98E-10	4.12E-10	6.98E-10	1.28E-09	1.90E-09	2.52E-09*	1.56E-09#	7.48E-10#
3.0E+02	9.74E-10	4.91E-10	7.59E-10	1.31E-09	1.93E-09	2.51E-09*	1.56E-09#	7.47E-10#
4.0E+02	1.20E-09	5.65E-10	8.17E-10	1.35E-09	1.96E-09	2.51E-09*	1.56E-09#	7.46E-10#
5.0E+02	1.39E-09	6.34E-10	8.72E-10	1.39E-09	1.98E-09	2.50E-09*	1.55E-09#	7.45E-10#
6.0E+02	1.56E-09	6.98E-10	9.24E-10	1.42E-09	2.01E-09	2.50E-09*	1.55E-09#	7.44E-10#
7.0E+02	1.71E-09	7.59E-10	9.74E-10	1.46E-09	2.03E-09	2.49E-09*	1.54E-09#	7.43E-10#
8.0E+02	1.85E-09	8.17E-10	1.02E-09	1.49E-09	2.05E-09	2.49E-09*	1.54E-09#	7.42E-10#
9.0E+02	1.98E-09	8.72E-10	1.07E-09	1.53E-09	2.08E-09	2.48E-09*	1.54E-09#	7.41E-10#
1.0E+03	2.10E-09	9.24E-10	1.11E-09	1.56E-09	2.10E-09	2.47E-09*	1.53E-09#	7.40E-10#
2.0E+03	2.76E-09*	1.35E-09	1.49E-09	1.85E-09	2.31E-09	2.42E-09*	1.50E-09#	7.29E-10#
4.0E+03	2.53E-09*	1.96E-09	2.05E-09	2.31E-09	2.61E-09	2.31E-09#	1.44E-09#	7.08E-10#
6.0E+03	1.99E-09#	2.38E-09	2.45E-09	2.61E-09	2.76E-09*	2.20E-09#	1.38E-09#	6.88E-10#
8.0E+03	1.57E-09#	2.65E-09	2.68E-09	2.76E-09*	2.81E-09*	2.09E-09#	1.3E-09#	6.69E-10#
1.0E+04	1.27E-09#	2.78E-09*	2.79E-09*	2.81E-09*	2.79E-09*	1.99E-09#	1.27E-09#	6.51E-10#
1.2E+04	1.04E-09#	2.82E-09*	2.81E-09*	2.79E-09*	2.73E-09*	1.90E-09#	1.22E-09#	6.34E-10#
1.5E+04	8.10E-10#	2.75E-09*	2.74E-09*	2.68E-09*	2.58E-09*	1.76E-09#	1.15E-09#	6.09E-10#
2.0E+04	5.71E-10#	2.51E-09*	2.49E-09*	2.42E-09*	2.31E-09#	1.57E-09#	1.04E-09#	5.71E-10#

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+02 eV, E_{max} = 2.0E+04 eV

H Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
100.	-41.3676	1.16527	-.103922	-.132163	-.0348422	-.0167981	-.00106673
200.	-40.8770	.821004	-.0109340	-.127376	-.0485804	-.0120056	7.63058E-04
500.	-40.2738	.442909	.00923768	-.0979147	-.0519441	-.0112276	.00362652
1000.	-39.8495	.187131	-.0287907	-.0768304	-.0399746	-.00851390	.00286700
5000.	-39.8387	-.195041	-.104148	-.0404533	-.0115750	-.00220721	-2.40196E-06
10000.	-40.7661	-.170203	-.0865943	-.0313220	-.00824330	-.00148203	-7.88573E-05
20000.	-42.1667	-.113121	-.0579411	-.0213073	-.00583215	-.00116246	-1.36085E-04
Equal Temp.	-41.3902	.353784	-.866154	-.123235	.00687538	.0519228	-.00558678

See appendix for Chebyshev fit details

$He^+ + H \rightarrow He^+ + H(2p)$

Maxwellian — Maxwellian



Spectral Line Emission Cross Sections for

$He^+ + H_2 \rightarrow H(Lyman\text{-}\alpha, Lyman\text{-}\beta)$

H(Lyman-α) Energy (eV/amu)	H(Lyman-α) Cross Section (cm^2)	H(Lyman-β) Energy (eV/amu)	H(Lyman-β) Cross Section (cm^2)
7.7E+00	4.33E-18	1.2E+01	1.06E-18
8.0E+00	5.16E-18	1.5E+01	2.15E-18
1.0E+01	1.05E-17	2.0E+01	3.68E-18
2.0E+01	3.39E-17	3.0E+01	5.82E-18
4.0E+01	5.51E-17	4.0E+01	7.07E-18
7.0E+01	6.19E-17	5.0E+01	7.81E-18
1.0E+02	6.01E-17	6.0E+01	8.21E-18
2.0E+02	4.57E-17	7.0E+01	8.57E-18
4.0E+02	3.56E-17	8.0E+01	8.64E-18
7.0E+02	3.20E-17	9.0E+01	8.56E-18
1.0E+03	3.13E-17	1.0E+02	8.46E-18
2.0E+03	3.35E-17	1.5E+02	7.67E-18
4.0E+03	4.45E-17	2.0E+02	6.78E-18
5.0E+03	4.61E-17		
7.0E+03	4.53E-17		
7.4E+03	4.48E-17		

References: 188, 195, 199, 700

Accuracy: 40%

Notes: (1) The Lyman-α transition is at 121.57nm and the Lyman-β transition is at 102.57nm. (2) Data on polarization of emissions may be found in Isler and Nathan (Ref. 199) and Young et al. (Ref. 700).

Chebyshev Fitting Parameters for Cross Sections

H(Lyman-α) E_{min} = 7.7E+00 eV/amu, E_{max} = 7.4E+03 eV/amu
H(Lyman-β) E_{min} = 1.2E+01 eV/amu, E_{max} = 2.0E+02 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H(Lyman-α)	-76.0522	.584543	-.527735	.587085	-.202654	-.0240480	-.0420581
H(Lyman-β)	-79.7592	.790251	-.524833	.114100	-.0482761	.0226572	-.00698738

The fit represents the H(Lyman-α) cross section with an rms deviation of 2.5%. The maximum deviation is 3.5% at 7.7E+00 eV/amu.

The fit represents the H(Lyman-β) cross section with an rms deviation of 0.5%. The maximum deviation is 1.0% at 6.0E+01 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + H_2 \rightarrow H(Lyman-\alpha, Lyman-\beta)$



Spectral Line Emission Cross Sections for

$He^+ + H_2 \rightarrow H(Balmer\text{-}\alpha, Balmer\text{-}\beta)$

H(Balmer-α) Energy (eV/amu)	H(Balmer-α) Cross Section (cm^2)	H(Balmer-β) Energy (eV/amu)	H(Balmer-β) Cross Section (cm^2)
1.0E+01	8.81E-19	1.5E+01	2.00E-19
2.0E+01	4.79E-18	2.0E+01	5.60E-19
4.0E+01	7.18E-18	4.0E+01	1.33E-18
7.0E+01	6.36E-18	5.0E+01	1.41E-18
1.0E+02	5.16E-18	7.0E+01	1.34E-18
2.0E+02	3.09E-18	1.0E+02	1.10E-18
4.0E+02	2.26E-18	2.0E+02	5.50E-19
7.0E+02	1.99E-18	4.0E+02	3.47E-19
1.0E+03	1.93E-18	7.0E+02	2.98E-19
2.0E+03	2.12E-18	1.0E+03	3.12E-19
4.0E+03	3.00E-18	2.0E+03	3.91E-19
7.0E+03	3.39E-18	4.0E+03	5.56E-19
1.0E+04	3.29E-18	5.0E+03	5.78E-19
		7.0E+03	5.28E-19
		8.0E+03	4.83E-19

<u>References:</u> 166, 199, 461, 464, 693, 694, 695

<u>Accuracy:</u> Unknown

<u>Notes:</u> (1) The Balmer-α emission is from the n=3->2 transition (656.28nm). (2) The Balmer-β emission is from the n=4->2 transition (486.13nm). (3) Data on polarization of emissions may be found in Isler and Nathan (Refs. 199 and 694).

Chebyshev Fitting Parameters for Cross Sections

H(Balmer-α) E_{min} = 1.0E+01 eV/amu, E_{max} = 1.0E+04 eV/amu
H(Balmer-β) E_{min} = 1.5E+01 eV/amu, E_{max} = 8.0E+03 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H(Balmer-α)	-80.7189	.0329782	-.173449	.608331	-.365301	.0179138	-.0168644
H(Balmer-β)	-84.1448	-.135846	-.0820835	.566238	-.436982	-.00523445	-.0131479

The fit represents the H(Balmer-α) cross section with an rms deviation of 2.2%. The maximum deviation is 4.5% at 2.0E+02 eV/amu.

The fit represents the H(Balmer-β) cross section with an rms deviation of 2.9%. The maximum deviation is 4.9% at 4.0E+01 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + H_2 \rightarrow H(\text{Balmer}-\alpha, \text{Balmer}-\beta)$



Dissociative Excitation Cross Sections for

$He^+ + H_2 \rightarrow He^+ + H(2s,2p) + H(\Sigma)$

H(2s) Energy (eV/amu)	H(2s) Cross Section (cm^2)	H(2p) Energy (eV/amu)	H(2p) Cross Section (cm^2)
1.4E+03	4.92E-18	1.3E+01	2.11E-17
1.5E+03	4.93E-18	2.0E+01	3.62E-17
1.7E+03	4.93E-18	4.0E+01	5.59E-17
2.0E+03	4.92E-18	6.0E+01	5.81E-17
2.5E+03	4.93E-18	7.0E+01	5.80E-17
3.0E+03	4.91E-18	1.0E+02	5.61E-17
3.5E+03	4.93E-18	2.0E+02	4.75E-17
4.0E+03	4.91E-18	4.0E+02	3.75E-17
4.5E+03	4.91E-18	7.0E+02	3.31E-17
5.0E+03	4.91E-18	1.0E+03	3.24E-17
5.5E+03	4.91E-18	2.0E+03	3.32E-17
6.0E+03	4.91E-18	4.0E+03	3.73E-17
6.4E+03	4.92E-18	5.0E+03	3.81E-17
		7.0E+03	3.75E-17
		7.4E+03	3.72E-17

References: 183, 195, 700, 701

Accuracy: Within a factor of 2

Notes: (1) The excitation cross sections for H(2s) and H(2p) are determined by the field-induced emission of Lyman-α (2s -> 2p -> 1s), and by Lyman-α emission (2p -> 1s), respectively. If cascade can be neglected, these are equal to the cross sections for the formation of the H(2s) and H(2p) states. (2) The notation H(Σ) designates all possible states for the other fragment (unspecified).

Chebyshev Fitting Parameters for Cross Sections

H(2s) E_{min} = 1.4E+03 eV/amu, E_{max} = 6.4E+03 eV/amu
H(2p) E_{min} = 1.3E+01 eV/amu, E_{max} = 7.4E+03 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H(2s)	-79.7066	-.00160282	1.58249E-05	7.60977E-04	2.44985E-04	6.18338E-04	
H(2p)	-75.6225	.00962025	-.178354	.293814	-.115895	-.0206130	-.00983973

The fit represents the H(2s) cross section with an rms deviation of 0.1%. The maximum deviation is 0.2% at 3.0E+03 eV/amu.

The fit represents the H(2p) cross section with an rms deviation of 0.8%. The maximum deviation is 0.9% at 7.0E+01 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + H_2 \rightarrow He^+ + H(2s,2p) + H(\Sigma)$



Dissociative Excitation Cross Sections for

$$He^+ + H_2 \rightarrow He^+ + H(3s,3p) + H(\Sigma)$$

H(3s) Energy (eV/amu)	H(3s) Cross Section (cm^2)	H(3p) Energy (eV/amu)	H(3p) Cross Section (cm^2)
2.5E+03	2.16E-19	2.5E+03	2.36E-18
3.0E+03	2.28E-19	3.0E+03	2.29E-18
3.5E+03	2.34E-19	3.5E+03	2.32E-18
4.0E+03	2.37E-19	4.0E+03	2.41E-18
4.5E+03	2.37E-19	4.5E+03	2.55E-18
5.0E+03	2.35E-19	5.0E+03	2.70E-18
5.5E+03	2.33E-19	5.5E+03	2.77E-18
6.0E+03	2.28E-19	6.0E+03	2.76E-18
7.0E+03	2.19E-19	7.0E+03	2.56E-18
		7.6E+03	2.29E-18

References: 183, 702

Accuracy: 50%

Notes: (1) The H(3s) and H(3p) cross sections are deduced from spectral-line emission measurements on the (unsubstantiated) assumption that cascade can be neglected. (2) The notation H(Σ) designates all possible states for the other fragment (unspecified).

Chebyshev Fitting Parameters for Cross Sections

H(3s) E_{min} = 2.5E+03 eV/amu, E_{max} = 7.0E+03 eV/amu
H(3p) E_{min} = 2.5E+03 eV/amu, E_{max} = 7.6E+03 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H(3s)	-85.8579	.00631068	-.0437007	-2.23538E-05	-1.27246E-04	5.58521E-04	6.28175E-04
H(3p)	-81.0948	.0512586	-.0390305	-.0692130	-.0155312	.00341300	-5.63877E-04

The fit represents the H(3s) cross section with an rms deviation of 0.1%. The maximum deviation is 0.2% at 5.0E+03 eV/amu.

The fit represents the H(3p) cross section with an rms deviation of 0.3%. The maximum deviation is 0.5% at 6.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + H_2 \rightarrow He^+ + H(3s,3p) + H(\Sigma)$



Projectile Excitation Cross Sections for

$He^+ + H_2 \rightarrow He^+(2s) + H_2$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.5E+03	6.95E+07	1.38E-20
3.0E+03	7.61E+07	2.18E-20
3.5E+03	8.22E+07	3.25E-20
4.0E+03	8.79E+07	4.58E-20
4.5E+03	9.32E+07	6.18E-20
5.0E+03	9.82E+07	8.10E-20
5.5E+03	1.03E+08	1.04E-19
6.0E+03	1.08E+08	1.30E-19
7.0E+03	1.16E+08	1.92E-19
7.6E+03	1.21E+08	2.31E-19

Reference: 706

Accuracy: Factor of 2

Notes: None

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 2.5E+03 eV/amu, E_{max} = 7.6E+03 eV/amu

A0	A1	A2	A3	A4	A5	A6
-88.6255	1.42095	-.00467445	-.00728965	-.00322271	-.00472755	-3.43794E-05

The fit represents the above cross section with an rms deviation of 0.1%. The maximum deviation is 0.1% at 6.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + H_2 \rightarrow He^+(2s) + H_2$



Excitation Cross Sections for

$He^+ + He \rightarrow He^+ + He^*(2^1S, 2^1P)$

$He^*(2^1S)$		$He^*(2^1P)$	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
7.0E+03	2.98E-18	6.0E+03	9.41E-18
8.0E+03	2.41E-18	7.0E+03	7.31E-18
9.0E+03	2.03E-18	1.0E+04	4.60E-18
1.0E+04	1.76E-18	1.5E+04	2.90E-18
1.5E+04	1.09E-18	2.0E+04	2.52E-18
2.0E+04	9.39E-19	4.0E+04	4.97E-18
2.5E+04	1.09E-18	7.0E+04	9.22E-18
3.0E+04	1.41E-18	1.0E+05	1.08E-17
		1.5E+05	1.13E-17
		2.5E+05	1.07E-17

References: 203, 684, 697, 699, 704, 705, 707, 717

Accuracy: 40%

Notes: (1) For relative cross sections see Refs. 697, 705, and 707. (2) For cross sections $He^+(1s) + He(1s^2)^1S \rightarrow He^+(1s) + He(1s2p)^{1,3}P$ see Bordenave-Montesquieu et al. (Ref. 704). (3) See Thomas (Ref. 684) for more details.

Chebyshev Fitting Parameters for Cross Sections

$He^*(2^1S)$	E_{min} =	7.0E+03	eV/amu,	E_{max} =	3.0E+04	eV/amu
$He^*(2^1P)$	E_{min} =	6.0E+03	eV/amu,	E_{max} =	2.5E+05	eV/amu

	A0	A1	A2	A3	A4	A5	A6
$He^*(2^1S)$	-82.0676	-.438515	.304992	.0789808	.0643720	-.0146204	-.00639777
$He^*(2^1P)$	-79.0520	.385720	.388654	-.432765	-.00474225	.126047	-.0106668

The fit represents the $He^*(2^1S)$ cross section with an rms deviation of 0.0%. The maximum deviation is 0.0% at 1.0E+04 eV/amu.

The fit represents the $He^*(2^1P)$ cross section with an rms deviation of 3.4%. The maximum deviation is 5.8% at 2.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + He \rightarrow He^+ + He(2^1S,2^1P)$



Excitation Cross Sections for

$He^+ + He \rightarrow He^+ + He^*(2^3S, 2^3P)$

$He^*(2^3S)$ Energy (eV/amu)	$He^*(2^3S)$ Cross Section (cm^2)	$He^*(2^3P)$ Energy (eV/amu)	$He^*(2^3P)$ Cross Section (cm^2)
6.4E+03	2.82E-18	6.5E+03	4.49E-18
7.0E+03	3.22E-18	7.0E+03	5.91E-18
8.0E+03	3.92E-18	8.0E+03	8.72E-18
9.0E+03	4.57E-18	8.4E+03	9.11E-18
1.0E+04	5.07E-18	9.0E+03	8.19E-18
1.5E+04	5.81E-18	1.0E+04	5.45E-18
2.0E+04	4.77E-18	1.5E+04	3.23E-18
2.5E+04	3.38E-18	2.0E+04	2.53E-18
3.0E+04	2.20E-18	2.5E+04	2.01E-18
		3.0E+04	1.18E-18
		3.2E+04	1.00E-18

<u>References:</u> 617, 684, 697, 699, 704, 708, 717

<u>Accuracy:</u> $E < 7.5 \times 10^3$ eV/amu - 75%
$E > 7.5 \times 10^3$ eV/amu - 40%

<u>Notes:</u> (1) For relative cross sections at lower energies see Bordenave-Montesquieu et al. (Ref. 704) and Dworetsky et al. (708). (2) For cross sections: $He^+(1s) + He^+(1s) \rightarrow He^+(1s) + He(1s2p)^{1,3}P$ see Ref. 704. (3) For more details see Thomas (Ref. 684).

<u>Chebyshev Fitting Parameters for Cross Sections</u>

$He^*(2^3S)$ E_{min} = 6.4E+03 eV/amu, E_{max} = 3.0E+04 eV/amu
$He^*(2^3P)$ E_{min} = 6.5E+03 eV/amu, E_{max} = 3.2E+04 eV/amu

	A0	A1	A2	A3	A4	A5	A6
$He^*(2^3S)$	-80.2166	-.0867600	-.430158	-.0323601	.00266805	-.00483153	.00172788
$He^*(2^3P)$	-80.5638	-.897275	-.226853	.0854789	-.228846	.0840160	.0168192

The fit represents the $He^*(2^3S)$ cross section with an rms deviation of 0.1%. The maximum deviation is 0.2% at 1.0E+04 eV/amu.

The fit represents the $He^*(2^3P)$ cross section with an rms deviation of 6.8%. The maximum deviation is 14.1% at 1.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + He \rightarrow He^+ + He(2^3S,2^3P)$



Excitation Cross Sections for

$He^+ + He \rightarrow He^+ + He^*(3^1S, 3^1P, 3^1D)$

He*(3¹S)		He*(3¹P)		He*(3¹D)	
Energy (eV/amu)	σ (cm²)	Energy (eV/amu)	σ (cm²)	Energy (eV/amu)	σ (cm²)
3.0E+03	2.96E-19	1.8E+03	4.84E-19	2.5E+03	3.98E-19
3.5E+03	2.78E-19	2.0E+03	5.24E-19	3.0E+03	5.16E-19
3.7E+03	2.75E-19	3.5E+03	6.61E-19	4.0E+03	8.25E-19
4.0E+03	2.93E-19	4.0E+03	6.50E-19	5.0E+03	1.02E-18
4.5E+03	3.33E-19	5.0E+03	6.28E-19	6.0E+03	1.04E-18
5.0E+03	3.64E-19	7.0E+03	8.19E-19	7.0E+03	1.03E-18
5.5E+03	3.83E-19	9.0E+03	9.18E-19	7.6E+03	1.02E-18
6.0E+03	3.98E-19	1.0E+04	9.04E-19		
7.0E+03	4.03E-19	1.5E+04	6.73E-19		
8.0E+03	4.02E-19	2.0E+04	5.62E-19		
		4.0E+04	1.49E-18		
		7.0E+04	2.34E-18		
		1.0E+05	2.81E-18		
		1.5E+05	3.07E-18		
		2.0E+05	3.14E-18		
		3.0E+05	2.90E-18		

References: 204, 696, 697, 698, 705, 708

Accuracy: 30%

Notes: (1) For relative cross sections at lower energies see Refs. 697, 705, and 708. (2) For more details see Ref. 684. (3) Acceptable polynomial fits were not possible for the He*(3¹P) data.

Chebyshev Fitting Parameters for Cross Sections

He*(3¹S) E_{min} = 3.0E+03 eV/amu, E_{max} = 8.0E+03 eV/amu
He*(3¹D) E_{min} = 2.5E+03 eV/amu, E_{max} = 7.6E+03 eV/amu

	A0	A1	A2	A3	A4	A5	A6
He*(3¹S)	-85.0346	.210040	-.00809700	-.0618979	.0267187	.00449587	-.0121214
He*(3¹D)	-83.4873	.491131	-.174973	-.0270949	.0308684	.00651934	-.00951236

The fit represents the He*(3¹S) cross section with an rms deviation of 0.6%. The maximum deviation is 1.2% at 3.7E+03 eV/amu.

No fit to He*(3¹P) data.

The fit represents the He*(3¹D) cross section with an rms deviation of 0.0%. The maximum deviation is 0.0% at 0.0E+00 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + He \rightarrow He^+ + He(3^1S, 3^1P, 3^1D)$



Excitation Cross Sections for

$He^+ + He \rightarrow He^+ + He^*(3^3S, 3^3P, 3^3D)$

$He^*(3^3S)$		$He^*(3^3P)$		$He^*(3^3D)$	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
2.5E+03	6.21E-19	2.5E+02	2.05E-19	2.5E+02	1.49E-18
3.0E+03	5.84E-19	3.2E+02	1.89E-19	3.7E+02	1.23E-18
3.5E+03	5.93E-19	4.0E+02	2.19E-19	4.0E+02	1.24E-18
4.0E+03	6.58E-19	7.0E+02	3.28E-19	6.0E+02	1.44E-18
4.5E+03	7.44E-19	7.4E+02	3.30E-19	7.0E+02	1.35E-18
5.0E+03	8.49E-19	1.0E+03	1.93E-19	1.0E+03	9.74E-19
5.5E+03	9.21E-19	1.1E+03	1.78E-19	1.5E+03	7.92E-19
6.0E+03	9.87E-19	1.5E+03	3.17E-19	2.0E+03	9.07E-19
7.0E+03	1.05E-18	2.0E+03	4.30E-19	3.0E+03	7.59E-19
7.7E+03	1.08E-18	3.7E+03	5.78E-19	4.0E+03	9.50E-19
		4.0E+03	5.65E-19	6.0E+03	1.47E-18
		5.0E+03	5.12E-19	7.0E+03	1.30E-18
		7.0E+03	6.98E-19	1.0E+04	8.73E-19
		1.0E+04	9.85E-19	2.0E+04	3.36E-19
		1.2E+04	1.07E-18	4.0E+04	1.35E-19
		2.0E+04	8.19E-19		
		4.0E+04	4.05E-19		

<u>References:</u> 204, 696, 697, 705, 707, 708

<u>Accuracy:</u> Unknown

<u>Notes:</u> (1) Relative cross sections may be found in Refs. 697, 705, 707, and 708. (2) Accuracies reported in Refs. 204 and 696 are: systematic error < 10%; random error < 5% on each; however, results differ by a factor of 2 at some energies common to both experiments. (3) Acceptable polynomial fits to the $He^*(3^3P)$ and $He^*(3^3D)$ data were not possible.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

$He^*(3^3S)$ E_{min} = 2.5E+03 eV/amu, E_{max} = 7.7E+03 eV/amu

	A0	A1	A2	A3	A4	A5	A6
$He^*(3^3S)$	-83.4063	.336147	.0620301	-.0763959	-.00323136	.0169475	-.00192645

The fit represents the $He^*(3^3S)$ cross section with an rms deviation of 0.3%. The maximum deviation is 0.5% at 4.5E+03 eV/amu.

No fit to $He^*(3^3P)$ data.

No fit to $He^*(3^3D)$ data.

See appendix for Chebyshev fit details.

$He^+ + He \rightarrow He^+ + He(3^3S, 3^3P, 3^3D)$



Excitation Cross Sections for

$He^+ + He \rightarrow He^+ + He^*(4^1S, 4^1P, 4^1D)$

$He^*(4^1S)$		$He^*(4^1P)$		$He^*(4^1D)$	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
5.0E+02	7.34E-21	3.0E+03	2.03E-19	2.5E+02	8.39E-20
5.8E+02	6.87E-21	3.5E+03	2.63E-19	3.0E+02	9.35E-20
7.0E+02	8.28E-21	4.0E+03	2.93E-19	3.5E+02	9.18E-20
1.0E+03	1.49E-20	4.5E+03	3.07E-19	4.0E+02	8.22E-20
1.2E+03	1.81E-20	5.0E+03	2.95E-19	5.0E+02	6.48E-20
1.5E+03	1.72E-20	5.5E+03	2.70E-19	6.0E+02	5.93E-20
2.0E+03	3.22E-20	6.0E+03	2.44E-19	7.0E+02	5.92E-20
2.7E+03	4.40E-20	6.6E+03	2.35E-19	1.0E+03	6.79E-20
3.5E+03	4.21E-20	7.0E+03	2.43E-19	1.5E+03	6.27E-20
4.0E+03	4.27E-20	8.0E+03	2.82E-19	2.0E+03	7.15E-20
5.0E+03	4.92E-20	9.0E+03	3.12E-19	3.0E+03	1.03E-19
6.0E+03	6.04E-20	1.0E+04	3.19E-19	4.0E+03	1.55E-19
7.0E+03	6.58E-20	1.5E+04	2.60E-19	7.0E+03	4.10E-19
1.0E+04	9.65E-20	2.0E+04	2.11E-19	8.5E+03	4.92E-19
1.3E+04	1.17E-19			1.0E+04	4.31E-19
2.0E+04	9.49E-20			1.5E+04	2.02E-19
4.0E+04	2.37E-19			2.0E+04	1.21E-19
7.0E+04	2.99E-19			2.3E+04	1.16E-19
1.0E+05	2.82E-19			3.0E+04	1.36E-19
2.0E+05	1.95E-19			4.0E+04	1.74E-19
3.0E+05	1.41E-19			5.5E+04	1.90E-19
				7.0E+04	1.82E-19
				1.0E+05	1.51E-19
				1.5E+05	1.10E-19
				2.0E+05	8.42E-20
				3.0E+05	5.94E-20

<u>References:</u> 204, 696, 698, 703, 705

<u>Accuracy:</u> Unknown

<u>Notes:</u> (1) Relative cross sections may be found in Ref. 705. (2) Accuracies reported in Refs. 204 and 696 are: systematic error < 10%; random error < 5% on each; however, results differ by a factor of 2 at some energies common to both experiments. (3) Acceptable polynomial fits to the $He^*(4^1D)$ data were not possible.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

$He^*(4^1S)$ E_{min} = 5.0E+02 eV/amu, E_{max} = 3.0E+05 eV/amu
$He^*(4^1P)$ E_{min} = 3.0E+03 eV/amu, E_{max} = 2.0E+04 eV/amu

	A0	A1	A2	A3	A4	A5	A6
$He^*(4^1S)$	-88.4744	1.72074	-.628284	-.185014	-.137545	-.0287794	.0855418
$He^*(4^1P)$	-85.6466	-.0488093	-.204458	-.0516457	-.0853529	.120960	.0910818

The fit represents the $He^*(4^1S)$ cross section with an rms deviation of 13.5%. The maximum deviation is 38.5% at 2.0E+04 eV/amu.

The fit represents the $He^*(4^1P)$ cross section with an rms deviation of 3.0%. The maximum deviation is 5.1% at 1.0E+04 eV/amu.

No fit to $He^*(4^1D)$ data.

See appendix for Chebyshev fit details.

$He^+ + He \rightarrow He^+ + He(4^1S, 4^1P, 4^1D)$



Excitation Cross Sections for

$He^+ + He \rightarrow He^+ + He^*(4^3S, 4^3P, 4^3D)$

$He^*(4^3S)$		$He^*(4^3P)$		$He^*(4^3D)$	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
1.2E+03	2.24E-20	2.5E+02	2.15E-20	2.5E+02	2.91E-19
1.8E+03	4.12E-20	3.3E+02	2.30E-20	3.6E+02	3.48E-19
2.0E+03	3.93E-20	4.0E+02	2.13E-20	4.0E+02	3.37E-19
2.4E+03	3.38E-20	4.7E+02	1.96E-20	6.0E+02	1.96E-19
3.6E+03	7.03E-20	7.0E+02	2.97E-20	7.0E+02	1.82E-19
4.0E+03	6.50E-20	1.0E+03	2.06E-20	9.5E+02	2.06E-19
5.2E+03	4.84E-20	1.5E+03	2.63E-20	1.0E+03	2.05E-19
7.0E+03	6.87E-20	2.0E+03	3.87E-20	1.5E+03	1.59E-19
1.0E+04	1.28E-19	4.0E+03	1.15E-19	2.0E+03	1.36E-19
1.5E+04	1.96E-19	4.5E+03	1.17E-19	2.2E+03	1.33E-19
2.0E+04	2.54E-19	5.5E+03	1.13E-19	3.0E+03	1.53E-19
2.6E+04	2.25E-19	7.0E+03	1.24E-19	3.7E+03	1.50E-19
		1.0E+04	1.84E-19	4.0E+03	1.54E-19
		1.5E+04	2.60E-19	7.0E+03	3.55E-19
		2.0E+04	2.06E-19	8.0E+03	3.73E-19
		2.7E+04	1.41E-19	1.0E+04	3.40E-19
		3.2E+04	1.70E-19	2.0E+04	1.57E-19
		3.7E+04	1.41E-19	3.5E+04	6.48E-20

References: 204, 696, 703, 705, 708

Accuracy: Unknown

Notes: (1) Relative cross sections may be found in Refs. 705 and 708. (2) Accuracies reported in Refs. 204 and 696 are: systematic error < 10%; random error < 5% on each; however, results differ by a factor of 2 at some energies common to both experiments. (3) Acceptable polynomial fits to these data were not possible.

Chebyshev Fitting Parameters for Cross Sections

No fits to the cross sections.

See appendix for Chebyshev fit details.

$He^+ + He \rightarrow He^+ + He(4^3S, 4^3P, 4^3D)$



Projectile Excitation Cross Sections for

$He^+ + He \rightarrow He^+(n=2) + He$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.0E+03	4.39E+07	3.84E-18
1.7E+03	5.73E+07	8.16E-18
2.0E+03	6.21E+07	7.62E-18
3.0E+03	7.61E+07	5.30E-18
4.0E+03	8.79E+07	4.35E-18
5.0E+03	9.82E+07	3.80E-18
6.0E+03	1.08E+08	3.43E-18
7.0E+03	1.16E+08	3.16E-18
8.0E+03	1.24E+08	2.96E-18
9.0E+03	1.32E+08	2.80E-18
1.0E+04	1.39E+08	2.68E-18
2.0E+04	1.96E+08	2.30E-18
2.5E+04	2.20E+08	2.11E-18
3.0E+04	2.41E+08	1.91E-18
4.0E+04	2.78E+08	1.55E-18

References: 699, 704, 705, 713, 714

Accuracy: E < 70%

Notes: (1) See Ref. 705 for relative cross sections for excitation of He^+ 2^2P. (2) For combined excitation of target and projectile see Ref. 714.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.0E+03 eV/amu, E_{max} = 4.0E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-80.4224	-.698320	-.116751	.174729	-.194669	.0663496	-.0346561

The fit represents the above cross section with an rms deviation of 1.4%. The maximum deviation is 2.7% at 3.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + He \rightarrow He^+(n=2) + He$



Projectile Excitation Cross Sections for

$He^+ + He \rightarrow He^+(n=3) + He$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
7.4E+03	1.19E+08	2.93E-19
8.0E+03	1.24E+08	4.13E-19
9.0E+03	1.32E+08	6.76E-19
1.0E+04	1.39E+08	8.88E-19
1.5E+04	1.70E+08	1.66E-18
2.0E+04	1.96E+08	2.43E-18
2.5E+04	2.20E+08	3.54E-18

References: 705, 713

Accuracy: Unknown

Note: Ref. 705 gives relative cross sections for excitation of 3^2P level.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 7.4E+03 eV/amu, E_{max} = 2.5E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-82.4466	1.12862	-.190981	.124483	-.0202534		

The fit represents the above cross section with an rms deviation of 1.4%. The maximum deviation is 2.9% at 8.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + He \rightarrow He^+(n=3) + He$



Spectral Line Emission Cross Sections for

He + H_2 -> H(Balmer-α,Balmer-β)

H(Balmer-α) Energy (eV/amu)	H(Balmer-α) Cross Section (cm^2)	H(Balmer-β) Energy (eV/amu)	H(Balmer-β) Cross Section (cm^2)
1.5E+02	6.77E-19	1.2E+02	6.93E-20
2.0E+02	7.82E-19	1.5E+02	1.34E-19
3.0E+02	9.61E-19	2.0E+02	2.18E-19
4.0E+02	1.08E-18	3.0E+02	3.34E-19
5.0E+02	1.18E-18	4.0E+02	4.11E-19
6.0E+02	1.26E-18	5.0E+02	4.56E-19
7.0E+02	1.34E-18	6.0E+02	4.85E-19
8.0E+02	1.40E-18	7.0E+02	5.07E-19
9.0E+02	1.45E-18	8.0E+02	5.25E-19
1.0E+03	1.51E-18	9.0E+02	5.35E-19
2.0E+03	1.80E-18	1.0E+03	5.44E-19
3.0E+03	1.89E-18	1.3E+03	5.52E-19
4.0E+03	1.89E-18	2.0E+03	5.47E-19
5.0E+03	1.84E-18	3.0E+03	5.24E-19
6.0E+03	1.76E-18	4.0E+03	4.97E-19
7.0E+03	1.65E-18	5.0E+03	4.75E-19
7.6E+03	1.58E-18	6.0E+03	4.54E-19
		7.0E+03	4.36E-19
		8.0E+03	4.20E-19

Reference: 461

Accuracy: Unknown

Notes: (1) The hydrogen Balmer line emission results from dissociative excitation of H_2. (2) The Balmer-α emission is the n=3->2 transition (656.28nm). (3) The Balmer-β emission is the n=4->2 transition (486.13nm).

Chebyshev Fitting Parameters for Cross Sections

H(Balmer-α) E_{min} = 1.5E+02 eV/amu, E_{max} = 7.6E+03 eV/amu
H(Balmer-β) E_{min} = 1.2E+02 eV/amu, E_{max} = 8.0E+03 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H(Balmer-α)	-82.3960	.472324	-.197315	-.0438413	-.0176366	-.00308248	5.34250E-05
H(Balmer-β)	-85.1264	.678943	-.561863	.182560	-.0716643	.0321037	-.0158573

The fit represents the H(Balmer-α) cross section with an rms deviation of 0.3%. The maximum deviation is 0.5% at 2.0E+02 eV/amu.

The fit represents the H(Balmer-β) cross section with an rms deviation of 0.9%. The maximum deviation is 1.4% at 2.0E+02 eV/amu.

See appendix for Chebyshev fit details.

He + H_2 -> H(Balmer-α, Balmer-β)



Excitation Cross Sections for

He + He -> He(2^1S,2^1P)

He(2^1S) Energy (eV/amu)	He(2^1S) Cross Section (cm^2)	He(2^1P) Energy (eV/amu)	He(2^1P) Cross Section (cm^2)
1.6E+01	1.00E-19	1.3E+01	4.83E-18
2.0E+01	6.25E-19	1.5E+01	7.85E-18
2.5E+01	1.17E-18	2.0E+01	1.52E-17
3.0E+01	1.52E-18	2.5E+01	2.39E-17
3.5E+01	1.73E-18	3.0E+01	3.03E-17
4.0E+01	1.90E-18	3.5E+01	3.59E-17
4.5E+01	2.07E-18	4.0E+01	4.05E-17
5.0E+01	2.20E-18	5.0E+01	4.70E-17
5.5E+01	2.33E-18	6.0E+01	5.25E-17
6.0E+01	2.44E-18	7.0E+01	5.62E-17
7.0E+01	2.65E-18	8.0E+01	5.95E-17
7.4E+01	2.72E-18	9.0E+01	6.19E-17
		1.0E+02	6.39E-17
		1.5E+02	7.24E-17

References: 719, 721, 722

Accuracy: Unknown

Notes: (1) The data for 2^1S state are theoretical data of Olson et al. (Ref. 721). The 2^1P state data are the experimental data of Kempter et al. (Ref. 719) and are approximately a factor of 2 greater than those of Olson et al. (2) The experimental data were determined by measuring emission line intensities and correcting for branching ratios using known transition probabilities. With He, cascading contributions should be less than 3%.

Chebyshev Fitting Parameters for Cross Sections

He(2^1S) E_{min} = 1.6E+01 eV/amu, E_{max} = 7.4E+01 eV/amu
He(2^1P) E_{min} = 1.3E+01 eV/amu, E_{max} = 1.5E+02 eV/amu

	A0	A1	A2	A3	A4	A5	A6
He(2^1S)	-82.7427	1.30331	-.579333	.299739	-.133372	.0473667	-.0142447
He(2^1P)	-76.1505	1.23531	-.420131	.116081	-.0158137	-.00167796	-.0033629

The fit represents the He(2^1S) cross section with an rms deviation of 0.5%. The maximum deviation is 0.9% at 2.5E+01 eV/amu.

The fit represents the He(2^1P) cross section with an rms deviation of 1.2%. The maximum deviation is 3.0% at 2.0E+01 eV/amu.

See appendix for Chebyshev fit details.

He + He –> He(2^1S,2^1P)



Excitation Cross Section for

He + He -> He(3^1P,3^1D,3^3P,3^3D)

He(3^1P)		He(3^1D)		He(3^3P)		He(3^3D)	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
2.5E+02	2.27E-18	2.5E+01	1.12E-19	2.5E+01	1.49E-19	2.5E+01	1.67E-19
3.0E+02	2.12E-18	3.0E+01	2.23E-19	4.0E+01	9.97E-19	3.0E+01	4.68E-19
4.0E+02	1.93E-18	4.0E+01	5.21E-19	7.0E+01	2.45E-18	4.0E+01	1.09E-18
5.0E+02	1.83E-18	5.0E+01	9.01E-19	1.0E+02	2.68E-18	5.0E+01	1.55E-18
6.0E+02	1.77E-18	6.0E+01	1.33E-18	2.0E+02	2.20E-18	6.0E+01	1.98E-18
7.0E+02	1.76E-18	7.0E+01	1.71E-18	4.0E+02	1.58E-18	7.0E+01	2.39E-18
8.0E+02	1.76E-18	8.0E+01	1.92E-18	7.0E+02	1.35E-18	8.0E+01	2.68E-18
9.0E+02	1.76E-18	9.0E+01	1.81E-18	1.0E+03	1.47E-18	9.0E+01	2.80E-18
1.0E+03	1.77E-18	1.0E+02	1.74E-18	1.2E+03	1.64E-18	1.0E+02	2.77E-18
1.2E+03	1.80E-18	1.1E+02	1.74E-18			1.5E+02	2.44E-18
		1.5E+02	2.31E-18				

References: 718, 719, 723

Accuracy: Unknown

Note: The recommended data for 3^3P state have been interpolated between the data of Kimura et al. (Ref. 718) (E > 2.5x10^2 eV/amu) and the low-energy data of Kempter et al. (Ref. 719). This extrapolation is uncertain due to different energy dependencies.

Chebyshev Fitting Parameters for Cross Sections

He(3^1P)	E_{min}	=	2.5E+02	eV/amu,	E_{max}	=	1.2E+03 eV/amu
He(3^1D)	E_{min}	=	2.5E+01	eV/amu,	E_{max}	=	1.5E+02 eV/amu
He(3^3P)	E_{min}	=	2.5E+01	eV/amu,	E_{max}	=	1.2E+03 eV/amu
He(3^3D)	E_{min}	=	2.5E+01	eV/amu,	E_{max}	=	1.5E+02 eV/amu

	A0	A1	A2	A3	A4	A5	A6
He(3^1P)	-81.5965	-.116911	.0573167	8.08164E-04	-.00364047	6.11763E-05	.00187028
He(3^1D)	-83.4302	1.37998	-.506774	.0626383	.0986308	.0721865	3.46628E-04
He(3^3P)	-82.5431	.625566	-.782070	.545218	-.100346	.0310014	.00459533
He(3^3D)	-82.5242	1.15604	-.602651	.117602	-.0538056	.0673126	.0230997

The fit represents the He(3^1P) cross section with an rms deviation of 0.1%. The maximum deviation is 0.3% at 6.0E+02 eV/amu.

The fit represents the He(3^1D) cross section with an rms deviation of 1.9%. The maximum deviation is 2.8% at 9.0E+01 eV/amu.

The fit represents the He(3^3P) cross section with an rms deviation of 0.7%. The maximum deviation is 1.1% at 1.0E+02 eV/amu.

The fit represents the He(3^3D) cross section with an rms deviation of 0.1%. The maximum deviation is 0.3% at 9.0E+01 eV/amu.

See appendix for Chebyshev fit details.

He + He -> He(3^1P,3^1D,3^3P,3^3D)



Excitation Cross Sections for

He + He -> He(4^1S,4^1P)

He(4^1S)		He(4^1S) cont.		He(4^1P)	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
1.7E+01	2.00E-20	5.0E+03	2.96E-19	5.0E+03	3.65E-19
2.0E+01	2.98E-20	6.0E+03	3.37E-19	5.5E+03	3.88E-19
3.0E+01	9.53E-20	7.0E+03	3.78E-19	6.0E+03	4.06E-19
4.0E+01	2.37E-19	8.0E+03	4.11E-19	7.0E+03	4.23E-19
5.0E+01	4.71E-19	9.0E+03	4.40E-19	8.0E+03	4.06E-19
6.0E+01	7.06E-19	1.0E+04	4.61E-19	9.0E+03	3.85E-19
7.0E+01	8.91E-19	1.5E+04	5.11E-19	1.0E+04	3.60E-19
7.9E+01	1.05E-18	2.0E+04	4.72E-19	1.5E+04	2.61E-19
9.0E+01	1.19E-18	2.5E+04	4.22E-19	2.0E+04	1.88E-19
1.0E+02	1.33E-18			2.4E+04	1.50E-19
1.5E+02	1.90E-18				
1.8E+02	2.15E-18				

<u>References:</u> 703, 719, 723

<u>Accuracy:</u> Unknown

<u>Note:</u> The 4^1S data in the low and high energy ranges are plotted individually to indicate the problems involved in comparing and interpolating between low and high energy data. Use these data with caution.

Chebyshev Fitting Parameters for Cross Sections

He(4^1S)	E_{min}	=	1.7E+01	eV/amu,	E_{max}	=	1.8E+02 eV/amu
He(4^1S)	E_{min}	=	5.0E+03	eV/amu,	E_{max}	=	2.5E+04 eV/amu
He(4^1P)	E_{min}	=	5.0E+03	eV/amu,	E_{max}	=	2.4E+04 eV/amu

	A0	A1	A2	A3	A4	A5	A6
He(4^1S)	-85.2188	2.41501	-.487597	-.0727179	.111514	-.00381501	-.0352846
He(4^1S)	-84.6731	.196616	-.160691	-.0239012	.00732614	.00447306	.00338388
He(4^1D)	-85.4091	-.479796	-.191704	.0341683	-.0126641	.00116120	.00977808

The fit represents the He(4^1S) cross section with an rms deviation of 1.2%. The maximum deviation is 2.3% at 4.0E+01 eV/amu.

The fit represents the He(4^1S) cross section with an rms deviation of 0.2%. The maximum deviation is 0.4% at 7.1E+03 eV/amu.

The fit represents the He(4^1P) cross section with an rms deviation of 0.3%. The maximum deviation is 0.6% at 8.0E+03 eV/amu.

See appendix for Chebyshev fit details.

He + He -> He(4 ^{1}S,4 ^{1}P)



Excitation Cross Sections for

He + He -> He(4^3S,4^3P)

He(4^3S)		He(4^3S) cont.		He(4^3P)	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
2.4E+01	2.53E-20	5.0E+03	4.37E-19	2.5E+02	2.78E-18
2.5E+01	2.90E-20	5.5E+03	4.47E-19	4.0E+02	1.38E-18
3.0E+01	5.97E-20	6.0E+03	4.66E-19	7.0E+02	8.79E-19
3.5E+01	1.07E-19	7.0E+03	4.90E-19	1.0E+03	8.23E-19
4.0E+01	1.59E-19	8.0E+03	5.29E-19	2.0E+03	7.68E-19
4.5E+01	2.08E-19	9.0E+03	5.47E-19	4.0E+03	7.14E-19
5.0E+01	2.62E-19	1.0E+04	5.14E-19	7.0E+03	7.07E-19
5.5E+01	3.15E-19	1.1E+04	4.88E-19	1.0E+04	5.62E-19
6.0E+01	3.65E-19	1.5E+04	5.30E-19	1.5E+04	3.11E-19
7.0E+01	4.61E-19	2.0E+04	6.12E-19	2.0E+04	1.95E-19
8.0E+01	5.43E-19	2.5E+04	6.77E-19		
8.8E+01	5.95E-19				

<u>References:</u> 703, 718, 719, 720, 723

<u>Accuracy:</u> Unknown

<u>Note:</u> No attempt has been made to interpolate between the low and high energy data for the 4^3S state. Use these data with caution.

Chebyshev Fitting Parameters for Cross Sections

He(4^3S)	E_{min}	=	2.4E+01	eV/amu,	E_{max}	=	8.8E+01 eV/amu
He(4^3S)	E_{min}	=	5.0E+03	eV/amu,	E_{max}	=	2.5E+04 eV/amu
He(4^3P)	E_{min}	=	2.5E+02	eV/amu,	E_{max}	=	2.0E+04 eV/amu

	A0	A1	A2	A3	A4	A5	A6
He(4^3S)	-86.5334	1.57661	-.301332	.0231287	.0147011	-.0218801	.0103488
He(4^3S)	-84.1641	.186801	.0274621	.0565092	-.00857002	-.0283410	.0102453
He(4^3P)	-83.4554	-.954640	-.0267074	-.366176	-.0251284	-.0183797	.0210686

The fit represents the He(4^3S) cross section with an rms deviation of 0.4%. The maximum deviation is 1.0% at 4.5E+01 eV/amu.

The fit represents the He(4^3S) cross section with an rms deviation of 2.1%. The maximum deviation is 3.6% at 1.1E+04 eV/amu.

The fit represents the He(4^3P) cross section with an rms deviation of 3.5%. The maximum deviation is 6.2% at 4.0E+03 eV/amu.

See appendix for Chebyshev fit details.

He + He -> He(4^3S,4^3P)



Excitation and Ion-Pair Production Cross Sections for He + He -> Autoionizing States

$2s^2\ ^1S$		$2s2p\ ^3P$		$2p^2\ ^1D$		$2s2p\ ^1P$		$He^-(1s2s^2)^2S$	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
2.5E+03	6.83E-20	2.5E+03	1.72E-20	2.5E+03	2.01E-19	2.5E+03	1.13E-19	3.7E+03	2.45E-19
3.0E+03	8.84E-20	3.0E+03	2.04E-20	3.0E+03	2.55E-19	3.0E+03	1.52E-19	4.0E+03	2.60E-19
4.0E+03	1.32E-19	4.0E+03	2.70E-20	4.0E+03	3.57E-19	4.0E+03	2.27E-19	5.0E+03	3.05E-19
5.0E+03	1.80E-19	5.0E+03	3.37E-20	5.0E+03	4.42E-19	5.0E+03	2.91E-19	6.0E+03	3.43E-19
6.0E+03	2.28E-19	6.0E+03	4.01E-20	6.0E+03	5.18E-19	6.0E+03	3.48E-19	7.0E+03	3.76E-19
7.0E+03	2.82E-19	7.0E+03	4.74E-20	7.0E+03	5.74E-19	7.0E+03	4.05E-19	7.7E+03	3.83E-19
8.0E+03	3.37E-19	8.0E+03	5.42E-20	8.0E+03	6.16E-19	8.0E+03	4.53E-19	8.0E+03	3.80E-19
9.0E+03	3.90E-19	9.0E+03	6.16E-20	9.0E+03	6.49E-19	9.0E+03	5.05E-19	9.0E+03	3.41E-19
1.0E+04	4.33E-19	1.0E+04	6.96E-20	1.0E+04	6.72E-19	1.0E+04	5.49E-19	1.0E+04	2.80E-19
1.4E+04	5.20E-19	1.5E+04	1.30E-19	1.2E+04	6.89E-19	1.3E+04	6.18E-19	1.5E+04	8.38E-20
1.5E+04	5.19E-19	2.0E+04	2.51E-19	1.5E+04	5.58E-19	1.5E+04	5.92E-19	2.0E+04	3.76E-20
2.0E+04	4.54E-19	3.0E+04	4.28E-19	2.0E+04	3.42E-19	2.0E+04	4.82E-19	3.0E+04	2.22E-20
3.0E+04	3.31E-19	3.5E+04	4.92E-19	3.0E+04	1.57E-19	3.0E+04	3.38E-19	3.5E+04	1.99E-20
3.5E+04	2.86E-19			3.5E+04	1.22E-19	3.5E+04	2.89E-19		

<u>References:</u> 724, 725, 726

<u>Accuracy:</u> Unknown

<u>Note:</u> The $1s2s^2\ ^2S$ state is the autodetaching state of He^-.

Chebyshev Fitting Parameters for Cross Sections

$2s^2\ ^1S$	E_{min} =	2.5E+03	eV/amu,	E_{max} =	3.5E+04	eV/amu
$2s2p\ ^3P$	E_{min} =	2.5E+03	eV/amu,	E_{max} =	3.5E+04	eV/amu
$2p^2\ ^1D$	E_{min} =	2.5E+03	eV/amu,	E_{max} =	3.5E+04	eV/amu
$2s2p\ ^1P$	E_{min} =	2.5E+03	eV/amu,	E_{max} =	3.5E+04	eV/amu
$He^-(1s2s^2)^2S$	E_{min} =	3.7E+03	eV/amu,	E_{max} =	3.5E+04	eV/amu

	A0	A1	A2	A3	A4	A5	A6
$2s^2\ ^1S$	-85.8524	.799247	-.526341	-.114359	.0439092	.0371210	-.00499344
$2s2p\ ^3P$	-87.8698	1.74012	.165859	-.0382066	-.0794670	-.0352277	.0132924
$2p^2\ ^1D$	-85.2799	-.217724	-.729656	-.0810841	.0662609	.0581753	.00536077
$2s2p\ ^1P$	-85.2994	.497649	-.530222	-.0630468	.0249202	.0444313	-5.43766E-04
$He^-(1s2s^2)^2S$	-87.4403	-1.52170	-.498463	.350749	.157944	-.0929551	-.0455722

The fit represents the $2s^2\ ^1S$ cross section with an rms deviation of 1.1%. The maximum deviation is 2.2% at 2.0E+04 eV/amu.

The fit represents the $2s2p\ ^3P$ cross section with an rms deviation of 2.4%. The maximum deviation is 5.2% at 1.5E+04 eV/amu.

The fit represents the $2p^2\ ^1D$ cross section with an rms deviation of 2.0%. The maximum deviation is 3.6% at 2.0E+04 eV/amu.

The fit represents the $2s2p\ ^1P$ cross section with an rms deviation of 1.8%. The maximum deviation is 3.7% at 2.0E+04 eV/amu.

The fit represents the $He^-(1s2s^2)^2S$ cross section with an rms deviation of 1.9%. The maximum deviation is 3.4% at 1.5E+04 eV/amu.

See appendix for Chebyshev fit details.

He + He –> He(2s2p^{2}P, 2s2p^{1}P, 2s^{21}S, 2p^{21}D)
He + He –> He$^-$(1s2s^{22}S)



Excitation Cross Sections for

He^{2+} + H -> H(2s,2p)

H(2s)		H(2p)	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
2.5E+04	1.41E-16	2.5E+04	1.68E-16
3.0E+04	1.16E-16	3.0E+04	1.75E-16
4.0E+04	8.70E-17	4.0E+04	1.84E-16
5.0E+04	6.97E-17	5.0E+04	1.92E-16
6.0E+04	5.77E-17	6.0E+04	1.97E-16
7.0E+04	4.98E-17	7.0E+04	2.01E-16
8.0E+04	4.37E-17	8.0E+04	2.05E-16
9.0E+04	3.86E-17	9.0E+04	2.06E-16
1.0E+05	3.50E-17	1.0E+05	2.08E-16
1.5E+05	2.35E-17	1.5E+05	2.07E-16
2.0E+05	1.79E-17	2.0E+05	1.97E-16
3.0E+05	1.22E-17	3.0E+05	1.69E-16
4.0E+05	9.42E-18	4.0E+05	1.45E-16
5.0E+05	7.75E-18	5.0E+05	1.31E-16
5.6E+05	7.06E-18	5.4E+05	1.29E-16

Reference: 174

Accuracy: Unknown

Note: The curves are results of theoretical calculations for the excitation of H(1s) by He^{2+} inpact. Ref. 174 compares the results of calculations using alternative basis sets together with a full explanation of the estimated accuracy at different He^{2+} energies.

Chebyshev Fitting Parameters for Cross Sections

H(2s) E_{min} = 2.5E+04 eV/amu, E_{max} = 5.6E+05 eV/amu
H(2p) E_{min} = 2.5E+04 eV/amu, E_{max} = 5.4E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H(2s)	-76.0595	-1.50152	.0300296	.00522842	.00296312	-4.05387E	04.00159617
H(2p)	-72.5902	-.122617	-.180664	-.0249549	.0157733	.0139456	.00499314

The fit represents H(2s) cross section with an rms deviation of 0.2%.
The maximum deviation is 0.5% at 6.0E+04 eV/amu.

The fit represents H(2p) cross section with an rms deviation of 0.2%.
The maximum deviation is 0.3% at 4.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$He^{2+} + H \rightarrow H(2s,2p)$



Projectile Excitation Cross Sections for

$He(2^3S) + He(1^1S) \rightarrow He(3^1P) + He$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
5.0E+01	9.82E+06	3.44E-18
6.0E+01	1.08E+07	4.55E-18
7.0E+01	1.16E+07	5.27E-18
8.0E+01	1.24E+07	5.66E-18
9.0E+01	1.32E+07	5.84E-18
1.0E+02	1.39E+07	5.93E-18
1.5E+02	1.70E+07	4.63E-18
2.0E+02	1.96E+07	3.94E-18
2.5E+02	2.20E+07	4.44E-18
3.0E+02	2.41E+07	5.57E-18
4.0E+02	2.78E+07	6.82E-18
5.0E+02	3.11E+07	6.97E-18
6.0E+02	3.40E+07	6.76E-18
7.0E+02	3.68E+07	6.35E-18
8.0E+02	3.93E+07	5.88E-18
9.0E+02	4.17E+07	5.35E-18
1.0E+03	4.39E+07	4.80E-18
1.2E+03	4.81E+07	3.50E-18

References: 728, 729

Accuracy: Unknown

Notes: (1) Results from Ref. 728 correspond to projectile emission measurements and no corrections were made for contamination of metastable atomic beam with ground state atoms. Results have been normalized by the authors to previous results on excitation. (2) Experimental details of Ref. 729 are unknown.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 5.0E+01 eV/amu, E_{max} = 1.2E+03 eV/amu

A0	A1	A2	A3	A4	A5	A6
-79.7131	.0521928	-.176441	-.0874202	-.203462	.0710365	.0424143

The fit represents the above cross section with an rms deviation of 4.9%. The maximum deviation is 10.5% at 2.0E+02 eV/amu.

See appendix for Chebyshev fit details.

$He(2^3S) + He(1^1S) \rightarrow He(3^1P) + He$



Projectile Excitation Cross Sections for

$He(2^3S) + He(1^1S) \rightarrow He(3^3P) + He$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
3.5E+01	8.22E+06	3.51E-17
4.0E+01	8.79E+06	4.07E-17
5.0E+01	9.82E+06	5.05E-17
6.0E+01	1.08E+07	5.96E-17
7.0E+01	1.16E+07	6.67E-17
8.0E+01	1.24E+07	7.23E-17
9.0E+01	1.32E+07	7.70E-17
1.0E+02	1.39E+07	8.11E-17
1.5E+02	1.70E+07	9.36E-17
2.0E+02	1.96E+07	9.94E-17
2.5E+02	2.20E+07	1.02E-16
3.0E+02	2.41E+07	1.03E-16
4.0E+02	2.78E+07	1.02E-16
5.0E+02	3.11E+07	1.00E-16
6.0E+02	3.40E+07	9.93E-17
7.0E+02	3.68E+07	9.82E-17
8.0E+02	3.93E+07	9.68E-17
9.0E+02	4.17E+07	9.49E-17
1.0E+03	4.39E+07	9.35E-17
1.3E+03	5.01E+07	8.48E-17

References: 728, 729

Accuracy: Unknown

Notes: (1) Results from Ref. 728 correspond to projectile emission measurements and no corrections were made for contamination of metastable atomic beam with ground state atoms. (2) Results have been normalized by the authors to previous results on excitation. (3) Experimental details of Ref. 729 are unknown.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 3.5E+01 eV/amu, E_{max} = 1.3E+03 eV/amu

A0	A1	A2	A3	A4	A5	A6
-74.2632	.399374	-.300481	.0525981	-.0111543	-.00859422	-.00474980

The fit represents the above cross section with an rms deviation of 0.4%. The maximum deviation is 0.8% at 5.0E+01 eV/amu.

See appendix for Chebyshev fit details.

$He(2^3S) + He(1^1S) \rightarrow He(3^3P) + He$



Projectile Excitation Cross Sections for

$He(2^3S) + He(1^1S) \rightarrow He(4^3S,4^3D) + He$

$He(4^3S)$ Energy (eV/amu)	Cross Section (cm^2)	$He(4^3D)$ Energy (eV/amu)	Cross Section (cm^2)
7.0E+01	1.17E-18	6.0E+01	6.96E-18
8.0E+01	9.49E-19	7.0E+01	7.98E-18
9.0E+01	8.39E-19	8.0E+01	9.02E-18
1.0E+02	7.81E-19	9.0E+01	9.94E-18
1.5E+02	1.22E-18	1.0E+02	1.07E-17
2.0E+02	1.62E-18	1.5E+02	1.36E-17
3.0E+02	1.99E-18	2.0E+02	1.49E-17
1.0E+04	2.70E-18	2.5E+02	1.52E-17
1.5E+04	2.16E-18	3.0E+02	1.42E-17
2.0E+04	1.76E-18	4.0E+02	1.16E-17
2.5E+04	1.49E-18	5.0E+02	1.05E-17
2.7E+04	1.39E-18	6.0E+02	1.09E-17
		7.0E+02	1.16E-17
		8.0E+02	1.21E-17
		9.0E+02	1.26E-17
		1.0E+03	1.29E-17
		1.5E+03	1.39E-17
		2.0E+03	1.39E-17
		3.0E+03	1.33E-17
		4.0E+03	1.22E-17
		5.0E+03	1.11E-17
		6.0E+03	1.00E-17
		7.0E+03	9.02E-18
		8.0E+03	8.13E-18
		9.0E+03	7.36E-18
		1.0E+04	6.61E-18
		1.5E+04	3.87E-18
		2.0E+04	2.28E-18
		2.5E+04	1.41E-18

References: 727, 728, 729, 730

Accuracy: Unknown

Notes: (1) Results from Ref. 728 correspond to projectile emission cross sections and no correction was made for contamination of metastable atomic beam with ground state atoms. (2) Results have been normalized by the authors to previous results on excitation. (3) Experimental details of Ref. 729 are unknown. (4) Dotted curve for 4^3S excitation between .3 and 10 keV/amu represents an interpolation between the data of Ref. 727 and Ref. 729, and has a large uncertainty.

Chebyshev Fitting Parameters for Cross Sections

$He(4^3S)$ E_{min} = 7.0E+01 eV/amu, E_{max} = 2.7E+04 eV/amu
$He(4^3D)$ E_{min} = 6.0E+01 eV/amu, E_{max} = 2.5E+04 eV/amu

	A0	A1	A2	A3	A4	A5	A6	A7	A8
$He(4^3S)$	-81.7091	.309903	-.473244	-.0741860	.0175929	-.112721	.0812889	-.0416811	.0437011
$He(4^3D)$	-78.8400	-.637167	-.719668	-.186552	-.198706	.0689624	.0116222	-.0618008	.0358345

The fit represents the $He(4^3S)$ cross section with an rms deviation of 2.6%. The maximum deviation is 7.0% at 1.0E+02 eV/amu.

The fit represents the $He(4^3D)$ cross section with an rms deviation of 3.7%. The maximum deviation is 9.8% at 5.0E+02 eV/amu.

See appendix for Chebyshev fit details.

$He(2^3S) + He \rightarrow He(4^3S,4^3D) + He$



Target Excitation Crosss Sections for

$He(2^3S) + He(1^1S) \rightarrow He + He(4^3S,4^3D)$

$He(4^3S)$ Energy (eV/amu)	Cross Section (cm^2)	$He(4^3D)$ Energy (eV/amu)	Cross Section (cm^2)
5.0E+03	6.41E-18	5.0E+03	1.17E-17
6.0E+03	7.20E-18	6.0E+03	1.39E-17
7.0E+03	7.72E-18	7.0E+03	1.48E-17
8.0E+03	7.92E-18	7.5E+03	1.49E-17
9.0E+03	8.01E-18	8.0E+03	1.47E-17
1.0E+04	7.89E-18	9.0E+03	1.38E-17
1.5E+04	5.87E-18	1.0E+04	1.22E-17
2.0E+04	3.35E-18	1.2E+04	6.16E-18
2.5E+04	1.50E-18	1.5E+04	3.51E-18
		2.0E+04	1.74E-18
		2.3E+04	1.28E-18

<u>References:</u> 727, 730

<u>Accuracy:</u> 40%

<u>Note:</u> Data are for excitation of the ground state He target atom by the metastable $He(2^3S)$ projectile atom.

Chebyshev Fitting Parameters for Cross Sections

$He + He(4^3S)$ E_{min} = 5.0E+03 eV/amu, E_{max} = 2.5E+04 eV/amu
$He + He(4^3D)$ E_{min} = 5.0E+03 eV/amu, E_{max} = 2.3E+04 eV/amu

	A0	A1	A2	A3	A4	A5	A6
$He + He(4^3S)$	-79.6926	-.631297	-.445191	-.0910388	-.0216050	-.00467890	.00182959
$He + He(4^3D)$	-79.4099	-1.23537	-.423173	.177825	.0609486	-.0537149	-.0270941

The fit represents $He + He(4^3S)$ cross section with an rms deviation of 0.2%. The maximum deviation is 0.4% at 8.0E+03 eV/amu.

The fit represents $He + He(4^3D)$ cross section with an rms deviation of 4.4%. The maximum deviation is 10.7% at 1.2E+04 eV/amu.

See appendix for Chebyshev fit details.

$He(2^3S) + He(1^1S) \rightarrow He(4^3S,4^3D) + He$



Excitation and Spectral Line Emission Cross Sections for

$H^+ + Li(2s) \rightarrow H^+ + Li(2p)$
$\rightarrow H^+ + Li(670.8nm)$

Li(2p)		Li(670.8nm)	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
5.0E+02	9.56E-16	2.0E+03	2.97E-15
7.0E+02	1.20E-15	2.5E+03	3.35E-15
1.0E+03	1.50E-15	3.0E+03	3.69E-15
1.5E+03	1.91E-15	4.0E+03	4.04E-15
2.0E+03	2.25E-15	5.0E+03	4.27E-15
4.0E+03	3.06E-15	6.0E+03	4.43E-15
7.0E+03	3.62E-15	7.0E+03	4.53E-15
1.0E+04	3.96E-15	8.0E+03	4.60E-15
1.3E+04	3.99E-15	9.0E+03	4.65E-15
2.0E+04	3.88E-15	1.0E+04	4.69E-15
4.0E+04	2.38E-15	1.5E+04	4.73E-15
7.0E+04	1.42E-15	2.0E+04	4.55E-15
1.0E+05	1.03E-15		
1.3E+05	8.17E-16		

References: Li(2p); 257, 285
Li(670.8nm) emission; 258, 732, 734

Accuracy: Unknown

Notes: (1) The ground state of Li is the 2s state. (2) The resonance line of Li is the 2p -> 2s transition at a wavelength of 670.8nm. (3) Myers et al. (Ref. 734) have also measured emission cross sections for this reaction in the energy region 100-1500 eV/amu. Their results are more than an order of magnitude lower than the recommended data in the energy overlap region.

Chebyshev Fitting Parameters for Cross Sections

Li(2p) E_{min} = 5.0E+02 eV/amu, E_{max} = 1.3E+05 eV/amu
Li(670.8 nm) E_{min} = 2.0E+03 eV/amu, E_{max} = 2.0E+04 eV/amu

	A0	A1	A2	A3	A4	A5	A6
Li(2p)	-67.9437	-.00943077	-.745522	-.112088	.0453106	.0512591	.0155081
Li(670.8 nm)	-66.2671	.208208	-.0976231	.00780915	-.00712239	-.00227550	.00106025

The fit represents Li(2p) cross section with an rms deviation of 1.9%. The maximum deviation is 3.6% at 4.0E+04 eV/amu.

The fit represents Li(670.8 nm) cross section with an rms deviation of 0.3%. The maximum deviation is 0.4% at 2.5E+03 eV/amu.

See appendix for Chebyshev fit details.

$H^+ + Li(2s) \rightarrow Li(2p, 670.8\ nm\ emission)$



Total Spectral Line Emission Cross Sections for

$He^+ + Li(2s) \rightarrow He^+ + Li(670.8nm)$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
5.0E+02	3.11E+07	5.58E-16
6.0E+02	3.40E+07	6.55E-16
7.0E+02	3.68E+07	7.48E-16
8.0E+02	3.93E+07	8.43E-16
9.0E+02	4.17E+07	9.33E-16
1.0E+03	4.39E+07	1.03E-15
1.5E+03	5.38E+07	1.47E-15
2.0E+03	6.21E+07	1.93E-15
2.5E+03	6.95E+07	2.43E-15
3.0E+03	7.61E+07	2.90E-15
4.0E+03	8.79E+07	3.39E-15
4.5E+03	9.32E+07	3.51E-15
5.0E+03	9.82E+07	3.52E-15

References: 267, 732, 734

Accuracy: Unknown

Notes: (1) The 2p -> 2s transition at 670.8nm is the resonance transition of Li. (2) The emission cross section includes cascading into the 2p state from the upper levels. (3) As far as we are aware no data for the cross sections for the 2p state excitation exist. (4) Myers et al. (Ref. 734) obtained data for the emission cross sections in the energy range 50-375 eV/amu. Their measurements are an order of magnitude less than the recommended data at 400 eV/amu.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 5.0E+02 eV/amu, E_{max} = 5.0E+03 eV/amu

A0	A1	A2	A3	A4	A5	A6
-68.2494	.976730	-.0528574	-.0520709	-.0299645	-.00494558	.00671922

The fit represents the above cross section with an rms deviation of 0.5%. The maximum deviation is 0.8% at 2.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + Li(2s) \rightarrow Li(2p)$ (670.8 nm emission)



Excitation and Spectral Line Emission Cross Sections for

He^{2+} + Li(2s) -> He^{2+} + Li(2p)
-> He^{2+} + Li(670.8nm)

Li(2p)		Li(2p) (cont.)		Li(670.8nm)	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
1.4E+02	3.90E-17	3.8E+03	2.79E-15	3.7E+03	2.92E-15
2.0E+02	6.43E-17	4.0E+03	2.94E-15	4.0E+03	3.13E-15
3.0E+02	1.11E-16	7.0E+03	4.93E-15	7.0E+03	5.21E-15
4.0E+02	1.62E-16	1.0E+04	6.57E-15	1.0E+04	6.96E-15
5.0E+02	2.18E-16	2.0E+04	7.93E-15	1.5E+04	8.45E-15
6.0E+02	2.73E-16	2.6E+04	8.12E-15	2.0E+04	9.07E-15
7.0E+02	3.28E-16	4.0E+04	7.50E-15	2.6E+04	9.21E-15
8.0E+02	3.84E-16	7.0E+04	6.04E-15	4.0E+04	8.69E-15
9.0E+02	4.36E-16	1.0E+05	5.10E-15	7.0E+04	6.94E-15
1.0E+03	4.90E-16	2.0E+05	3.25E-15	1.0E+05	5.81E-15
1.5E+03	7.52E-16	4.0E+05	1.82E-15	1.5E+05	4.70E-15
2.0E+03	9.89E-16	7.0E+05	1.09E-15		
		1.0E+06	7.68E-16		
		2.0E+06	3.93E-16		
		2.7E+06	2.95E-16		

<u>References:</u> Li(2p); 731, 733, 735
Li(670.8nm) emission; 259, 733, 735

<u>Accuracy:</u> Unknown

<u>Notes:</u> (1) The 2p -> 2s is the resonance transition of Li with λ = 670.8nm. (2) No attempt has been made to join the low energy data of Barrett and Leventhal (Ref. 259) to the data of Kadota et al. (Ref. 733). (3) The data of Kadota et al. have been extended to 2.5×10^6 eV/amu using the theoretical data of Ermolaev et al. (Ref. 731).

<u>Chebyshev Fitting Parameters for Cross Sections</u>

Li(2p)	E_{min} =	1.4E+02	eV/amu,	E_{max} =	2.0E+03	eV/amu
Li(2p)	E_{min} =	3.8E+03	eV/amu	E_{max} =	2.7E+06	eV/amu
Li(670.8 nm)	E_{min} =	3.7E+03	eV/amu,	E_{max} =	1.5E+05	eV/amu

	A0	A1	A2	A3	A4	A5	A6
Li(2p)	-72.1632	1.62490	-.0826001	-.0110092	.00206946	.00278599	-.00427450
Li(2p)	-67.5365	-1.31752	-.867548	.198289	-.0106036	.00118819	.0110240
Li(670.8 nm)	-65.6114	.210669	-.460211	.0351576	.0330529	-.00874795	.00505234

The fit represents Li(2p) excitation cross section with an rms deviation of 0.2%.
The maximum deviation is 0.4% at 4.0E+02 eV/amu.

The fit represents Li(2p) cross section with an rms deviation of 1.1%.
The maximum deviation is 2.3% at 7.0E+03 eV/amu.

The fit represents Li(670.8 nm) cross section with an rms deviation of 0.4%.
The maximum deviation is 0.5% at 7.0E+04 eV/amu.

He^{2+} + Li(2s) –> Li(2p, 670.8 nm emission)



D. Cross Sections for Ionization and Production of Charged Particles

(* denotes rate coefficient data also)

	Page
$H + H_2$ -> Total Slow Positive Ion Production . . .	D-2
-> Total Slow Electron Production	D-2
-> $H + H_2^+ + e$	D-4
$H^+ + H$ -> $H^+ + H^+ + e$	D-6*
$H^+ + H^-$ -> $H_2^+ + e$	D-12
$H^+ + H_2$ -> Total Slow Electron Production	D-14
-> $H^+ + H_2 + e$	D-16
-> Total Production of H_2^+	D-16
-> Total Production of H^+	D-16
$H^+ + He$ -> Electrons and H^-	D-18*
-> $H^+ + He^+ + e$	D-24*
-> $(H + He^+)$ or $(H^+ + He^+ + e)$	D-24*
-> $H^+ + He^{2+} + 2e$	D-34*
-> $(H^- + He^{2+})$ or $(H^+ + He^{2+} + 2e)$	D-34*
$He(2^1S) + H$ -> $He + H^+ + e$	D-42
-> $HeH^+ + e$	D-42
-> $(He + H^+ + e)$ or $(HeH^+ + e)$	D-42
$He(2^3S) + H$ -> $He + H^+ + e$	D-44
-> $HeH^+ + e$	D-44
-> $(He + H^+ + e)$ or $(HeH^+ + e)$	D-44
$He(2^3S) + H_2$ -> Total Ionization	D-46
-> Rearrangement Ionization	D-46
$He(2^1S) + H_2$ -> Total Ionization	D-48
-> Rearrangement Ionization	D-48
$H^+ + He^+$ -> $(H^+ + He^{2+} + e)$	D-50*
-> $(H^+ + He^{2+} + e)$ or $(H^o + He^{2+})$	D-50*
$He^+ + He^+$ -> $(He^+ + He^{2+} + e)$	D-58*
-> $(He + He^{2+})$ or $(He^+ + He^{2+} + e)$ . . .	D-58*
$He + H_2$ -> Total Slow Positive Ion Production . .	D-64
-> Total Slow Electron Production	D-64
-> $He + H_2^+ + e$	D-66
-> Total H^+ Production	D-66
$He + He$ -> Total Slow Positive Ion Production . . .	D-68
-> Total Slow Electron Production	D-68
-> $He + He^+ + e$	D-70
-> $He + He^{2+} + 2e$	D-70

D. (Cont'd)

$He^+ + H \rightarrow He^+ + H^+ + e$ D-72*

$He^+ + H_2$ -> Total Slow Positive Ion Production . . D-76
-> Total Slow Electron Production D-76

$He^+ + H_2$ -> $(He + H_2^+)$ or $(He^+ + H_2^+ + e)$ D-78
-> Total H^+ Production D-78
-> $He^+ + H^+ + H^+ + 2e$ D-78

$He^+ + He$ -> Total Slow Positive Ion Production . . D-80*
-> Total Slow Electron Production D-80*
-> $He^+ + He^+ + e$ D-90*
-> $He^+ + He^{2+} + 2e$ D-90*

$He^{2+} + H \rightarrow He^{2+} + H^+ + e$ D-96*

$He^{2+} + H_2$ -> Total Slow Positive Ion Production . . D-100
-> Total Slow Electron Production D-100
-> Total H^+ Production D-102
-> H^+ Production by Ionization Collisions D-102
-> Total H_2^+ Production D-102
-> H_2^+ Production by Single-Ionization . D-102

$He^{2+} + He$ -> Total Slow Positive Ion Production . . D-104*
-> Total Slow Electron Production D-104*
-> $He^{2+} + He^+ + e$ D-114*
-> $He^{2+} + He^{2+} + 2e$ D-114*

$H^+ + Li \rightarrow H^+ + Li^+ + e$ D-122*

$He^+ + Li \rightarrow He^+ + Li^+ + e$ D-126

$He^{2+} + Li \rightarrow He^{2+} + Li^+ + e$ D-128*

Cross Sections for Slow Positive-Ion and Electron Production in Collisions of Hydrogen Atoms with Molecular Hydrogen

$H + H_2$ -> Total Slow Positive Ion Prod.
$H + H_2$ -> Total Slow Electron Prod.

Slow Positive Ion Production		Electron Production	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
4.0E+01	5.51E-19	8.0E+01	2.83E-19
7.0E+01	1.65E-18	1.0E+02	5.71E-19
1.0E+02	2.51E-18	2.0E+02	2.65E-18
2.0E+02	3.37E-18	4.0E+02	9.67E-18
3.0E+02	3.17E-18	7.0E+02	2.38E-17
4.0E+02	3.26E-18	1.0E+03	3.98E-17
7.0E+02	4.73E-18	2.0E+03	9.33E-17
1.0E+03	7.28E-18	3.0E+03	1.24E-16
2.0E+03	1.97E-17	4.0E+03	1.29E-16
4.0E+03	5.06E-17	5.0E+03	1.32E-16
1.0E+04	1.27E-16	7.0E+03	1.50E-16
2.0E+04	1.75E-16	1.0E+04	2.00E-16
4.0E+04	1.60E-16	2.0E+04	2.90E-16
1.0E+05	1.15E-16	3.0E+04	3.08E-16
2.0E+05	8.14E-17	4.0E+04	2.98E-16
4.0E+05	5.02E-17	7.0E+04	2.56E-16
		1.0E+05	2.26E-16
		2.0E+05	1.58E-16
		4.0E+05	1.02E-16

References: 28, 30, 143, 287, 315, 323, 324

Accuracy: 20%

Notes: (1) The slow positive charge formation cross section is the sum of the ionization cross section and the electron capture cross section. $\sigma^+ = \sigma_1 (H_2^+ + e) + \sigma_{0-1} (H^- + H_2^+)$. The dissociative ionization cross section ($H + H_2$ -> $H + H^+ + H + e$) is only 2-5% of the H_2^+ formation cross section in this energy range. (2) At low energies most of the H_2^+ arises from electron capture collisions. (3) The electron production cross section is the sum of ionization and the electron loss or stripping cross section $\sigma^+ = \sigma_1 (H_2^+ + e) + \sigma_{01} (H^+ + e)$.

Chebyshev Fitting Parameters for Cross Sections

Positive Ions E_{min} = 4.0E+01 eV/amu, E_{max} = 4.0E+05 eV/amu
Electrons E_{min} = 8.0E+01 eV/amu, E_{max} = 4.0E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
Pos. Ions	-77.2533	2.45645	-.972849	-.533057	-.0537388	.365771	-.154454
Electrons	-75.9790	2.77048	-1.73727	.139659	-.0624901	-.00991339	.0680936

The fit represents the Positive Ions cross section with an rms deviation of 6.7%. The maximum deviation is 14.9% at 7.0E+01 eV/amu.

The fit represents the Electrons cross section with an rms deviation of 8.5%. The maximum deviation is 17.8% at 7.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$H + H_2$ –> Sum of Slow Positive Charges
$H + H_2$ –> Sum of Electrons Produced



Ionization Cross Sections for

$$H + H_2 \rightarrow H + H_2^+ + e$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.0E+02	1.96E+07	4.04E-19
4.0E+02	2.78E+07	1.42E-18
7.0E+02	3.68E+07	3.32E-18
1.0E+03	4.39E+07	5.48E-18
2.0E+03	6.21E+07	1.49E-17
4.0E+03	8.79E+07	3.71E-17
7.0E+03	1.16E+08	6.72E-17
1.0E+04	1.39E+08	9.30E-17
2.0E+04	1.96E+08	1.45E-16
2.8E+04	2.32E+08	1.54E-16
4.0E+04	2.78E+08	1.46E-16
7.0E+04	3.68E+08	1.23E-16
1.0E+05	4.39E+08	1.08E-16
2.0E+05	6.21E+08	8.13E-17
4.0E+05	8.78E+08	4.99E-17

References: 28, 30, 143, 308, 315, 323, 325

Accuracy: 30%

Notes: (1) The ionization cross section is determined from either of two relations: $\sigma_i = \sigma^- - \sigma_{01}$ or $\sigma_i = \sigma^+ - \sigma_{0-1}$, where σ^- is the cross section for formation of electrons, σ_{01} is the loss or stripping cross section, σ^+ is the cross section for slow ion formation, and σ_{0-1} is the electron capture cross section. (2) There is considerable scatter in the data among the various investigators.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 2.0E+02 eV/amu, E_{max} = 4.0E+05 eV/amu

A0	A1	A2	A3	A4	A5	A6
-77.1205	2.47663	-1.43551	-.126243	.100447	.0662120	-.0477593

The fit represents the above cross section with an rms deviation of 2.7%. The maximum deviation is 4.1% at 7.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$$H + H_2 \rightarrow H + H_2^+ + e$$



Ionization Cross Sections for

$$H^+ + H \rightarrow H^+ + H^+ + e$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
9.4E+03	1.35E+08	1.62E-17
1.0E+04	1.39E+08	1.84E-17
1.5E+04	1.70E+08	4.21E-17
2.0E+04	1.96E+08	7.20E-17
3.0E+04	2.41E+08	1.21E-16
4.0E+04	2.78E+08	1.39E-16
4.5E+04	2.95E+08	1.41E-16
5.0E+04	3.11E+08	1.40E-16
6.0E+04	3.40E+08	1.37E-16
7.0E+04	3.68E+08	1.30E-16
8.0E+04	3.93E+08	1.23E-16
9.0E+04	4.17E+08	1.16E-16
1.0E+05	4.39E+08	1.10E-16
1.5E+05	5.38E+08	8.72E-17
2.0E+05	6.21E+08	7.14E-17
3.0E+05	7.61E+08	5.14E-17
4.0E+05	8.78E+08	4.04E-17
7.0E+05	1.16E+09	2.54E-17
1.0E+06	1.39E+09	1.88E-17
1.5E+06	1.70E+09	1.34E-17

<u>References:</u> 4, 261, 335, 336, 337, 338, 339, 340, 341, 342, 383, 586

<u>Accuracy:</u> $E > 4x10^4$ eV/amu - 25%
$E < 4x10^4$ eV/amu - Unknown

<u>Notes:</u> (1) Data above $4x10^4$ eV/amu are the recommended data of Rudd, et al. (Ref. 335). Below this energy, the recent data of Shah et al. (Ref. 336) indicate a marked departure from the evaluated data of Rudd. The data plotted here in the low energy range are the recent results of Shah et al. (Ref. 336). (2) For a hydrogen atom target, the cross sections for slow electron production (σ^-) are equal to the ionization cross sections (σ_i). For slow ion production (σ^+) the cross sections are the sum of the electron capture cross sections (σ_{01}) and the ionization cross sections at high energies $\sigma^- = \sigma^+ = \sigma_i$.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

E_{min} = 9.4E+03 eV/amu, E_{max} = 1.5E+06 eV/amu

A0	A1	A2	A3	A4	A5	A6
-75.5053	-.409785	-.996553	.355360	-.049000	-.0349984	.0437825

The fit represents the above cross section with an rms deviation of 1.7%. The maximum deviation is 2.7% at 4.0E+05 eV/amu.

See appendix for Chebyshev fit details.

$$H^+ + H \rightarrow H^+ + H^+ + e$$



Ionization Rate Coefficients for

$$H^+ + H \rightarrow H^+ + H^+ + e$$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	Equal Temp.	H^+ Temp. (eV) 1000.	2000.	4000.	7000.	10000.	15000.	20000.
1.0E+03	9.60E-11#	9.60E-11#	5.16E-10#	2.35E-09*	6.44E-09*	1.08E-08	1.74E-08	2.25E-08
1.5E+03	5.16E-10#	2.59E-10#	8.64E-10#	2.96E-09*	7.18E-09	1.16E-08	1.79E-08	2.30E-08
2.0E+03	1.29E-09*	5.16E-10#	1.29E-09*	3.61E-09*	7.92E-09	1.23E-08	1.85E-08	2.34E-08
2.5E+03	2.35E-09*	8.64E-10#	1.79E-09*	4.29E-09*	8.66E-09	1.29E-08	1.91E-08	2.38E-08
3.0E+03	3.61E-09*	1.29E-09*	2.35E-09*	4.99E-09*	9.40E-09	1.36E-08	1.96E-08	2.42E-08
3.5E+03	4.99E-09*	1.79E-09*	2.96E-09*	5.71E-09*	1.01E-08	1.43E-08	2.01E-08	2.46E-08
4.0E+03	6.44E-09*	2.35E-09*	3.61E-09*	6.44E-09*	1.08E-08	1.49E-08	2.06E-08	2.50E-08
4.5E+03	7.92E-09	2.96E-09*	4.29E-09*	7.18E-09	1.16E-08	1.56E-08	2.11E-08	2.54E-08
5.0E+03	9.40E-09	3.61E-09*	4.99E-09*	7.92E-09	1.23E-08	1.62E-08	2.16E-08	2.58E-08
5.5E+03	1.08E-08	4.29E-09*	5.71E-09*	8.66E-09	1.29E-08	1.68E-08	2.21E-08	2.61E-08
6.0E+03	1.23E-08	4.99E-09*	6.44E-09*	9.40E-09	1.36E-08	1.74E-08	2.25E-08	2.65E-08
8.0E+03	1.74E-08	7.92E-09	9.40E-09	1.23E-08	1.62E-08	1.96E-08	2.42E-08	2.78E-08
1.0E+04	2.16E-08	1.08E-08	1.23E-08	1.49E-08	1.85E-08	2.16E-08	2.58E-08	2.90E-08
1.2E+04	2.50E-08	1.36E-08	1.49E-08	1.74E-08	2.06E-08	2.34E-08	2.72E-08	3.01E-08
1.4E+04	2.78E-08	1.62E-08	1.74E-08	1.96E-08	2.25E-08	2.50E-08	2.84E-08	3.10E-08
1.6E+04	3.01E-08	1.85E-08	1.96E-08	2.16E-08	2.42E-08	2.65E-08	2.95E-08	3.19E-08
1.8E+04	3.19E-08	2.06E-08	2.16E-08	2.34E-08	2.58E-08	2.78E-08	3.06E-08	3.27E-08
2.0E+04	3.35E-08	2.25E-08	2.34E-08	2.50E-08	2.72E-08	2.90E-08	3.15E-08	3.35E-08

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+03 eV, E_{max} = 2.0E+04 eV

H Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
1000.	-40.0051	2.76605	-.344647	-.0369496	.00899170	-3.564719E-04	4.064999E-04
2000.	-38.7629	1.97055	-.100798	-.0656769	.0461132	.00211656	-1.279077E-04
4000.	-37.4352	1.22903	.0356592	-.0490361	-.00390032	.00203472	2.011422E-04
7000.	-36.4087	.743427	.0691035	-.0248017	-.00589077	5.652350E-04	3.380127E-04
10000.	-35.8193	.504234	.0663753	-.0124429	-.00494956	-1.095647E-04	2.357336E-04
15000.	-35.2398	.301656	.0518339	-.00363904	-.00313907	-4.112259E-04	8.863478E-05
20000.	-34.8966	.198907	.0391475	-5.009338E-04	-.00198385	-3.886616E-04	1.159612E-05
Equal Temp.	-38.7626	2.81265	-.736000	.106667	-.0206816	.00733904	-.00185096

See appendix for Chebyshev fit details.

$$H^+ + H \rightarrow H^+ + H^+ + e$$

Maxwellian — Maxwellian



Ionization Rate Coefficients for

$$H + H^+ \rightarrow H^+ + H^+ + e$$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	H Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	2.56E-09	1.41E-08	3.86E-08	4.78E-08	4.83E-08	4.43E-08	3.30E-08
2.0E+00	2.56E-09	1.41E-08	3.86E-08	4.78E-08	4.83E-08	4.43E-08	3.30E-08
4.0E+00	2.53E-09	1.41E-08	3.85E-08	4.78E-08	4.83E-08	4.43E-08	3.30E-08
7.0E+00	2.47E-09	1.41E-08	3.85E-08	4.78E-08	4.83E-08	4.43E-08	3.30E-08
1.0E+01	2.40E-09	1.41E-08	3.85E-08	4.78E-08	4.83E-08	4.43E-08	3.30E-08
2.0E+01	2.27E-09*	1.41E-08	3.85E-08	4.77E-08	4.83E-08	4.43E-08	3.30E-08
4.0E+01	2.19E-09*	1.41E-08	3.84E-08	4.77E-08	4.83E-08	4.43E-08	3.30E-08
7.0E+01	2.18E-09*	1.41E-08	3.84E-08	4.77E-08	4.83E-08	4.43E-08	3.30E-08
1.0E+02	2.22E-09*	1.42E-08	3.83E-08	4.77E-08	4.83E-08	4.43E-08	3.30E-08
2.0E+02	2.42E-09*	1.44E-08	3.82E-08	4.77E-08	4.82E-08	4.43E-08	3.30E-08
4.0E+02	2.88E-09*	1.49E-08	3.81E-08	4.76E-08	4.82E-08	4.42E-08	3.30E-08
7.0E+02	3.58E-09*	1.56E-08	3.79E-08	4.75E-08	4.81E-08	4.42E-08	3.30E-08
1.0E+03	4.25E-09*	1.63E-08	3.77E-08	4.73E-08	4.81E-08	4.41E-08	3.30E-08
2.0E+03	6.40E-09	1.81E-08	3.74E-08	4.69E-08	4.79E-08	4.40E-08	3.30E-08
4.0E+03	1.02E-08	2.09E-08	3.71E-08	4.62E-08	4.76E-08	4.38E-08	3.30E-08
7.0E+03	1.49E-08	2.40E-08	3.70E-08	4.52E-08	4.70E-08	4.36E-08	3.29E-08
1.0E+04	1.86E-08	2.64E-08	3.72E-08	4.45E-08	4.64E-08	4.33E-08	3.29E-08
2.0E+04	2.73E-08	3.18E-08	3.82E-08	4.31E-08	4.47E-08	4.26E-08	3.29E-08

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

H Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-38.2968	1.12152	.644681	.100717	-.100166	-.0395957	.00265964
20000.	-35.7290	.356274	.202092	.0620743	-.00276894	-.0115918	-.00410154
40000.	-34.1679	-.0162401	3.246363E-04	.00754856	.00728055	.00420183	.00124748
70000.	-33.7606	-.0400418	-.0258461	-.0120904	-.00336001	2.548592E-04	9.663075E-04
100000.	-33.7204	-.0256605	-.0180094	-.0107132	-.00535053	-.00208862	-4.645604E-04
200000.	-33.8801	-.0146171	-.00902218	-.00477874	-.00229897	-9.828762E-04	-3.658987E-04
500000.	-34.4565	-7.913823E-04	-5.710595E-04	-3.486254E-04	-1.820028E-04	-8.154704E-05	-3.512063E-05

See appendix for Chebyshev fit details.

$$H + H^+ \rightarrow H^+ + H^+ + e$$



Associative Detachment Cross Sections for

$$H^+ + H^- \rightarrow H_2^+ + e$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.0E-03	6.21E+04	5.12E-14
4.0E-03	8.79E+04	2.23E-14
7.0E-03	1.16E+05	1.36E-14
1.0E-02	1.39E+05	1.08E-14
2.0E-02	1.96E+05	6.77E-15
4.0E-02	2.78E+05	3.77E-15
7.0E-02	3.68E+05	2.19E-15
1.0E-01	4.39E+05	1.55E-15
2.0E-01	6.21E+05	8.79E-16
4.0E-01	8.79E+05	4.96E-16
7.0E-01	1.16E+06	3.25E-16
1.0E+00	1.39E+06	2.32E-16
2.0E+00	1.96E+06	1.27E-16
4.0E+00	2.78E+06	5.18E-17
7.0E+00	3.68E+06	2.37E-17

References: 417, 418, 419

Accuracy: 30%

Notes: (1) The theoretical data of (Ref. 417) are within a few percent of the experimental data of (Ref. 418). (2) Measurements of $D^+ + D^- \rightarrow D_2^+ + e$ are consistently 30-50% below those for $H^+ + H^-$. At energies greater than 1 eV/amu the fluctuations in the data mask this difference.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 2.0E-03 eV/amu, E_{max} = 7.0E+00 eV/amu

A0	A1	A2	A3	A4	A5	A6
-68.6095	-3.63490	-.153293	-0.133084	-.0284174	-.0787410	.0439445

The fit represents the above cross section with an rms deviation of 2.2%. The maximum deviation is 3.4% at 7.0E-03 eV/amu.

See appendix for Chebyshev fit details.

$$H^+ + H \rightarrow H_2^+ + e$$



Total Slow Electron Production Cross Sections for

$H^+ + H_2 \rightarrow \Sigma e$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
5.0E+02	3.11E+07	1.44E-18
7.0E+02	3.68E+07	2.15E-18
1.0E+03	4.39E+07	3.32E-18
2.0E+03	6.21E+07	7.85E-18
4.0E+03	8.79E+07	1.84E-17
7.0E+03	1.16E+08	3.67E-17
1.0E+04	1.39E+08	5.66E-17
2.0E+04	1.96E+08	1.18E-16
4.0E+04	2.78E+08	1.96E-16
6.5E+04	3.54E+08	2.19E-16
7.0E+04	3.68E+08	2.19E-16
1.0E+05	4.39E+08	1.96E-16
2.0E+05	6.21E+08	1.33E-16
4.0E+05	8.78E+08	7.96E-17
7.0E+05	1.16E+09	5.08E-17
1.0E+06	1.39E+09	3.74E-17
2.0E+06	1.96E+09	2.07E-17
4.0E+06	2.77E+09	1.15E-17
5.0E+06	3.09E+09	9.50E-18

<u>References:</u> 30, 36, 42, 45, 89, 217, 315, 335, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 366

<u>Accuracy:</u> 25%

<u>Notes:</u> (1) The data are the recommended data of Rudd et al. (Ref. 335). See this reference for an analytical fit to the data at low and high energies. (2) The slow electron production cross sections are the sum of four reactions: (a) $H^+ + H_2 \rightarrow H^+ + H_2^+ + e$ (pure ionization); (b) $\rightarrow H^+ + H + H^+ + e$ (ionization with dissociation); (c) $\rightarrow H^+ + H^+ + e$ (double ionization); (d) $\rightarrow H + 2H^+ + e$ (electron capture with ionization).

<u>Chebyshev Fitting Parameters for Cross Sections</u>

E_{min} = 5.0E+02 eV/amu, E_{max} = 5.0E+06 eV/amu

A0	A1	A2	A3	A4	A5	A6
-76.9159	.969378	-1.91735	-.0245092	.313763	-.000351145	-.0992103

The fit represents the above cross section with an rms deviation of 3.1%. The maximum deviation is 6.2% at 2.0E+05 eV/amu.

See appendix for Chebyshev fit details.

$H^+ + H_2 \rightarrow \Sigma e$



Cross Sections for the Formation of Positive Ions in Collisions of H^+ with H_2

$H^+ + H_2 \rightarrow H^+ + H_2^+ + e$ (Single Ionization)
$H^+ + H_2 \rightarrow$ Total Production of H_2^+ (or Total Ion Formation)
$H^+ + H_2 \rightarrow$ Total Production of H^+

Single Ionization		H_2^+ Production		H^+ Production	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
5.0E+03	1.23E-17	6.0E+03	8.23E-16	5.0E+03	3.71E-17
7.0E+03	1.78E-17	7.0E+03	8.37E-16	6.0E+03	4.93E-17
1.0E+04	2.63E-17	1.0E+04	8.03E-16	7.0E+03	6.11E-17
1.5E+04	4.24E-17	2.0E+04	6.09E-16	8.0E+03	7.25E-17
2.0E+04	6.23E-17	4.0E+04	4.13E-16	9.0E+03	8.29E-17
3.0E+04	1.30E-16	7.0E+04	2.93E-16	1.0E+04	9.16E-17
4.0E+04	1.83E-16	1.0E+05	2.32E-16	1.5E+04	1.17E-16
6.0E+04	2.20E-16	1.5E+05	1.75E-16	1.7E+04	1.20E-16
7.0E+04	2.17E-16	2.0E+05	1.43E-16	2.0E+04	1.17E-16
1.0E+05	2.04E-16	4.0E+05	8.69E-17	2.5E+04	1.02E-16
1.5E+05	1.74E-16	7.0E+05	5.51E-17	3.0E+04	8.36E-17
2.0E+05	1.44E-16	1.0E+06	4.05E-17	3.5E+04	7.03E-17
4.0E+05	8.71E-17	1.5E+06	2.85E-17	4.0E+04	6.03E-17
7.0E+05	5.54E-17	2.0E+06	2.24E-17	4.5E+04	5.24E-17
1.0E+06	4.07E-17	3.0E+06	1.56E-17	5.0E+04	4.62E-17
2.0E+06	2.24E-17	3.5E+06	1.37E-17		
3.0E+06	1.57E-17				
3.5E+06	1.37E-17				
3.8E+06	1.28E-17				

<u>References:</u> 30, 36, 40, 89, 315, 346, 348, 350, 353, 354, 356, 360, 361, 362, 363, 364, 365, 366

<u>Accuracy:</u> 30%

<u>Notes:</u> (1) The interaction of H^+ with H_2 to produce H_2^+ ions may follow one of two pathways: (a) $H^+ + H_2 \rightarrow H^+ + H_2^+ + e$, single ionization; (b) $H^+ + H_2 \rightarrow H + H_2^+$, electron capture. Five reactions are involved in the production of H^+ ions; (c) $H^+ + H_2^+ \rightarrow H^+ + H + H^+ + e$, dissociative ionization; (d) $\rightarrow H^+ + H^+ + H^+ + 2e$, double ionization; (e) $\rightarrow H + H + H^+$, electron capture with dissociation; (f) $\rightarrow H + H^+ + H^+ + e$, transfer ionization; (g) $\rightarrow H^- + 2H^+$, double electron capture. (2) At energies less than 150 keV/amu the total charge production cross sections (σ_t^+) are dominated by electron capture. (3) Within the experimental error the total charge production cross section is the same as the total cross section for producing H_2^+. (4) Above 150 keV the single ionization cross section (reaction a) is the dominant process to form positive charges or H_2^+. (5) Cross sections are available for the double ionization (Ref. 36, 348, 356, 360, 365) (reaction c). However there are large discrepancies and these data are not included here. The cross sections are at least an order of magnitude smaller than the single ionization cross sections.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

Sing. Ion.	E_{min} =	5.0E+03 eV/amu,	E_{max} =	3.8E+06 eV/amu
H_2^+ Prod.	E_{min} =	6.0E+03 eV/amu,	E_{max} =	3.5E+06 eV/amu
H^+ Prod.	E_{min} =	5.0E+03 eV/amu,	E_{max} =	5.0E+04 eV/amu

	A0	A1	A2	A3	A4	A5	A6
Sing. Ion	-75.4752	-.100192	-1.35816	.230586	.187780	-.157087	.0102380
H_2^+ Prod.	-72.9773	-2.12495	-.260502	.0542968	-.0279667	.0221059	-.00372930
H^+ Prod.	-74.4647	.0986066	-.525436	.00183837	.0396132	.0121041	-.00373826

The fit represents the Single Ion. cross section with an rms deviation of 7.1%. The maximum deviation is 17.3% at 2.0E+04 eV/amu.
The fit represents the H_2^+ Prod. cross section with an rms deviation of 0.4%. The maximum deviation is 0.7% at 2.0E+05 eV/amu.
The fit represents the H^+ Prod. cross section with an rms deviation of 0.6%. The maximum deviation is 1.2% at 3.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$H^+ + H_2 \rightarrow H^+ + H_2^+ + e$
$He^- + H_2 \rightarrow$ Total H^+ Production
$He^- + H_2 \rightarrow$ Total H_2^+ Production

Cross Section vs. Energy



Slow Negative Charge Production Cross Sections for

H^+ + He -> electrons and H^-

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2
4.4E+02	2.91E+07	9.65E-20
5.0E+02	3.11E+07	1.16E-19
7.0E+02	3.68E+07	1.92E-19
1.0E+03	4.39E+07	3.28E-19
2.0E+03	6.21E+07	9.40E-19
4.0E+03	8.79E+07	2.73E-18
7.0E+03	1.16E+08	6.35E-18
1.0E+04	1.39E+08	1.09E-17
2.0E+04	1.96E+08	2.96E-17
4.0E+04	2.78E+08	6.68E-17
7.0E+04	3.68E+08	9.07E-17
1.0E+05	4.39E+08	9.28E-17
2.0E+05	6.21E+08	6.86E-17
4.0E+05	8.78E+08	4.24E-17
7.0E+05	1.16E+09	2.77E-17
1.0E+06	1.39E+09	2.09E-17
1.5E+06	1.70E+09	1.53E-17
2.0E+06	1.96E+09	1.20E-17
4.0E+06	2.77E+09	6.79E-18
5.0E+06	3.09E+09	5.63E-18

References: 40, 83, 89, 315, 335, 347, 353, 354, 355, 357, 358, 359, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380

Accuracy: $E > 4x10^4$ eV/amu - 20%
$E < 4x10^4$ eV/amu - Unknown - see note (1)

Notes: (1) These data are the recommended data of Rudd et al. (Ref. 335). The data below $4x10^4$ eV/amu are strongly weighted by the experimental results of Rudd et al. (Ref. 83) taken in 1983. Several investigators have investigated the low energy region, all of them reporting cross sections higher than the recommended data. (2) Reactions that produce slow negative charges are: (a) H^+ + He -> H^+ + He^+ + e, single ionization; (b) H^+ + He -> H^+ + He^{2+} + 2e, double ionization; (c) H^+ + He -> H + He^+ -> H^+ + He^+ + e, capture with ionization or transfer ionization; (d) H^+ + He -> H^- + He^{2+} double electron capture.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 4.4E+02 eV/amu, E_{max} = 5.0E+06 eV/amu

A0	A1	A2	A3	A4	A5	A6
-79.6795	2.14499	-2.18192	-.146158	.362843	.0528127	-.123316

The fit represents the above cross section with an rms deviation of 4.4%. The maximum deviation is 7.3% at 2.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$H^+ + He \rightarrow electrons + H^-$



Slow Negative Charge Production Rate Coefficients for

H^+ + He -> electrons and H^-

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H Temp. (eV)	Equal Temp.	He Temp. (eV) 10.	750.	1000.	1250.	5000.	15000.	20000.
1.0E+02	3.59E-13#	1.61E-13#	3.94E-12*	6.17E-12*	8.83E-12*	1.02E-10	3.79E-10	1.39E-09
7.5E+02	4.86E-11	3.10E-11	4.86E-11	5.54E-11	6.26E-11	2.24E-10	5.88E-10	1.74E-09
1.0E+03	8.71E-11	5.57E-11	7.85E-11	8.71E-11	9.61E-11	2.84E-10	6.79E-10	1.87E-09
1.3E+03	1.37E-10	8.74E-11	1.16E-10	1.26E-10	1.37E-10	3.50E-10	7.76E-10	2.02E-09
1.5E+03	1.97E-10	1.26E-10	1.60E-10	1.72E-10	1.84E-10	4.23E-10	8.79E-10	2.16E-09
1.8E+03	2.69E-10	1.72E-10	2.11E-10	2.24E-10	2.39E-10	5.02E-10	9.87E-10	2.31E-09
2.0E+03	3.50E-10	2.25E-10	2.69E-10	2.84E-10	3.00E-10	5.88E-10	1.10E-09	2.47E-09
3.0E+03	7.76E-10	5.03E-10	5.66E-10	5.88E-10	6.10E-10	9.87E-10	1.60E-09	3.11E-09
4.0E+03	1.34E-09	8.80E-10	9.59E-10	9.87E-10	1.01E-09	1.47E-09	2.16E-09	3.78E-09
5.0E+03	2.02E-09	1.34E-09	1.44E-09	1.47E-09	1.50E-09	2.02E-09	2.78E-09	4.48E-09
6.0E+03	2.78E-09	1.88E-09	1.98E-09	2.02E-09	2.05E-09	2.62E-09	3.44E-09	5.20E-09
7.0E+03	3.61E-09	2.47E-09	2.58E-09	2.62E-09	2.66E-09	3.27E-09	4.13E-09	5.93E-09
8.0E+03	4.48E-09	3.11E-09	3.23E-09	3.27E-09	3.31E-09	3.95E-09	4.84E-09	6.66E-09
9.0E+03	5.38E-09	3.78E-09	3.91E-09	3.95E-09	3.99E-09	4.66E-09	5.56E-09	7.40E-09
1.0E+04	6.30E-09	4.48E-09	4.61E-09	4.66E-09	4.70E-09	5.38E-09	6.30E-09	8.13E-09
1.2E+04	8.13E-09	5.93E-09	6.07E-09	6.11E-09	6.16E-09	6.85E-09	7.76E-09	9.57E-09
1.5E+04	1.08E-08	8.13E-09	8.27E-09	8.31E-09	8.36E-09	9.04E-09	9.92E-09	1.17E-08
2.0E+04	1.48E-08	1.17E-08	1.18E-08	1.18E-08	1.19E-08	1.25E-08	1.33E-08	1.48E-08

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+02 eV, E_{max} = 2.0E+04 eV

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
10.	-46.5650	5.49971	-.475864	.0795255	-.103842	.0146906	5.701499E-04
750.	-44.8867	4.19397	.198206	-.184815	-.0174624	-.00795734	.00438872
1000.	-44.5760	3.97780	.272829	-.192799	-.0215899	-.00569587	.00394217
1250.	-44.3093	3.79820	.326679	-.193047	-.0272627	-.00347228	.00358121
5000.	-42.1265	2.50907	.515276	-.0994986	-.0616701	-.00489874	.00584793
10000.	-40.7163	1.83020	.481655	-.0331261	-.0467863	-.0183645	.00768600
20000.	-39.1385	1.18831	.377951	.0154712	-.0231939	-.0206727	.00437997
Equal Temp.	-45.6407	5.29550	-.441278	.00871718	-.0827904	.0101600	.00309937

See appendix for Chebyshev fit details.

$H^+ + He \rightarrow electrons + H^-$

Maxwellian — Maxwellian



Slow Negative Charge Production Rate Coefficients for

He + H^+ -> electrons and H^-

Beam - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	150000.	200000.
1.0E+00	1.51E-09	5.59E-09	1.84E-08	3.11E-08	4.07E-08	4.18E-08	4.26E-08
2.0E+00	1.51E-09	5.80E-09	1.85E-08	3.32E-08	4.07E-08	4.18E-08	4.26E-08
4.0E+00	1.52E-09	5.81E-09	1.85E-08	3.33E-08	4.07E-08	4.18E-08	4.26E-08
7.0E+00	1.52E-09	5.81E-09	1.85E-08	3.32E-08	4.07E-08	4.18E-08	4.26E-08
1.0E+01	1.52E-09	5.81E-09	1.85E-08	3.32E-08	4.06E-08	4.18E-08	4.26E-08
2.0E+01	1.53E-09	5.81E-09	1.84E-08	3.32E-08	4.06E-08	4.18E-08	4.25E-08
4.0E+01	1.54E-09	5.82E-09	1.84E-08	3.31E-08	4.05E-08	4.18E-08	4.25E-08
7.0E+01	1.56E-09	5.85E-09	1.84E-08	3.31E-08	4.05E-08	4.18E-08	4.25E-08
1.0E+02	1.58E-09	5.87E-09	1.83E-08	3.30E-08	4.04E-08	4.18E-08	4.25E-08
2.0E+02	1.65E-09	5.97E-09	1.83E-08	3.30E-08	4.02E-08	4.18E-08	4.24E-08
4.0E+02	1.80E-09	6.17E-09	1.84E-08	3.28E-08	4.01E-08	4.18E-08	4.23E-08
7.0E+02	2.03E-09	6.48E-09	1.85E-08	3.28E-08	3.99E-08	4.18E-08	4.22E-08
1.0E+03	2.26E-09	6.80E-09	1.87E-08	3.27E-08	3.97E-08	4.18E-08	4.22E-08
2.0E+03	3.04E-09	7.83E-09	1.92E-08	3.27E-08	3.93E-08	4.18E-08	4.20E-08
4.0E+03	4.66E-09	9.70E-09	2.04E-08	3.26E-08	3.87E-08	4.17E-08	4.17E-08
7.0E+03	7.06E-09	1.21E-08	2.19E-08	3.25E-08	3.82E-08	4.13E-08	4.14E-08
1.0E+04	9.37E-09	1.43E-08	2.31E-08	3.25E-08	3.77E-08	4.09E-08	4.11E-08
2.0E+04	1.59E-08	1.98E-08	2.62E-08	3.29E-08	3.69E-08	3.98E-08	4.02E-08

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-39.3816	1.02698	.576534	.188719	4.425971E-05	-.0355917	-.0198253
20000.	-37.3350	.517958	.301257	.123942	.00937576	-.0100372	-.0129438
40000.	-35.4715	.129405	.0937138	.0477169	.0118588	-4.888201E-04	-.003254337E-03
70000.	-34.4703	.00355792	-.0123483	.0153783	-.00669156	.00817185	-.00392000
100000.	-34.0898	-.0436120	-.0204578	-.00631264	-9.902555E-04	3.700060E-04	5.429520E-04
150000.	-33.9953	-.0150052	-.0121264	-.00832720	-.00470324	-.00198547	-3.364408E-04
200000.	-33.9705	-.0219825	-.0119088	-.00528740	-.00226756	-.00102041	-4.501627E-04

See appendix for Chebyshev fit details.

$He + H^+ \rightarrow electrons + H^-$



Ionization and Total He^+ Production Cross Sections for

$H^+ + He \rightarrow H^+ + He^+ + e$
$H^+ + He \rightarrow (H + He^+)$ or $(H^+ + He^+ + e)$

Ionization		Total He^+ Production	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
1.0E+04	2.04E-17	5.0E+03	3.82E-17
1.5E+04	3.49E-17	7.0E+03	7.64E-17
2.0E+04	4.94E-17	1.0E+04	1.33E-16
3.0E+04	6.61E-17	2.0E+04	2.39E-16
4.0E+04	7.37E-17	2.3E+04	2.43E-16
7.0E+04	8.28E-17	4.0E+04	2.17E-16
9.0E+04	8.42E-17	7.0E+04	1.58E-16
1.0E+05	8.38E-17	1.0E+05	1.22E-16
1.5E+05	7.65E-17	2.0E+05	7.14E-17
2.0E+05	6.69E-17	4.0E+05	4.27E-17
4.0E+05	4.26E-17	7.0E+05	2.84E-17
7.0E+05	2.82E-17	1.0E+06	2.18E-17
1.0E+06	2.18E-17	1.5E+06	1.61E-17
1.5E+06	1.61E-17	2.0E+06	1.29E-17
2.0E+06	1.29E-17	4.0E+06	6.75E-18
3.0E+06	8.92E-18	5.0E+06	5.31E-18
4.0E+06	6.71E-18		
5.0E+06	5.32E-18		

<u>References:</u> 83, 87, 89, 315, 353, 354, 359, 368, 371, 372, 375, 376, 381, 382, 383, 384, 385, 386, 387, 388

<u>Accuracy:</u> 30%

<u>Notes:</u> (1) The total cross section for slow He^+ ion production is the sum of the electron capture cross ($H^+ + He \rightarrow H + He^+$) and the ionization cross section ($H^+ + He \rightarrow H^+ + H^+ + e$). (2) The cross section for production of He^{2+} is two orders of magnitude less than that for He^+ and contributes a negligible amount to the slow ion formation. (3) At energies less than 2×10^5 eV/amu the electron capture cross section is the dominant reaction in producing slow ions. Above this energy the cross section for He^+ formation is the same as the ionization cross section to form He^+. (4) For cross sections involving the formation of slow He^{2+} see the following set of data.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

Ionization E_{min} = 1.0E+04 eV/amu, E_{max} = 5.0E+06 eV/amu
Tot. He^+ Prod. E_{min} = 5.0E+03 eV/amu, E_{max} = 5.0E+06 eV/amu

	A0	A1	A2	A3	A4	A5	A6
Ionization	-76.3648	-.874453	-.884552	.221104	-.0292448	-.0234184	-.0152807
Tot. He^+ Prod.	-75.4062	-1.41315	-.923745	.441895	-.193224	-.00636635	.0315740

The fit represents the Ionization cross section with an rms deviation of 1.6%. The maximum deviation is 2.5% at 7.0E+04 eV/amu.

The fit represents the Tot. He^+ Prod. cross section with an rms deviation of 1.3%. The maximum deviation is 1.7% at 1.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$H^+ + He \rightarrow H^+ + He^+ + e$
$H^+ + He \rightarrow (H + He^+)$ or $(H^+ + He^+ + e)$



Ionization Rate Coefficients for

$H^+ + He \rightarrow H^+ + He^+ + e$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	Equal Temp.	He Temp. (eV) 5000.	6000.	7000.	8000.	10000.	15000.	20000.
1.0E+03	4.07E-12#	1.33E-10#	2.07E-10#	2.99E-10#	4.08E-10#	6.69E-10#	1.51E-09*	2.50E-09*
1.5E+03	5.51E-11#	2.99E-10#	4.08E-10#	5.31E-10#	6.69E-10#	9.78E-10#	1.89E-09*	2.92E-09*
2.0E+03	2.07E-10#	5.31E-10#	6.69E-10#	8.18E-10#	9.78E-10#	1.32E-09#	2.30E-09*	3.35E-09*
2.5E+03	4.68E-10#	8.18E-10#	9.78E-10#	1.15E-09#	1.32E-09#	1.70E-09*	2.71E-09*	3.77E-09*
3.0E+03	8.18E-10#	1.15E-09#	1.32E-09#	1.51E-09*	1.70E-09*	2.09E-09*	3.13E-09*	4.20E-09*
3.5E+03	1.24E-09#	1.51E-09*	1.70E-09*	1.89E-09*	2.09E-09*	2.50E-09*	3.56E-09*	4.62E-09*
4.0E+03	1.70E-09*	1.89E-09*	2.09E-09*	2.30E-09*	2.50E-09*	2.92E-09*	3.99E-09*	5.05E-09*
4.5E+03	2.20E-09*	2.30E-09*	2.50E-09*	2.71E-09*	2.92E-09*	3.35E-09*	4.41E-09*	5.47E-09*
5.0E+03	2.71E-09*	2.71E-09*	2.92E-09*	3.13E-09*	3.35E-09*	3.77E-09*	4.84E-09*	5.89E-09*
5.5E+03	3.24E-09*	3.13E-09*	3.35E-09*	3.56E-09*	3.77E-09*	4.20E-09*	5.26E-09*	6.30E-09*
6.0E+03	3.77E-09*	3.56E-09*	3.77E-09*	3.99E-09*	4.20E-09*	4.62E-09*	5.68E-09*	6.72E-09*
8.0E+03	5.89E-09*	5.26E-09*	5.47E-09*	5.68E-09*	5.89E-09*	6.30E-09*	7.32E-09	8.29E-09
1.0E+04	7.91E-09	6.92E-09*	7.12E-09*	7.32E-09	7.52E-09	7.91E-09	8.87E-09	9.79E-09
1.2E+04	9.79E-09	8.48E-09	8.68E-09	8.87E-09	9.06E-09	9.43E-09	1.03E-08	1.12E-08
1.4E+04	1.15E-08	9.97E-09	1.01E-08	1.03E-08	1.05E-08	1.08E-08	1.17E-08	1.25E-08
1.6E+04	1.31E-08	1.14E-08	1.15E-08	1.17E-08	1.19E-08	1.22E-08	1.30E-08	1.37E-08
1.8E+04	1.46E-08	1.27E-08	1.28E-08	1.30E-08	1.31E-08	1.34E-08	1.42E-08	1.49E-08
2.0E+04	1.60E-08	1.39E-08	1.40E-08	1.42E-08	1.43E-08	1.46E-08	1.53E-08	1.60E-08

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+03 eV, E_{max} = 2.0E+04 eV

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
5000.	-40.3451	2.34883	-.257537	-.0218607	.0131131	-.00170960	-2.72133E-04
6000.	-40.0282	2.14145	-.186832	-.0325594	.0117041	-7.73826E-04	-3.91919E-04
7000.	-39.7552	1.96806	-.132538	-.0385303	.00991961	-1.95606E-05	-4.64455E-04
8000.	-39.5164	1.82079	-.0901635	-.0415358	.00805275	5.25545E-04	-4.72996E-04
10000.	-39.1158	1.58360	-.0298040	-.0424986	.00463840	.00112196	-3.85570E-04
15000.	-38.3952	1.19217	.0455887	-.0348519	-.00101950	.00132410	-2.81514E-05
20000.	-37.8995	.951741	.0744487	-.0255952	-.00350613	8.71591E-04	-4.39401E-05
Equal Temp.	-41.9272	3.86757	-1.06890	.261511	-.0564975	.00926856	-.00119300

See appendix for Chebyshev fit details

$$H^+ + He \rightarrow H^+ + He^+ + e$$

Maxwellian — Maxwellian



Ionization Rate Coefficients for

$He + H^+ \rightarrow He^+ + H^+ + e$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	1.45E-09*	9.69E-09	2.05E-08	3.04E-08	3.68E-08	4.15E-08	3.55E-08
2.0E+00	1.47E-09*	9.68E-09	2.05E-08	3.04E-08	3.68E-08	4.15E-08	3.55E-08
4.0E+00	1.49E-09*	9.67E-09	2.05E-08	3.04E-08	3.68E-08	4.15E-08	3.55E-08
7.0E+00	1.52E-09*	9.66E-09	2.05E-08	3.04E-08	3.68E-08	4.15E-08	3.55E-08
1.0E+01	1.54E-09*	9.66E-09	2.04E-08	3.04E-08	3.68E-08	4.15E-08	3.55E-08
2.0E+01	1.60E-09*	9.65E-09	2.04E-08	3.04E-08	3.68E-08	4.15E-08	3.55E-08
4.0E+01	1.68E-09*	9.64E-09	2.04E-08	3.04E-08	3.67E-08	4.15E-08	3.55E-08
7.0E+01	1.77E-09*	9.64E-09	2.04E-08	3.04E-08	3.67E-08	4.15E-08	3.55E-08
1.0E+02	1.85E-09*	9.65E-09	2.04E-08	3.04E-08	3.67E-08	4.15E-08	3.55E-08
2.0E+02	2.08E-09*	9.71E-09	2.04E-08	3.04E-08	3.66E-08	4.14E-08	3.55E-08
4.0E+02	2.45E-09*	9.85E-09	2.05E-08	3.04E-08	3.66E-08	4.13E-08	3.55E-08
7.0E+02	2.91E-09*	1.01E-08	2.06E-08	3.04E-08	3.65E-08	4.13E-08	3.54E-08
1.0E+03	3.33E-09*	1.03E-08	2.06E-08	3.04E-08	3.65E-08	4.12E-08	3.55E-08
2.0E+03	4.53E-09*	1.10E-08	2.09E-08	3.05E-08	3.63E-08	4.11E-08	3.55E-08
4.0E+03	6.51E-09*	1.23E-08	2.15E-08	3.07E-08	3.62E-08	4.08E-08	3.55E-08
7.0E+03	8.99E-09	1.42E-08	2.25E-08	3.09E-08	3.60E-08	4.05E-08	3.54E-08
1.0E+04	1.12E-08	1.58E-08	2.34E-08	3.11E-08	3.58E-08	4.02E-08	3.54E-08
2.0E+04	1.70E-08	2.04E-08	2.59E-08	3.18E-08	3.55E-08	3.94E-08	3.53E-08

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-39.1279	1.16505	.469737	.0813519	-.0228287	-.0182847	-.00258180
20000.	-36.5805	.285887	.190126	.0869452	.0237093	4.34865E-04	-3.87928E-03
40000.	-35.3193	.0816359	.0597659	.0332395	.0137513	.00383099	1.22161E-04
70000.	-34.6011	.0145540	.0116645	.00621244	.00251084	6.93786E-04	4.47956E-04
100000.	-34.2559	-.0155100	-.00743994	-.00254453	-6.97240E-04	-1.61490E-04	2.33078E-05
200000.	-34.0175	-.0200973	-.0118588	-.00582167	-.00256998	-9.87558E-04	-2.32402E-04
500000.	-34.3105	-.00181505	-.00131097	-8.28718E-04	-5.05268E-04	-2.95214E-04	-1.01434E-04

See appendix for Chebyshev fit details

$He + H^+ \rightarrow He^+ + H^+ + e$



Total He+ Production Rate Coefficients for

$H^+ + He \rightarrow (H + He^+)$ or $(H^+ + He^+ + e)$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	Equal Temp.	He Temp. (eV) 1000.	2000.	4000.	7000.	10000.	15000.	20000.
1.0E+03	3.32E-10#	3.32E-10#	6.71E-10#	1.68E-09#	3.78E-09*	6.26E-09*	1.06E-08*	1.49E-08
1.5E+03	1.39E-09#	1.13E-09#	1.68E-09#	3.02E-09*	5.41E-09*	8.00E-09*	1.24E-08	1.65E-08
2.0E+03	3.02E-09*	2.32E-09*	3.02E-09*	4.58E-09*	7.12E-09*	9.77E-09*	1.41E-08	1.81E-08
2.5E+03	4.98E-09*	3.78E-09*	4.58E-09*	6.26E-09*	8.88E-09*	1.15E-08	1.57E-08	1.96E-08
3.0E+03	7.12E-09*	5.41E-09*	6.26E-09*	8.00E-09*	1.06E-08*	1.32E-08	1.73E-08	2.10E-08
3.5E+03	9.33E-09*	7.12E-09*	8.00E-09*	9.77E-09*	1.24E-08	1.49E-08	1.88E-08	2.24E-08
4.0E+03	1.15E-08	8.88E-09*	9.77E-09*	1.15E-08	1.41E-08	1.65E-08	2.03E-08	2.37E-08
4.5E+03	1.37E-08	1.06E-08*	1.15E-08	1.32E-08	1.57E-08	1.81E-08	2.17E-08	2.49E-08
5.0E+03	1.57E-08	1.24E-08	1.32E-08	1.49E-08	1.73E-08	1.96E-08	2.30E-08	2.61E-08
5.5E+03	1.77E-08	1.41E-08	1.49E-08	1.65E-08	1.88E-08	2.10E-08	2.43E-08	2.72E-08
6.0E+03	1.96E-08	1.57E-08	1.65E-08	1.81E-08	2.03E-08	2.24E-08	2.55E-08	2.83E-08
8.0E+03	2.61E-08	2.17E-08	2.24E-08	2.37E-08	2.55E-08	2.72E-08	2.98E-08	3.20E-08
1.0E+04	3.12E-08	2.67E-08	2.72E-08	2.83E-08	2.98E-08	3.12E-08	3.32E-08	3.51E-08
1.2E+04	3.51E-08	3.07E-08	3.12E-08	3.20E-08	3.32E-08	3.44E-08	3.61E-08	3.76E-08
1.4E+04	3.82E-08	3.40E-08	3.44E-08	3.51E-08	3.61E-08	3.70E-08	3.84E-08	3.97E-08
1.6E+04	4.06E-08	3.67E-08	3.70E-08	3.76E-08	3.84E-08	3.92E-08	4.04E-08	4.14E-08
1.8E+04	4.25E-08	3.90E-08	3.92E-08	3.97E-08	4.04E-08	4.10E-08	4.19E-08	4.28E-08
2.0E+04	4.40E-08	4.08E-08	4.10E-08	4.14E-08	4.19E-08	4.25E-08	4.33E-08	4.40E-08

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+03 eV, E_{max} = 2.0E+04 eV

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
1000.	-37.7759	2.35765	-.526776	.0461542	-.00599557	.00186274	5.61892E-05
2000.	-37.3613	2.05214	-.389347	.00372548	.00380397	3.22896E-04	1.92742E-07
4000.	-36.7649	1.63391	-.227162	-.0328342	.00799982	4.62850E-04	-3.57204E-04
7000.	-36.1737	1.24696	-.108095	-.0449948	.00515942	.00147386	-4.77249E-04
10000.	-35.7678	.998971	-.0496155	-.0432215	.00174125	.00180493	-2.03326E-04
15000.	-35.3028	.734516	-.00387920	-.0346346	-.00171864	.00144813	-1.44086E-05
20000.	-34.9844	.566598	.0150211	-.0267530	-.00313077	.00108401	2.07674E-04
Equal Temp.	-37.4591	2.34257	-.631509	.0945062	-.0203712	.00642863	-.00134327

See appendix for Chebyshev fit details

$H^+ + He \rightarrow (H + He^+)$ or $(H^+ + He^+ + e)$



Total He^+ Production Rate Coefficients for

$He + H^+ \rightarrow (He^+ + H)$ or $(He^+ + H^+ + e)$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	1.84E-08	4.68E-08	6.02E-08	5.80E-08	5.36E-08	4.44E-08	3.56E-08
2.0E+00	1.84E-08	4.68E-08	6.02E-08	5.80E-08	5.36E-08	4.44E-08	3.56E-08
4.0E+00	1.84E-08	4.67E-08	6.02E-08	5.80E-08	5.36E-08	4.44E-08	3.56E-08
7.0E+00	1.83E-08	4.66E-08	6.01E-08	5.80E-08	5.36E-08	4.44E-08	3.56E-08
1.0E+01	1.83E-08	4.66E-08	6.01E-08	5.80E-08	5.36E-08	4.44E-08	3.56E-08
2.0E+01	1.83E-08	4.64E-08	6.00E-08	5.80E-08	5.36E-08	4.44E-08	3.56E-08
4.0E+01	1.83E-08	4.62E-08	5.99E-08	5.79E-08	5.36E-08	4.44E-08	3.56E-08
7.0E+01	1.83E-08	4.60E-08	5.98E-08	5.79E-08	5.35E-08	4.44E-08	3.56E-08
1.0E+02	1.84E-08	4.59E-08	5.97E-08	5.79E-08	5.35E-08	4.44E-08	3.56E-08
2.0E+02	1.87E-08	4.55E-08	5.94E-08	5.78E-08	5.35E-08	4.44E-08	3.56E-08
4.0E+02	1.95E-08	4.51E-08	5.91E-08	5.77E-08	5.35E-08	4.43E-08	3.56E-08
7.0E+02	2.07E-08	4.48E-08	5.87E-08	5.75E-08	5.34E-08	4.44E-08	3.56E-08
1.0E+03	2.18E-08	4.46E-08	5.85E-08	5.74E-08	5.34E-08	4.44E-08	3.56E-08
2.0E+03	2.51E-08	4.44E-08	5.76E-08	5.71E-08	5.32E-08	4.44E-08	3.56E-08
4.0E+03	3.00E-08	4.47E-08	5.62E-08	5.64E-08	5.30E-08	4.44E-08	3.56E-08
7.0E+03	3.50E-08	4.56E-08	5.46E-08	5.54E-08	5.25E-08	4.44E-08	3.56E-08
1.0E+04	3.86E-08	4.64E-08	5.35E-08	5.45E-08	5.21E-08	4.44E-08	3.56E-08
2.0E+04	4.53E-08	4.85E-08	5.16E-08	5.21E-08	5.05E-08	4.42E-08	3.56E-08

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-35.1311	.412881	.228878	.0583838	-.0173875	-.0216039	-.00620616
20000.	-33.7784	-.00396664	.0221013	.0208873	.00921146	.00114081	-.00146388
40000.	-33.3338	-.0658860	-.0346993	-.0127985	-.00242907	.00117581	.00179130
70000.	-33.3704	-.0391288	-.0249648	-.0133215	-.00605125	-.00211349	-2.49745E-04
100000.	-33.5061	-.0193669	-.0135261	-.00817621	-.00432298	-.00189118	-6.09453E-04
200000.	-33.8626	-6.29120E-04	-9.61867E-04	-9.42621E-04	-7.21387E-04	-4.47888E-04	-2.11683E-04
500000.	-34.3009	-7.64617E-04	-4.54422E-04	-2.24604E-04	-1.49745E-04	-1.31679E-04	-5.40513E-05

See appendix for Chebyshev fit details

$He + H^+ \rightarrow (He^+ + H)$ or $(He^+ + H^+ + e)$

Beam – Maxwellian



Ionization and Total He^{2+} Production Cross Sections for

$H^+ + He \rightarrow H^+ + He^{2+} + 2e$
$H^+ + He \rightarrow (H^- + He^{2+})$ or $(H^+ + He^{2+} + 2e)$

Ionization		Total He^{2+} Production	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
2.0E+04	1.40E-19	1.0E+04	4.97E-19
3.0E+04	4.46E-19	1.5E+04	9.85E-19
4.0E+04	6.68E-19	2.0E+04	1.49E-18
6.0E+04	9.69E-19	3.0E+04	2.17E-18
8.0E+04	1.07E-18	4.0E+04	2.38E-18
1.0E+05	9.95E-19	5.0E+04	2.41E-18
1.5E+05	7.42E-19	7.0E+04	2.30E-18
2.0E+05	5.59E-19	1.0E+05	1.80E-18
3.0E+05	3.67E-19	1.5E+05	1.04E-18
4.0E+05	2.52E-19	2.0E+05	6.71E-19
6.0E+05	1.38E-19	3.0E+05	3.83E-19
8.0E+05	9.63E-20	4.0E+05	2.52E-19
1.0E+06	7.49E-20	7.0E+05	1.14E-19
1.5E+06	4.95E-20	1.0E+06	7.54E-20
2.0E+06	3.72E-20	1.5E+06	4.98E-20
3.0E+06	2.46E-20	2.0E+06	3.74E-20
4.0E+06	1.85E-20	4.0E+06	1.86E-20
5.0E+06	1.46E-20	5.0E+06	1.48E-20

References: 83, 87, 89, 315, 353, 354, 359, 368, 371, 372, 375, 376, 381, 382, 383, 384, 385, 386, 387, 388

Accuracy: 20%

Notes: (1) The total cross section for producing slow He^{2+} ions is the sum of the double ionization cross section ($H^+ + He \rightarrow H^+ + He^{2+} + 2e$) and the electron capture cross section ($H^+ + He \rightarrow H^- + He^{2+}$). (2) At energies less than 3×10^5 eV/amu the production of He^{2+} is dominated by double electron capture collisions. (3) For cross sections involving the formation of He^+ see the preceding set of data.

Chebyshev Fitting Parameters for Cross Sections

Ionization E_{min} = 2.0E+04 eV/amu, E_{max} = 5.0E+06 eV/amu
Tot. He^{2+} Prod. E_{min} = 1.0E+04 eV/amu, E_{max} = 5.0E+06 eV/amu

	A0	A1	A2	A3	A4	A5	A6
Ionization	-86.8125	-1.69704	-.988419	.579481	-.140358	-.0229323	-.00915691
Tot. He^{2+} Prod.	-85.7168	-2.26739	-.987093	.602380	-.0759791	-.104008	.0149516

The fit represents the Ionization cross section with an rms deviation of 2.5%. The maximum deviation is 4.3% at 4.0E+04 eV/amu.

The fit represents the Tot. He^{2+} Prod. cross section with an rms deviation of 2.7%. The maximum deviation is 4.8% at 2.0E+05 eV/amu.

See appendix for Chebyshev fit details.

$H^+ + He \rightarrow H^+ + He^{2+} + 2e$
$H^+ + He \rightarrow (H^- + He^{2+})$ or $(H^+ + He^{2+} + 2e)$



Double Ionization Rate Coefficients for

$He + H^+ \rightarrow He^{2+} + H^+ + 2e$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	He Energy (eV/amu) 20000.	40000.	60000.	80000.	100000.	200000.	500000.
1.0E+00	1.42E-11*	1.85E-10	3.29E-10	4.20E-10	4.37E-10	3.47E-10	1.78E-10
2.0E+00	1.44E-11*	1.85E-10	3.29E-10	4.20E-10	4.37E-10	3.47E-10	1.78E-10
4.0E+00	1.46E-11*	1.85E-10	3.29E-10	4.19E-10	4.36E-10	3.47E-10	1.78E-10
7.0E+00	1.49E-11*	1.85E-10	3.29E-10	4.19E-10	4.36E-10	3.47E-10	1.78E-10
1.0E+01	1.52E-11*	1.85E-10	3.28E-10	4.19E-10	4.36E-10	3.47E-10	1.78E-10
2.0E+01	1.58E-11*	1.85E-10	3.28E-10	4.18E-10	4.36E-10	3.47E-10	1.78E-10
4.0E+01	1.68E-11*	1.85E-10	3.27E-10	4.17E-10	4.35E-10	3.47E-10	1.78E-10
7.0E+01	1.80E-11*	1.85E-10	3.27E-10	4.16E-10	4.34E-10	3.47E-10	1.78E-10
1.0E+02	1.90E-11*	1.85E-10	3.26E-10	4.15E-10	4.34E-10	3.47E-10	1.78E-10
2.0E+02	2.20E-11*	1.85E-10	3.26E-10	4.13E-10	4.32E-10	3.47E-10	1.78E-10
4.0E+02	2.70E-11*	1.87E-10	3.25E-10	4.10E-10	4.31E-10	3.46E-10	1.78E-10
7.0E+02	3.34E-11*	1.89E-10	3.25E-10	4.07E-10	4.28E-10	3.47E-10	1.78E-10
1.0E+03	3.90E-11*	1.92E-10	3.25E-10	4.04E-10	4.26E-10	3.46E-10	1.78E-10
2.0E+03	5.50E-11*	2.00E-10	3.24E-10	3.97E-10	4.19E-10	3.46E-10	1.78E-10
4.0E+03	8.11E-11*	2.15E-10	3.23E-10	3.85E-10	4.08E-10	3.45E-10	1.78E-10
7.0E+03	1.13E-10	2.32E-10	3.21E-10	3.73E-10	3.94E-10	3.42E-10	1.79E-10
1.0E+04	1.40E-10	2.44E-10	3.20E-10	3.64E-10	3.83E-10	3.39E-10	1.79E-10
2.0E+04	2.03E-10	2.71E-10	3.18E-10	3.45E-10	3.58E-10	3.28E-10	1.80E-10

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
20000.	-48.1754	1.30300	.485174	.0480600	-.0457483	-.0219390	-7.65210E-04
40000.	-44.6402	.155683	.100721	.0406747	.00551141	-.00485389	-.00420546
60000.	-43.6938	.0163921	-.00478529	-.00179360	-.00131646	-1.05431E-04	8.93568E-04
80000.	-43.2865	-.0824386	-.0435066	-.0165975	-.00387115	5.14172E-04	1.18040E-03
100000.	-43.1958	-.0767199	-.0463123	-.0215537	-.00727208	-9.94415E-04	8.20812E-04
200000.	-43.5811	-.0170301	-.0127181	-.00866682	-.00530410	-.00278153	-1.19855E-03
500000.	-44.8971	.00488974	.00333081	.00168204	5.29597E-04	-3.67440E-06	-7.72152E-05

See appendix for Chebyshev fit details

$He + H^+ \rightarrow He^{2+} + H^+ + 2e$

Beam – Maxwellian



Total He^{2+} Production Rate Coefficients for

$H^+ + He \rightarrow (H^- + He^{2+}) + (H^+ + He^{2+} + 2e)$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	Equal Temp.	He Temp. (eV) 5000.	7500.	10000.	12500.	15000.	17500.	20000.
7.5E+02	7.68E-15#	2.03E-12#	6.81E-12#	1.47E-11#	2.53E-11#	3.80E-11*	5.22E-11*	6.75E-11*
1.0E+03	1.04E-13#	3.55E-12#	9.61E-12#	1.87E-11#	3.01E-11#	4.35E-11*	5.82E-11*	7.38E-11*
1.3E+03	5.03E-13#	5.59E-12#	1.29E-11#	2.30E-11#	3.53E-11*	4.92E-11*	6.44E-11*	8.03E-11*
1.5E+03	1.45E-12#	8.15E-12#	1.66E-11#	2.77E-11#	4.07E-11*	5.52E-11*	7.07E-11*	8.68E-11*
1.8E+03	3.12E-12#	1.12E-11#	2.08E-11#	3.27E-11#	4.63E-11*	6.12E-11*	7.70E-11*	9.33E-11*
2.0E+03	5.59E-12#	1.47E-11#	2.53E-11#	3.80E-11*	5.22E-11*	6.75E-11*	8.35E-11*	1.00E-10*
3.0E+03	2.30E-11#	3.27E-11#	4.63E-11*	6.12E-11*	7.70E-11*	9.33E-11*	1.10E-10*	1.27E-10*
4.0E+03	4.92E-11*	5.52E-11*	7.07E-11*	8.68E-11*	1.03E-10*	1.20E-10*	1.37E-10*	1.54E-10*
5.0E+03	8.03E-11*	8.03E-11*	9.66E-11*	1.13E-10*	1.30E-10*	1.47E-10*	1.63E-10*	1.80E-10*
6.0E+03	1.13E-10*	1.07E-10*	1.23E-10*	1.40E-10*	1.57E-10*	1.73E-10*	1.89E-10*	2.05E-10
7.0E+03	1.47E-10*	1.33E-10*	1.50E-10*	1.67E-10*	1.83E-10*	1.99E-10	2.14E-10	2.30E-10
8.0E+03	1.80E-10*	1.60E-10*	1.76E-10*	1.92E-10*	2.08E-10	2.24E-10	2.38E-10	2.53E-10
9.0E+03	2.11E-10	1.86E-10*	2.02E-10	2.18E-10	2.33E-10	2.47E-10	2.61E-10	2.75E-10
1.0E+04	2.41E-10	2.11E-10	2.27E-10	2.41E-10	2.56E-10	2.70E-10	2.83E-10	2.96E-10
1.2E+04	2.96E-10	2.59E-10	2.72E-10	2.86E-10	2.99E-10	3.12E-10	3.24E-10	3.35E-10
1.5E+04	3.66E-10	3.21E-10	3.33E-10	3.44E-10	3.55E-10	3.66E-10	3.76E-10	3.86E-10
2.0E+04	4.50E-10	4.04E-10	4.12E-10	4.21E-10	4.29E-10	4.36E-10	4.44E-10	4.50E-10

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 7.5E+02 eV, E_{max} = 2.0E+04 eV

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
5000.	-47.9855	2.71190	-.299562	-.0631734	.0152650	-.00140966	-4.92425E-04
7500.	-47.0952	2.12886	-.119977	-.0784054	.00701306	.00151172	-6.61029E-04
10000.	-46.4732	1.74892	-.0292041	-.0751534	3.23238E-04	.00245385E-03	-3.15620E-04
12500.	-46.0040	1.47944	.0203814	-.0667770	-.00376242	.00232506	5.55977E-05
15000.	-45.6322	1.27684	.0490457	-.0581142	-.00655132	.00195763	1.43604E-04
17500.	-45.3276	1.11863	.0651645	-.0499554	-.00809607	.00145805	3.73270E-04
20000.	-45.0718	.991107	.0746366	-.0430706	-.00880789	.00106692	2.57551E-04
Equal Temp.	-50.6204	5.07900	-1.59808	.393560	-.101018	.0175307	-.00147330

See appendix for Chebyshev fit details

$$H^+ + He \rightarrow (H^- + He^{2+}) + (H^+ + He^{2+} + 2e)$$

Maxwellian — Maxwellian



Total He^{2+} Production Rate Coefficients for

$He + H^+ \rightarrow (He^{2+} + H^-) + (He^{2+} + H^+ + 2e)$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	150000.	200000.
1.0E+00	3.56E-11*	2.92E-10	6.61E-10	8.44E-10	7.90E-10	5.59E-10	4.17E-10
2.0E+00	3.60E-11*	2.92E-10	6.61E-10	8.44E-10	7.89E-10	5.59E-10	4.17E-10
4.0E+00	3.67E-11*	2.92E-10	6.60E-10	8.43E-10	7.89E-10	5.59E-10	4.17E-10
7.0E+00	3.74E-11*	2.92E-10	6.60E-10	8.43E-10	7.88E-10	5.59E-10	4.17E-10
1.0E+01	3.80E-11*	2.91E-10	6.60E-10	8.42E-10	7.88E-10	5.59E-10	4.17E-10
2.0E+01	3.96E-11*	2.91E-10	6.59E-10	8.41E-10	7.86E-10	5.59E-10	4.17E-10
4.0E+01	4.20E-11*	2.91E-10	6.59E-10	8.39E-10	7.85E-10	5.59E-10	4.17E-10
7.0E+01	4.49E-11*	2.91E-10	6.58E-10	8.37E-10	7.83E-10	5.58E-10	4.17E-10
1.0E+02	4.73E-11*	2.92E-10	6.58E-10	8.36E-10	7.81E-10	5.58E-10	4.18E-10
2.0E+02	5.41E-11*	2.95E-10	6.57E-10	8.32E-10	7.78E-10	5.58E-10	4.18E-10
4.0E+02	6.55E-11*	3.01E-10	6.56E-10	8.26E-10	7.72E-10	5.57E-10	4.19E-10
7.0E+02	8.03E-11*	3.10E-10	6.55E-10	8.20E-10	7.66E-10	5.57E-10	4.19E-10
1.0E+03	9.37E-11*	3.18E-10	6.53E-10	8.15E-10	7.62E-10	5.57E-10	4.20E-10
2.0E+03	1.33E-10*	3.43E-10	6.50E-10	8.00E-10	7.48E-10	5.56E-10	4.22E-10
4.0E+03	1.99E-10	3.84E-10	6.46E-10	7.73E-10	7.26E-10	5.54E-10	4.24E-10
7.0E+03	2.77E-10	4.32E-10	6.41E-10	7.40E-10	6.98E-10	5.49E-10	4.27E-10
1.0E+04	3.39E-10	4.69E-10	6.36E-10	7.12E-10	6.74E-10	5.42E-10	4.28E-10
2.0E+04	4.75E-10	5.41E-10	6.21E-10	6.47E-10	6.14E-10	5.17E-10	4.24E-10

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

<u>Chebyshev Fitting Parameters for Rate Coefficients</u>

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-46.3757	1.27650	.480627	.0489252	-.0511164	-.0308841	-.00542828
20000.	-43.5984	.264368	.158736	.0549494	.00162732	-.00969406	-.00629046
40000.	-42.3060	-.0238894	-.0125884	-.00618733	-.00280784	-.00102524	-4.84730E-04
70000.	-41.9087	-.101626	-.0604596	-.0289603	-.0112319	-.00285512	3.95976E-04
100000.	-42.0360	-.0958542	-.0558944	-.0266235	-.0108604	-.00350631	-4.83059E-04
150000.	-42.6352	-.0233424	-.0167110	-.0116669	-.00760143	-.00419389	-.00181500
200000.	-43.1811	.0113231	.00418441	-8.73051E-04	-.00251835	-.00215399	-.00136312

See appendix for Chebyshev fit details

$He + H^+ \rightarrow (He^{2+} + H^-)$ or $(He^{2+} + H^+ + 2e)$



Penning, Associative, and Total Ionization Cross Sections for

$He(2^1S) + H \rightarrow He + H^+ + e$ (σ_{PI})
$He(2^1S) + H \rightarrow HeH^+ + e$ (σ_{AI})
$He(2^1S) + H \rightarrow$ (σ_{PI}) + (σ_{AI}) (Total Ion.)

Total Ion. (TI)		Penn. Ion. (PI)		Assoc. Ion. (AI)	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
2.0E-02	7.93E-15	1.8E-02	6.02E-15	1.9E-02	1.83E-15
2.5E-02	7.35E-15	2.0E-02	5.89E-15	2.0E-02	1.77E-15
3.0E-02	6.89E-15	3.0E-02	5.34E-15	2.5E-02	1.51E-15
4.0E-02	6.19E-15	4.0E-02	4.93E-15	3.0E-02	1.29E-15
6.0E-02	5.16E-15	6.0E-02	4.29E-15	4.0E-02	9.78E-16
8.0E-02	4.51E-15	8.0E-02	3.83E-15	6.0E-02	5.79E-16
1.0E-01	4.02E-15	1.0E-01	3.47E-15	8.0E-02	3.75E-16
1.3E-01	3.49E-15	1.3E-01	3.11E-15	1.0E-01	2.60E-16
				1.3E-01	1.80E-16

<u>References:</u> 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433

<u>Accuracy:</u> 100%

<u>Notes:</u> (1) The total ionization of H by low energy $He(2^1S)$ is the sum of Penning ionization $He(2^1S) + H \rightarrow He + H^+ + e$ and associative ionization $He(2^1S) + H \rightarrow HeH^+ + e$. (2) All data have been normalized to theory at an interaction energy of 0.04 eV. (3) Total ionization is sometimes referred to as chemionization. (4) The energy of the 2^1S state is 20.61 eV with a lifetime of $2x10^{-2}$ sec.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

Total Ion. (TI)	E_{min} =	2.0E-02	eV/amu,	E_{max} =	1.3E-01	eV/amu
Penn. Ion. (PI)	E_{min} =	1.8E-02	eV/amu,	E_{max} =	1.3E-01	eV/amu
Assoc. Ion. (AI)	E_{min} =	1.9E-02	eV/amu,	E_{max} =	1.3E-01	eV/amu

	A0	A1	A2	A3	A4	A5	A6
Total Ion. (TI)	-65.7017	-.409875	-.0288546	.000799090	.000526085	-.00141688	.000701497
Penn. Ion. (PI)	-66.0843	-.332960	-.0347058	.00142222	.00272425	.00118710	.000264732
Assoc. Ion. (AI)	-69.9529	-1.18775	-.135804	.0232877	.0173897	.00454102	.000920812

The fit represents the (TI) cross section with an rms deviation of 0.1%. The maximum deviation is 0.2% at 3.0E-02 eV/amu.

The fit represents the (PI) cross section with an rms deviation of 0.1%. The maximum deviation is 0.1% at 6.0E-02 eV/amu.

The fit represents the (AI) cross section with an rms deviation of 0.3%. The maximum deviation is 0.4% at 3.0E-02 eV/amu.

See appendix for Chebyshev fit details.

$He(2^1S) + H \rightarrow He + H^+ + e \quad (\sigma_{PI})$
$He(2^1S) + H \rightarrow HeH^+ + e \quad (\sigma_{AI})$



Penning, Associative, and Total Ionization Cross Sections for

$He(2^3S) + H \rightarrow He + H^+ + e$ (σ_{PI})
$He(2^3S) + H \rightarrow HeH^+ + e$ (σ_{AI})
$He(2^3S) + H \rightarrow$ (σ_{PI}) + (σ_{AI}) (Total Ion.)

Total Ion. (TI)		Penn. Ion. (PI)		Assoc. Ion. (AI)	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
1.3E-02	4.26E-15	1.2E-02	2.95E-15	1.2E-02	1.34E-15
1.5E-02	4.12E-15	1.5E-02	2.80E-15	1.5E-02	1.25E-15
2.0E-02	3.84E-15	2.0E-02	2.61E-15	2.0E-02	1.13E-15
4.0E-02	3.05E-15	4.0E-02	2.20E-15	4.0E-02	8.08E-16
6.0E-02	2.55E-15	6.0E-02	1.95E-15	6.0E-02	6.14E-16
8.0E-02	2.24E-15	8.0E-02	1.77E-15	8.0E-02	4.90E-16
1.0E-01	2.02E-15	1.0E-01	1.61E-15	1.0E-01	3.99E-16
1.5E-01	1.64E-15	1.5E-01	1.36E-15	1.5E-01	2.67E-16
2.0E-01	1.40E-15	2.0E-01	1.19E-15	2.0E-01	1.95E-16
4.0E-01	9.79E-16	3.0E-01	9.77E-16	4.0E-01	9.18E-17
6.0E-01	7.72E-16	5.0E-01	7.58E-16	6.0E-01	5.98E-17
8.0E-01	6.31E-16	7.0E-01	6.13E-16	8.0E-01	4.54E-17
1.0E+00	5.25E-16	1.0E+00	4.88E-16	1.0E+00	3.69E-17
1.5E+00	3.92E-16	1.5E+00	3.87E-16	1.5E+00	2.53E-17
2.0E+00	3.44E-16	2.0E+00	3.36E-16	2.0E+00	1.93E-17
4.0E+00	2.54E-16	4.0E+00	2.49E-16	4.0E+00	9.51E-18
6.0E+00	2.13E-16	6.0E+00	2.10E-16	6.0E+00	6.12E-18
8.0E+00	1.89E-16	8.0E+00	1.86E-16	8.0E+00	4.11E-18
1.0E+01	1.71E-16	1.0E+01	1.71E-16	1.0E+01	2.17E-18
1.3E+01	1.53E-16	1.3E+01	1.51E-16	1.3E+01	9.08E-19

<u>References:</u> 420, 421, 422, 423, 424, 423, 425, 426, 427, 428, 429, 430, 431, 432, 433

<u>Accuracy:</u> 100%

<u>Notes:</u> (1) The total ionization of H by low energy metastable $He(2^3S)$ is the sum of the cross section for Penning ionization σ_{PI} - $He(2^3S) + H \rightarrow He + H^+ + e$ and associative ionization σ_{AI} - $He(2^3S) + H \rightarrow HeH^+ + e$. (2) Much of the data have been taken such that the entire metastable beam is in the 2^3S state. Measurements have shown that the ratio of singlet to triplet states in ion beams is approximately 0.1. This assumption results in an error well within the experimental errors. (3) Total ionization is also referred to as chemionization. (4) The energy level of the 2^3S is 19.82 eV with a lifetime of 7900 sec.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

Total Ion. (TI) E_{min} = 1.3E-02 eV/amu, E_{max} = 1.3E+01 eV/amu
Penn. Ion. (PI) E_{min} = 1.2E-02 eV/amu, E_{max} = 1.3E+01 eV/amu
Assoc. Ion. (AI) E_{min} = 1.2E-02 eV/amu, E_{max} = 1.3E+01 eV/amu

	A0	A1	A2	A3	A4	A5	A6
Total Ion. (TI)	-69.3748	-1.75540	-.0609031	.106912	.0109231	-.0140956	-.0257512
Penn. Ion. (PI)	-69.6823	-1.55608	-.107728	.0932212	.0140800	-.0224249	-.0110671
Assoc. Ion. (AI)	-74.3566	-3.45306	-.466805	-.0598336	-.211902	-.116545	-.0298545

The fit represents the (TI) cross section with an rms deviation of 1.9%. The maximum deviation is 4.7% at 1.5E+00 eV/amu.
The fit represents the (PI) cross section with an rms deviation of 0.9%. The maximum deviation is 1.5% at 1.5E+00 eV/amu.
The fit represents the (AI) cross section with an rms deviation of 3.2%. The maximum deviation is 8.6% at 4.0E+00 eV/amu.

See appendix for Chebyshev fit details.

$He(2^3S) + H \rightarrow He + H^+ + e \quad (\sigma_{PI})$
$He(2^3S) + H \rightarrow HeH^+ + e \quad (\sigma_{AI})$



Total, Penning, and Rearrangement Ionization Cross Sections for $He(2^3S) + H_2$ -> Ion Formation

Total Ionization (TI)		Rearrangement Ionization (RI)	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
2.0E-02	6.19E-17	3.0E-02	1.12E-17
3.0E-02	1.02E-16	4.0E-02	1.22E-17
4.0E-02	1.45E-16	5.0E-02	1.37E-17
5.0E-02	1.89E-16	6.0E-02	1.55E-17
6.0E-02	2.36E-16	7.0E-02	1.74E-17
7.0E-02	2.84E-16	8.0E-02	1.92E-17
8.0E-02	3.34E-16	9.0E-02	2.06E-17
9.0E-02	3.86E-16	1.0E-01	2.19E-17
1.0E-01	4.39E-16	1.5E-01	2.79E-17
1.5E-01	7.21E-16	2.0E-01	3.35E-17
1.7E-01	8.04E-16	3.0E-01	4.47E-17
		4.0E-01	5.55E-17
		5.0E-01	6.58E-17
		6.0E-01	7.62E-17
		7.0E-01	8.70E-17
		7.5E-01	9.18E-17

References: 412, 421, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456

Accuracy: Unknown

Notes: (1) The measurement of metastable helium in H_2 is very difficult and the experimental results are inconsistent. Therefore the data should not be taken as absolute. The total ionization cross sections at thermal energies which are plotted at 0.03 eV/amu vary by a factor of six among eight measurements in glow discharges, flowing afterglow and merged beams. (2) The processes to form ions are: (a) Penning ionization (PI), $He(2^3S) + H_2 \rightarrow He + H^{2+} + e$; (b) associative ionization (AI) $He(2^3S) + H_2 \rightarrow HeH^+ + e$; (c) dissociative ionization (DI) $He(2^3S) + H_2 \rightarrow He + H^+ + H + e$; (d) Rearrangement ionization (RI) $He(2^3S) + H_2 \rightarrow HeH^+ + H + e$; (e) Ion pair production (IPP) $He(2^3S) + H_2 \rightarrow He + H^+ + H^-$. The sum of these processes is know as total ionization or chemionization; (3) Note that the total ionization is the same as Penning ionization within the experimental errors; (4) The energy level of the 2^3S is 19.82 eV with a lifetime of 7900 sec.

Chebyshev Fitting Parameters for Cross Sections

Total Ion. (TI) E_{min} = 2.0E-02 eV/amu, E_{max} = 1.7E-01 eV/amu
Rearrang. Ion. (RI) E_{min} = 3.0E-02 eV/amu, E_{max} = 7.5E-01 eV/amu

	A0	A1	A2	A3	A4	A5	A6
Total Ion. (TI)	-72.0362	1.29388	-.00614640	-.00568072	-.00984080	-.00595660	-.00475204
Rearrang. Ion. (RI)	-76.1234	1.06951	.0610251	-.000279924	.0164987	-.0176903	.00723031

The fit represents the (TI) cross section with an rms deviation of 0.2%. The maximum deviation is 0.3% at 5.0E-02 eV/amu.

The fit represents the (RI) cross section with an rms deviation of 0.5%. The maximum deviation is 0.9% at 6.0E-02 eV/amu.

See appendix for Chebyshev fit details.

$He(2^3S) + H_2 \rightarrow$ Ion Formation



Total, Penning, and Rearrangement Ionization Cross Sections for $He(2^1S) + H_2 \rightarrow$ Ion Formation

Total Ionization (TI) Energy (eV/amu)	Cross Section (cm^2)	Rearrangement Ionization (RI) Energy (eV/amu)	Cross Section (cm^2)
2.8E-02	9.97E-17	3.0E-02	1.19E-17
3.0E-02	1.11E-16	4.0E-02	1.47E-17
3.5E-02	1.35E-16	5.0E-02	1.69E-17
4.0E-02	1.58E-16	6.0E-02	1.90E-17
5.0E-02	2.04E-16	7.0E-02	2.08E-17
6.0E-02	2.50E-16	8.0E-02	2.26E-17
7.0E-02	2.94E-16	9.0E-02	2.41E-17
8.0E-02	3.40E-16	1.0E-01	2.51E-17
9.0E-02	3.77E-16	1.1E-01	2.55E-17
1.0E-01	4.07E-16	1.5E-01	2.19E-17
1.5E-01	5.00E-16	2.0E-01	1.83E-17
1.8E-01	5.39E-16	2.5E-01	1.57E-17
		3.0E-01	1.38E-17
		4.0E-01	1.12E-17
		5.0E-01	9.61E-18
		6.0E-01	8.38E-18
		7.0E-01	7.47E-18
		7.5E-01	7.12E-18

<u>References:</u> 412, 421, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456

<u>Accuracy:</u> Unknown

<u>Notes:</u> (1) The measurement of metastable helium in H_2 is very difficult and the results are inconsistent. Therefore, the data should not be taken as absolute. (2) The processes to form ions are: (a) Penning ionization (PI) $He(2^1S) + H_2 \rightarrow He + H_2^+ + e$; (b) Associative ionization (AI) $He(2^1S) + H_2 \rightarrow HeH_2^+ + e$; (c) Dissociative ionization (DI) $He(2^1S) + H_2 \rightarrow He + H^+ + H + e$; (d) Rearrangement ionization (RI) $He(2^1S) + H_2 \rightarrow HeH^+ + H + e$; (e) Ion pair production (IPP) $He(2^1S) + H_2 \rightarrow He + H^+ + H^-$. The sum of these processes is known as total ionization or chemionization. (3) Note that the Penning ionization and the total ionization are the same within experimental errors; (4) The energy of the 2^1S state is 20.61 eV with a lifetime of $2x10^{-2}$ sec.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

Total Ion. (TI) E_{min} = 2.8E-02 eV/amu, E_{max} = 1.8E-01 eV/amu
Rearrang. Ion. (RI) E_{min} = 3.0E-02 eV/amu, E_{max} = 7.5E-01 eV/amu

	A0	A1	A2	A3	A4	A5	A6
Total Ion. (TI)	-71.7445	.851023	-.128729	-.0202443	.00116383	.0124597	-.000179356
Rearrang. Ion. (RI)	-77.6786	-.311173	-.434644	.0826330	.0582759	-.0400365	-.0150564

The fit represents the (TI) cross section with an rms deviation of 0.3%. The maximum deviation is 0.6% at 7.0E-02 eV/amu.

The fit represents the (RI) cross section with an rms deviation of 2.1%. The maximum deviation is 2.9% at 7.0E-02 eV/amu.

See appendix for Chebyshev fit details.

$He(2^1S) + H_2 \rightarrow$ Ion Formation



Ionization and He^{2+} Production Cross Sections for

$H^+ + He^+ \rightarrow (H^+ + He^{2+} + e)$ (σ_1)
$H^+ + He^+ \rightarrow (H^+ + He^{2+} + e)$ or $(H^0 + He^{2+})$ $(\sigma_{He^{2+}})$

Ionization		He^{2+} Production	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
2.0E+04	4.20E-19	3.7E+03	9.25E-20
2.5E+04	7.50E-19	4.0E+03	1.08E-19
3.0E+04	1.18E-18	5.0E+03	1.62E-19
3.5E+04	1.67E-18	6.0E+03	2.41E-19
4.0E+04	2.21E-18	7.0E+03	3.55E-19
4.5E+04	2.80E-18	8.0E+03	5.35E-19
5.0E+04	3.42E-18	9.0E+03	7.92E-19
6.0E+04	4.69E-18	1.0E+04	1.12E-18
7.0E+04	5.92E-18	1.5E+04	4.31E-18
8.0E+04	6.98E-18	2.0E+04	9.33E-18
9.0E+04	8.00E-18	2.5E+04	1.44E-17
1.0E+05	8.97E-18	3.0E+04	1.89E-17
1.5E+05	1.09E-17	4.0E+04	2.52E-17
2.0E+05	1.01E-17	4.5E+04	2.71E-17
3.0E+05	7.74E-18	5.0E+04	2.86E-17
4.0E+05	6.36E-18	6.0E+04	2.96E-17
5.0E+05	5.19E-18	7.0E+04	2.93E-17
		8.0E+04	2.81E-17
		9.0E+04	2.64E-17
		1.0E+05	2.42E-17
		1.5E+05	1.75E-17
		2.0E+05	1.36E-17
		3.0E+05	8.97E-18
		4.0E+05	6.56E-18
		5.0E+05	5.12E-18

<u>References:</u> 69, 73, 75, 76, 77, 81, 278, 279, 458

<u>Accuracy:</u> Ionization - 60%
He^{2+} Production - 30%

<u>Notes:</u> (1) The production of He^{2+} arises from two processes: (a) Charge exchange ($H + He^+ \rightarrow H + He^{2+}$) and (b) Ionization ($H^+ + He^+ \rightarrow H^+ + He^{2+} + e$). (2) At energies less than approximately 2.0 - 3.0×10^5 eV/amu the charge exchange process is dominant. Above this energy the ionization reaction dominates.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

Ionization E_{min} = 2.0E+04 eV/amu, E_{max} = 5.0E+05 eV/amu
He^{2+} Production E_{min} = 3.7E+03 eV/amu, E_{max} = 5.0E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
Ionization	-80.4308	1.24048	-.890202	.00345922	.0570045	.0193037	-.00746147
He^{2+} Production	-80.5893	2.10007	-1.81896	-.00398887	.325128	-.125749	-.00793823

The fit represents the Ionization cross section with an rms deviation of 1.1%. The maximum deviation is 2.1% at 3.0E+05 eV/amu.

The fit represents the He^{2+} Production cross section with an rms deviation of 3.2%. The maximum deviation is 6.2% at 3.7E+03 eV/amu.

See appendix for Chebyshev fit details.

$$H^+ + He^+ \rightarrow (H^+ + He^{2+} + e) \text{ or } (H + He^{2+}) \; (\sigma_{He^{2+}})$$
$$H^+ + He^+ \rightarrow H^+ + He^{2+} + e \quad (\sigma_i)$$



Ionization Rate Coefficients for

$He^+ + H^+ \rightarrow He^{2+} + H^+ + e$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	He^+ Energy (eV/amu) 20000.	40000.	60000.	80000.	100000.	200000.	500000.
1.0E+00	4.25E-11*	6.14E-10	1.60E-09	2.70E-09	3.89E-09	6.27E-09	2.55E-09#
2.0E+00	4.30E-11*	6.14E-10	1.60E-09	2.74E-09	3.93E-09	6.27E-09	2.55E-09#
4.0E+00	4.37E-11*	6.14E-10	1.59E-09	2.74E-09	3.93E-09	6.27E-09	2.55E-09#
7.0E+00	4.46E-11*	6.14E-10	1.59E-09	2.74E-09	3.93E-09	6.27E-09	2.55E-09#
1.0E+01	4.52E-11*	6.15E-10	1.59E-09	2.74E-09	3.93E-09	6.26E-09	2.55E-09#
2.0E+01	4.70E-11*	6.16E-10	1.60E-09	2.74E-09	3.92E-09	6.26E-09	2.55E-09#
4.0E+01	4.98E-11*	6.18E-10	1.60E-09	2.74E-09	3.92E-09	6.25E-09	2.55E-09#
7.0E+01	5.30E-11*	6.21E-10	1.60E-09	2.75E-09	3.91E-09	6.25E-09	2.54E-09#
1.0E+02	5.57E-11*	6.24E-10	1.60E-09	2.75E-09	3.91E-09	6.24E-09	2.54E-09#
2.0E+02	6.36E-11*	6.35E-10	1.61E-09	2.76E-09	3.90E-09	6.23E-09	2.54E-09#
4.0E+02	7.68E-11*	6.56E-10	1.63E-09	2.78E-09	3.89E-09	6.21E-09	2.54E-09#
7.0E+02	9.48E-11*	6.87E-10	1.67E-09	2.81E-09	3.89E-09	6.19E-09	2.54E-09#
1.0E+03	1.12E-10*	7.17E-10	1.70E-09	2.83E-09	3.89E-09	6.18E-09	2.54E-09#
2.0E+03	1.69E-10*	8.17E-10	1.80E-09	2.89E-09	3.91E-09	6.13E-09	2.53E-09#
4.0E+03	2.88E-10	1.01E-09	1.98E-09	3.01E-09	3.96E-09	6.05E-09	2.52E-09#
7.0E+03	4.80E-10	1.27E-09	2.22E-09	3.17E-09	4.02E-09	5.95E-09	2.51E-09#
1.0E+04	6.80E-10	1.51E-09	2.42E-09	3.30E-09	4.07E-09	5.86E-09	2.49E-09#
2.0E+04	1.34E-09	2.16E-09	2.93E-09	3.62E-09	4.21E-09	5.60E-09	2.44E-09#

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

He^+ Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
20000.	-45.6943	1.57907	.710583	.175768	-.00705451	-.0284130	-.0139117
40000.	-41.8177	.512347	.311944	.126340	.0238811	-.00849678	-.00968659
60000.	-40.2324	.239690	.153320	.0661005	.0168874	-.00195984	-.00376827
80000.	-39.3054	.111744	.0654773	.0331289	.00699774	.00254954	-.00222073
100000.	-38.6923	.0227031	.0195819	.0149078	.00271827	.00119878	-.00147599
200000.	-37.8257	-.0418823	-.0248902	-.0123878	-.00550032	-.00208680	-5.276582E-04
500000.	-39.5936	-.0146717	-.00937139	-.00524336	-.00267631	-.00119132	-4.070685E-04

See appendix for Chebyshev fit details

$He^+ + H^+ \rightarrow He^{2+} + H^+ + e$

Beam – Maxwellian



He^{2+} Production Rate Coefficients for

$H^+ + He^+ \rightarrow (H^+ + He^{2+} + e)$ or $(H + He^{2+})$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	Equal Temp.	He^+ Temp. (eV) 1000.	2000.	4000.	7000.	10000.	15000.	20000.
3.0E+02	1.95E-15#	4.34E-14#	3.74E-13#	3.02E-12#	1.65E-11*	4.93E-11	1.62E-10	3.47E-10
5.0E+02	9.77E-14#	2.70E-13#	1.04E-12#	5.22E-12*	2.30E-11*	6.23E-11	1.87E-10	3.83E-10
7.0E+02	5.74E-13#	8.33E-13#	2.21E-12#	8.32E-12*	3.11E-11*	7.71E-11	2.14E-10	4.20E-10
9.0E+02	1.70E-12#	1.86E-12#	4.02E-12*	1.25E-11*	4.08E-11	9.39E-11	2.42E-10	4.59E-10
1.0E+03	2.59E-12#	2.59E-12#	5.22E-12*	1.51E-11*	4.64E-11	1.03E-10	2.57E-10	4.78E-10
1.3E+03	6.08E-12*	5.22E-12*	9.26E-12*	2.30E-11*	6.23E-11	1.28E-10	2.97E-10	5.29E-10
1.5E+03	1.19E-11*	9.26E-12*	1.51E-11*	3.34E-11	8.11E-11	1.56E-10	3.39E-10	5.82E-10
1.8E+03	2.08E-11*	1.51E-11*	2.30E-11*	4.64E-11	1.03E-10	1.87E-10	3.83E-10	6.36E-10
2.0E+03	3.34E-11	2.30E-11*	3.34E-11	6.23E-11	1.28E-10	2.21E-10	4.30E-10	6.92E-10
4.0E+03	2.97E-10	1.87E-10	2.21E-10	2.97E-10	4.30E-10	5.82E-10	8.68E-10	1.18E-09
6.0E+03	8.08E-10	5.29E-10	5.82E-10	6.92E-10	8.68E-10	1.05E-09	1.38E-09	1.72E-09
8.0E+03	1.45E-09	9.92E-10	1.05E-09	1.18E-09	1.38E-09	1.58E-09	1.92E-09	2.25E-09
1.0E+04	2.12E-09	1.51E-09	1.58E-09	1.72E-09	1.92E-09	2.12E-09	2.45E-09	2.77E-09
1.2E+04	2.77E-09	2.05E-09	2.12E-09	2.25E-09	2.45E-09	2.65E-09	2.96E-09	3.27E-09
1.4E+04	3.38E-09	2.58E-09	2.65E-09	2.77E-09	2.96E-09	3.14E-09	3.44E-09	3.72E-09
1.6E+04	3.94E-09	3.08E-09	3.14E-09	3.27E-09	3.44E-09	3.61E-09	3.89E-09	4.15E-09
1.8E+04	4.44E-09	3.56E-09	3.61E-09	3.72E-09	3.89E-09	4.04E-09	4.30E-09	4.54E-09
2.0E+04	4.89E-09	3.99E-09	4.04E-09	4.15E-09	4.30E-09	4.44E-09	4.67E-09	4.89E-09

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 3.0E+02 eV, E_{max} = 2.0E+04 eV

He^+ Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
1000.	-48.8313	5.83309	-.599670	-.148905	-.0411485	.0351803	.00376359
2000.	-47.5113	4.87892	-.205206	-.261188	-.00849826	.0291892	-3.733670E-04
4000.	-45.9304	3.85019	.0664286	-.260544	-.0126575	.0232481	-8.778358E-04
7000.	-44.4276	2.96296	.192841	-.194959	-.0273665	.0145816	.00158914
10000.	-43.3804	2.38527	.238913	-.143127	-.0334989	.00836597	.00280697
15000.	-42.1743	1.76616	.255545	-.0870174	-.0337594	.00196449	.00271327
20000.	-41.3491	1.37656	.243899	-.0536024	-.0298721	-.00120071	.00208341
Equal Temp.	-49.5753	7.04496	-1.51108	.221849	-.184942	.102821	-.0193609

See appendix for Chebyshev fit details.

$H^+ + He^+ \rightarrow (H^+ + He^{2+} + e)$ or $(H + He^{2+})$

Maxwellian — Maxwellian



He^{2+} Production Rate Coefficients for

$He^{+} + H^{+} \rightarrow (He^{2+} + H^{+} + e)$ or $(He^{2+} + H)$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	He^+ Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	1.56E-10	1.83E-09	6.99E-09	1.08E-08	1.06E-08	8.45E-09	2.51E-09#
2.0E+00	1.56E-10	1.83E-09	6.99E-09	1.08E-08	1.06E-08	8.45E-09	2.51E-09#
4.0E+00	1.57E-10	1.83E-09	6.99E-09	1.08E-08	1.06E-08	8.44E-09	2.51E-09#
7.0E+00	1.57E-10	1.82E-09	6.98E-09	1.08E-08	1.06E-08	8.44E-09	2.51E-09#
1.0E+01	1.58E-10	1.82E-09	6.98E-09	1.08E-08	1.06E-08	8.44E-09	2.52E-09#
2.0E+01	1.61E-10	1.83E-09	6.98E-09	1.07E-08	1.06E-08	8.44E-09	2.52E-09#
4.0E+01	1.66E-10	1.83E-09	6.97E-09	1.07E-08	1.06E-08	8.44E-09	2.52E-09#
7.0E+01	1.74E-10	1.85E-09	6.97E-09	1.07E-08	1.06E-08	8.43E-09	2.52E-09#
1.0E+02	1.83E-10	1.86E-09	6.97E-09	1.07E-08	1.06E-08	8.43E-09	2.52E-09#
2.0E+02	2.12E-10	1.92E-09	6.97E-09	1.07E-08	1.06E-08	8.42E-09	2.52E-09#
4.0E+02	2.73E-10	2.02E-09	6.99E-09	1.06E-08	1.06E-08	8.41E-09	2.52E-09#
7.0E+02	3.69E-10	2.18E-09	7.01E-09	1.06E-08	1.06E-08	8.40E-09	2.53E-09#
1.0E+03	4.69E-10	2.32E-09	7.03E-09	1.05E-08	1.06E-08	8.40E-09	2.53E-09#
2.0E+03	8.12E-10	2.76E-09	7.11E-09	1.03E-08	1.05E-08	8.37E-09	2.53E-09#
4.0E+03	1.49E-09	3.47E-09	7.24E-09	1.00E-08	1.03E-08	8.33E-09	2.54E-09#
7.0E+03	2.42E-09	4.29E-09	7.40E-09	9.66E-09	1.00E-08	8.29E-09	2.55E-09#
1.0E+04	3.24E-09	4.92E-09	7.52E-09	9.41E-09	9.77E-09	8.24E-09	2.56E-09#
2.0E+04	5.23E-09	6.29E-09	7.81E-09	8.92E-09	9.18E-09	8.01E-09	2.58E-09#

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

He^+ Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-42.9783	1.71856	.769214	.0945643	-.105175	-.0612282	-1.282705E-04
20000.	-39.5695	.556768	.303458	.0843952	-.0122473	-.0224889	-.00919993
40000.	-37.5112	.0421662	.0299866	.0131929	.00299530	-2.171997E-04	-6.745161E-04
70000.	-36.7855	-.0761446	-.0461124	-.0196107	-.00417912	.00142605	.00181386
100000.	-36.7766	-.0515173	-.0368252	-.0201855	-.00804421	-.00194056	1.214410E-04
200000.	-37.2003	-.0177798	-.0110628	-.00629544	-.00355051	-.00193013	-.00100821
500000.	-39.5894	.0104222	.00514715	.00174490	3.638925E-04	-8.415182E-06	-5.520857E-05

See appendix for Chebyshev fit details.

$He^+ + H^+ \rightarrow (He^{2+} + H^+ + e)$ or $(He^{2+} + H)$

Beam – Maxwellian



Ionization and He^{2+} Production Cross Sections for

$He^+ + He^+ \rightarrow (He^+ + He^{2+} + e)$ (σ_i)
$He^+ + He^+ \rightarrow (He + He^{2+})$ or $(He^{+} + He^{2+} + e)$ ($\sigma_{He^{2+}}$)

Ionization		He^{2+} Production	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
1.0E+04	6.35E-19	5.9E+03	3.46E-18
1.5E+04	1.72E-18	6.0E+03	3.67E-18
2.0E+04	3.12E-18	7.0E+03	4.94E-18
2.5E+04	4.75E-18	8.0E+03	6.38E-18
3.0E+04	6.72E-18	9.0E+03	7.84E-18
3.5E+04	8.79E-18	1.0E+04	9.34E-18
4.0E+04	1.08E-17	1.5E+04	1.55E-17
4.5E+04	1.32E-17	2.0E+04	1.98E-17
5.0E+04	1.56E-17	2.5E+04	2.25E-17
5.5E+04	1.81E-17	3.0E+04	2.50E-17
5.8E+04	1.99E-17	3.5E+04	2.69E-17
		4.0E+04	2.82E-17
		4.5E+04	2.94E-17
		5.0E+04	3.00E-17
		5.5E+04	3.02E-17
		6.0E+04	3.03E-17
		6.7E+04	2.98E-17

References: 69, 277, 457

Accuracy: σ_i - Factor of 2
$\sigma_{He^{2+}}$ - 50%

Notes: (1) The production of He^{2+} arises from two processes: (a) Charge exchange ($He^+ + He^+ \rightarrow He + He^{2+}$); (b) Ionization ($He^+ + He^+ \rightarrow He^+ + He^{2+} + e$). (2) At energies less than 2.0 - 3.0x10^5 eV/amu the charge exchange reaction dominates. (3) The incident He^+ energy in eV/amu is plotted vs. the stationary He^+ target.

Chebyshev Fitting Parameters for Cross Sections

Ionization E_{min} = 1.0E+04 eV/amu, E_{max} = 5.8E+04 eV/amu
He^{2+} Prod. E_{min} = 5.9E+03 eV/amu, E_{max} = 6.7E+04 eV/amu

	A0	A1	A2	A3	A4	A5	A6
Ionization	-80.1208	1.69568	-.114597	.0198330	-.00494221	.00648764	.00123081
He^{2+} Prod.	-77.5761	1.04465	-.328288	.0385208	-.00883058	-.0130928	.00201232

The fit represents the Ionization cross section with an rms deviation of 0.5%. The maximum deviation is 1.0% at 4.0E+04 eV/amu.

The fit represents the He^{2+} Prod. cross section with an rms deviation of 0.5%. The maximum deviation is 1.2% at 5.9E+03 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + He^+ \rightarrow (He^+ + He^{2+} + e)$ or $(He + He^{2+})$ $(\sigma_{He^{2+}})$
$He^+ + He^+ \rightarrow He^+ + He^{2+} + e$ (σ_i)



Single Ionization Rate Coefficients for

$He^+ + He^+ \rightarrow He^+ + He^{2+} + e$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^+ Temp. (eV)	He^+ Energy (eV/amu) 10000.	20000.	30000.	40000.	50000.	52000.	55000.
1.0E+00	4.50E-11*	5.85E-10	1.62E-09	2.96E-09	4.85E-09	5.18E-09	5.62E-09
2.0E+00	4.53E-11*	6.12E-10	1.62E-09	3.00E-09	4.85E-09	5.25E-09	5.89E-09
4.0E+00	4.59E-11*	6.13E-10	1.62E-09	3.00E-09	4.85E-09	5.25E-09	5.90E-09
7.0E+00	4.65E-11*	6.13E-10	1.62E-09	3.00E-09	4.85E-09	5.25E-09	5.90E-09
1.0E+01	4.69E-11*	6.13E-10	1.62E-09	3.00E-09	4.85E-09	5.25E-09	5.90E-09
2.0E+01	4.82E-11*	6.13E-10	1.62E-09	3.01E-09	4.85E-09	5.26E-09	5.91E-09
4.0E+01	5.00E-11*	6.14E-10	1.62E-09	3.01E-09	4.85E-09	5.26E-09	5.90E-09
7.0E+01	5.22E-11*	6.15E-10	1.62E-09	3.01E-09	4.85E-09	5.26E-09	5.81E-09
1.0E+02	5.39E-11*	6.16E-10	1.62E-09	3.02E-09	4.86E-09	5.27E-09	5.67E-09
2.0E+02	5.89E-11*	6.21E-10	1.63E-09	3.03E-09	4.87E-09	5.24E-09	5.20E-09*
4.0E+02	6.68E-11*	6.31E-10	1.64E-09	3.06E-09	4.84E-09	5.05E-09	4.60E-09*
7.0E+02	7.70E-11*	6.46E-10	1.67E-09	3.09E-09	4.68E-09	4.70E-09*	4.10E-09*
1.0E+03	8.64E-11*	6.62E-10	1.69E-09	3.12E-09	4.49E-09*	4.40E-09*	3.79E-09*
2.0E+03	1.15E-10*	7.16E-10	1.76E-09	3.18E-09	3.93E-09*	3.75E-09*	3.24E-09*
4.0E+03	1.71E-10*	8.25E-10	1.91E-09	3.11E-09	3.28E-09*	3.10E-09*	2.74E-09#
7.0E+03	2.60E-10*	9.89E-10	2.04E-09	2.86E-09*	2.75E-09*	2.62E-09#	2.36E-09#
1.0E+04	3.57E-10	1.14E-09	2.07E-09*	2.62E-09*	2.44E-09#	2.33E-09#	2.13E-09#
2.0E+04	6.89E-10	1.42E-09*	1.95E-09*	2.09E-09*	1.88E-09#	1.81E-09#	1.69E-09#

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

He^+ Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-46.1185	1.19034	.579487	.183843	.0279475	-.00786851	-.00843904
20000.	-42.0417	.348208	.209457	.102371	.0130005	-.00449377	-.0168872
30000.	-40.3424	.114393	.0591680	.00327281	-.0216154	-.0232829	-.0142410
40000.	-39.3369	-.0868409	-.0958221	-.0743825	-.0480228	-.0161228	-1.557744E-05
50000.	-38.7358	-.388572	-.250357	-.102423	-.00799043	.0201351	.00955123
52000.	-38.6794	-.462898	-.271231	-.0815950	.00845828	.0221265	-7.933020E-04
55000.	-38.6693	-.588564	-.267081	-.0118366	.0203911	-.00168618	-.0171873

See appendix for Chebyshev fit details.

$He^+ + He^+ \rightarrow He^+ + He^{2+} + e$

Beam – Maxwellian



He^{2+} Production Rate Coefficients for

$He^{+} + He^{+} \rightarrow (He^{+} + He^{2+} + e)$ or $(He + He^{2+})$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^{+} Temp. (eV)	He^{+} Energy (eV/amu) 10000.	20000.	30000.	40000.	50000.	60000.	65000.
1.0E+00	1.30E-09	3.89E-09	5.71E-09	7.84E-09	9.32E-09	1.03E-08	1.06E-08
2.0E+00	1.30E-09	3.89E-09	5.99E-09	7.83E-09	9.32E-09	1.03E-08	1.06E-08
4.0E+00	1.30E-09	3.89E-09	6.01E-09	7.83E-09	9.32E-09	1.03E-08	1.06E-08
7.0E+00	1.29E-09	3.88E-09	6.01E-09	7.83E-09	9.31E-09	1.03E-08	1.06E-08
1.0E+01	1.29E-09	3.88E-09	6.01E-09	7.84E-09	9.31E-09	1.03E-08	1.06E-08
2.0E+01	1.29E-09	3.88E-09	6.01E-09	7.84E-09	9.31E-09	1.03E-08	1.05E-08
4.0E+01	1.29E-09	3.88E-09	6.01E-09	7.84E-09	9.31E-09	1.03E-08	1.02E-08
7.0E+01	1.30E-09	3.88E-09	6.01E-09	7.84E-09	9.31E-09	1.03E-08	9.57E-09*
1.0E+02	1.30E-09	3.88E-09	6.01E-09	7.84E-09	9.31E-09	1.03E-08	9.11E-09*
2.0E+02	1.31E-09	3.88E-09	6.02E-09	7.85E-09	9.30E-09	1.03E-08	8.19E-09*
4.0E+02	1.33E-09	3.89E-09	6.03E-09	7.85E-09	9.30E-09	9.99E-09	7.34E-09*
7.0E+02	1.36E-09	3.90E-09	6.04E-09	7.86E-09	9.29E-09	9.46E-09	6.76E-09*
1.0E+03	1.39E-09	3.92E-09	6.06E-09	7.87E-09	9.28E-09	9.00E-09*	6.44E-09*
2.0E+03	1.49E-09	3.98E-09	6.10E-09	7.89E-09	9.11E-09	7.96E-09*	5.87E-09*
4.0E+03	1.70E-09	4.11E-09	6.20E-09	7.87E-09	8.50E-09	6.90E-09*	5.36E-09*
7.0E+03	1.99E-09	4.31E-09	6.30E-09	7.63E-09	7.66E-09*	6.10E-09*	4.95E-09#
1.0E+04	2.26E-09	4.50E-09	6.32E-09	7.29E-09*	7.02E-09*	5.60E-09*	4.67E-09#
2.0E+04	3.05E-09	4.89E-09	6.01E-09*	6.26E-09*	5.71E-09*	4.67E-09#	4.08E-09#

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

He^{+} Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-40.5559	.324800	.215507	.101128	.0305437	.00225930	-.00348368
20000.	-38.6393	.0830639	.0600817	.0314309	.0105748	.00101472	-.00206469
30000.	-37.8497	.0259326	-.00436636	.00382678	-.0161047	-.00327000	-.00929120
40000.	-37.3876	-.0574256	-.0510653	-.0409152	-.0269282	-.0144190	-.00522979
50000.	-37.1678	-.166608	-.125810	-.0745545	-.0317364	-.00593417	.00489997
60000.	-37.1929	-.348477	-.198373	-.0604564	.00910326	.0165438	4.816223E-04
65000.	-37.4160	-.490762	-.135825	.0331097	.00708503	-.0235912	-.00325326

See appendix for Chebyshev fit details

$He^+ + He^+ \rightarrow (He^+ + He^{2+} + e)$ or $(He + He^{2+})$



Cross Sections for Producting Slow Ions and Electrons for

$He + H_2 \rightarrow \Sigma$ slow ions (σ^+)
$He + H_2 \rightarrow \Sigma$ slow electrons (σ^-)

(σ^+) Energy (eV/amu)	Cross Section (cm^2)	(σ^-) Energy (eV/amu)	Cross Section (cm^2)
7.6E+02	3.28E-17	1.2E+00	2.30E-20
8.0E+02	3.45E-17	1.3E+00	2.00E-19
1.0E+03	4.21E-17	1.4E+00	5.36E-19
2.0E+03	6.90E-17	1.5E+00	4.86E-19
4.0E+03	1.02E-16	1.7E+00	4.08E-19
6.0E+03	1.26E-16	2.0E+00	8.65E-19
8.0E+03	1.46E-16	2.5E+00	2.36E-18
1.0E+04	1.63E-16	3.0E+00	3.97E-18
2.0E+04	2.17E-16	4.0E+00	6.94E-18
4.0E+04	2.54E-16	5.0E+00	1.02E-17
6.0E+04	2.50E-16	6.0E+00	1.35E-17
8.0E+04	2.35E-16	7.0E+00	1.75E-17
1.0E+05	2.18E-16	8.0E+00	2.07E-17
2.0E+05	1.43E-16	9.0E+00	2.31E-17
2.5E+05	1.17E-16	1.0E+01	2.47E-17
		4.0E+01	5.52E-17
		7.0E+01	7.17E-17
		1.0E+02	8.42E-17
		4.0E+02	1.46E-16
		7.0E+02	1.77E-16
		1.0E+03	2.00E-16
		4.0E+03	3.06E-16
		7.0E+03	3.53E-16
		1.0E+04	3.77E-16
		4.0E+04	4.46E-16
		7.0E+04	4.15E-16
		1.0E+05	3.66E-16
		2.5E+05	2.06E-16

References: σ^+ - 143, 459, 463
σ^- - 143, 460

Accuracy: σ^+ - 20%
σ^- - 1.0 eV/amu to 12 eV/amu - 20%
12 eV/amu to 3.7×10^4 eV/amu - Unknown
3.7×10^4 eV/amu to 2.5×10^4 eV/amu - 20%

Notes: (1) Note the small resonance at 1.4 eV/amu results from metastable He. (2) The cross section σ^+ is defined as the sum of cross sections for all processes producing slow singly-charged ions plus twice the cross section for producing doubly-charged slow ions. Similarly the σ^- cross section is defined as the cross section for ionization plus the cross section for electron loss or stripping. In those reactions where electron capture is important, the σ^+ cross section is the sum of the ionization cross section and the electron capture cross section.

Chebyshev Fitting Parameters for Cross Sections

σ^+ E_{min} = 7.6E+02 eV/amu, E_{max} = 2.5E+05 eV/amu
σ^- E_{min} = 1.2E+00 eV/amu, E_{max} = 2.5E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
σ^+	-73.4532	.742441	-.556022	-.126285	-.0371599	.0218408	.000323040
σ^-	-75.1363	3.03724	-1.67952	.577117	-.601974	.234746	-.168549

The fit represents the σ^+ cross section with an rms deviation of 0.3%. The maximum deviation is 0.4% at 8.0E+04 eV/amu.

The fit represents the σ^- cross section with an rms deviation of 52.2%. The maximum deviation is 286.3% at 1.2E+00 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + He^+ \rightarrow \Sigma$ Slow Ions (σ^+)
$He^+ + He^+ \rightarrow \Sigma$ Slow Electrons (σ^-)



Ionization and H^+ Production Cross Sections for

$$He + H_2 \rightarrow He + H_2^+ + e$$
$$He + H_2 \rightarrow \Sigma H^+$$

Ionization (H_2^+ Production)		H^+ (Production)	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
7.4E+02	1.64E-17	7.4E+02	2.17E-17
8.0E+02	1.75E-17	8.0E+02	2.19E-17
9.0E+02	1.93E-17	9.0E+02	2.23E-17
1.0E+03	2.11E-17	1.0E+03	2.28E-17
1.5E+03	2.97E-17	1.5E+03	2.52E-17
2.0E+03	3.81E-17	2.0E+03	2.78E-17
3.0E+03	5.24E-17	3.0E+03	3.32E-17
4.0E+03	6.55E-17	4.0E+03	3.69E-17
5.0E+03	7.72E-17	4.3E+03	3.74E-17
6.0E+03	8.71E-17	5.0E+03	3.67E-17
7.0E+03	9.55E-17	6.0E+03	3.55E-17
8.0E+03	1.03E-16	7.0E+03	3.52E-17
9.0E+03	1.10E-16	8.0E+03	3.55E-17
1.0E+04	1.15E-16	9.0E+03	3.68E-17
1.2E+04	1.24E-16	1.0E+04	3.87E-17
		1.1E+04	4.20E-17

References: 459, 461, 462, 463, 464

Accuracy: Unknown

Notes: (1) The data of Refs. 461, 462, 463, and 464 are approximately a factor of 2-3 less than that of Ref. 459 for the production of H^+ ions. The data presented follows closely the results of Ref. 459.

Chebyshev Fitting Parameters for Cross Sections

H_2^+ Prod. E_{min} = 7.4E+02 eV/amu, E_{max} = 1.2E+04 eV/amu
H^+ Prod. E_{min} = 7.4E+02 eV/amu, E_{max} = 1.1E+04 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H_2^+ Prod.	-75.1222	1.03406	-.0748886	-.0222039	-.00237843	-.000779509	.000954515
H^+ Prod.	-76.0878	.319638	-.0448970	-.0183961	.0505953	.0349869	-.000131466

The fit represents the H_2^+ cross section with an rms deviation of 0.2%. The maximum deviation is 0.4% at 1.5E+03 eV/amu.

The fit represents the H^+ cross section with an rms deviation of 1.0%. The maximum deviation is 1.8% at 6.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$$He + H_2 \rightarrow He + H_2^+ + e \ (\sigma_{H_2^+})$$
$$He + H_2 \rightarrow \Sigma H^+ \ (\sigma_{H^+})$$



Electron and Slow Positive Ion Production Cross Sections for

He + He -> Σ slow electrons (σ^-)
He + He -> Σ slow ions (σ^+)

σ^+		σ^- (Lower Energy)		σ^- (Higher Energy)	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (ev/amu)	σ (cm^2)
5.0E+03	4.45E-17	2.7E+01	2.56E-20	3.7E+04	2.47E-16
6.0E+03	5.23E-17	3.0E+01	1.34E-19	4.0E+04	2.50E-16
7.0E+03	5.88E-17	3.5E+01	3.31E-19	5.0E+04	2.50E-16
8.0E+03	6.49E-17	4.0E+01	4.86E-19	6.0E+04	2.47E-16
9.0E+03	7.07E-17	5.0E+01	7.66E-19	7.0E+04	2.43E-16
1.0E+04	7.55E-17	6.0E+01	1.06E-18	8.0E+04	2.38E-16
1.5E+04	9.74E-17	7.0E+01	1.33E-18	9.0E+04	2.33E-16
2.0E+04	1.10E-16	8.0E+01	1.61E-18	1.0E+05	2.25E-16
3.0E+04	1.23E-16	9.0E+01	1.88E-18	1.5E+05	1.91E-16
4.0E+04	1.28E-16	1.0E+02	2.16E-18	2.0E+05	1.67E-16
5.0E+04	1.29E-16	1.5E+02	3.52E-18	2.5E+05	1.43E-16
6.0E+04	1.28E-16	2.0E+02	4.77E-18		
7.0E+04	1.26E-16	2.5E+02	5.89E-18		
8.0E+04	1.22E-16	3.0E+02	6.96E-18		
9.0E+04	1.19E-16				
1.0E+05	1.16E-16				
1.5E+05	9.91E-17				
2.0E+05	8.49E-17				
2.5E+05	7.44E-17				

References: 143, 403, 462, 463, 472, 473, 474, 475

Accuracy: Unknown

Notes: (1) The processes to form slow positive ions and electrons are: (a) He + He -> He^+ + He + e, electron loss; (b) He + He -> He^+ + He^+ + 2e (electron loss with target ionization); (c) He + He -> He + He^+ + e, single ionization; (d) He + He -> He + He^{2+} + 2e, double ionization; (e) He + He -> He^- + He^+, electron capture. (2) Cross sections for slow positive ion formation (σ^+) are the sum of reactions σ(b) + σ(c) + σ(d) + σ(e). (3) For electrons, (σ^-) is the sum of σ(a) + 2σ(b) + σ(c) + 2σ(d). For higher energies σ(e) is very small and thus σ^+ is approximately equal to the sum of the single and double ionization cross sections. (4) No attempt has been made to interpolate between the high-energy σ^- and the low-energy data (E < 300 eV/amu).

Chebyshev Fitting Parameters for Cross Sections

σ^+ E_{min} = 5.0E+03 eV/amu, E_{max} = 2.5E+05 eV/amu
σ^- (lower E) E_{min} = 2.7E+01 eV/amu, E_{max} = 3.0E+02 eV/amu
σ^- (Higher E) E_{min} = 3.7E+04 eV/amu, E_{max} = 2.5E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
σ^+	-74.0092	.259096	-.395285	-.00216683	.00569771	-.00263905	.00150963
σ^- (lower E)	-82.5433	2.19805	-.641453	.385156	-.256300	.170524	-.106115
σ^- (Higher E)	-72.2261	-.269539	-.0908126	-.000390444	-.000723589	-.00319421	-.00598024

The fit represents the σ^+ cross section with an rms deviation of 0.4%.
The maximum deviation is 0.9% at 1.0E+04 eV/amu.
The fit represents the σ^- (lower E) cross section with an rms deviation of 6.3%.
The maximum deviation is 10.0% at 4.0E+01 eV/amu.
The fit represents the σ^- (Higher E) cross section with an rms deviation of 0.1%.
The maximum deviation is 0.2% at 8.0E+04 eV/amu.

See appendix for Chebyshev fit details.

He + He –> ΣSlow Electrons (σ^{**})
He + He –> ΣSlow Ions (σ^{+})



Single and Double Ionization Cross Sections for

$$He + He \rightarrow He + He^{+} + e$$
$$He + He \rightarrow He + He^{2+} + 2e$$

Single Ionization		Double Ionization	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
1.2E+04	8.26E-17	1.3E+04	9.93E-19
1.5E+04	1.03E-16	1.5E+04	1.40E-18
2.0E+04	1.32E-16	2.0E+04	2.14E-18
2.5E+04	1.58E-16	2.5E+04	2.69E-18
3.0E+04	1.80E-16	3.0E+04	3.15E-18
3.5E+04	1.99E-16	3.5E+04	3.59E-18
4.0E+04	2.13E-16	4.0E+04	4.03E-18
4.5E+04	2.25E-16	4.5E+04	4.44E-18

References: 462, 463

Accuracy: Unknown

Notes: None

Chebyshev Fitting Parameters for Cross Sections

Single Ion. E_{min} = 1.2E+04 eV/amu, E_{max} = 4.5E+04 eV/amu
Double Ion. E_{min} = 1.3E+04 eV/amu, E_{max} = 4.5E+04 eV/amu

	A0	A1	A2	A3	A4	A5	A6
Single Ion.	-72.9678	.503766	-.0453017	-.00416573	-.00286603	.00135469	.000526378
Double Ion.	-81.1695	.703077	-.106105	.0433845	-.0115100	.00236396	-.00233450

The fit represents the Single Ion. cross section with an rms deviation of 0.1%. The maximum deviation is 0.2% at 3.0E+04 eV/amu.

The fit represents the Double Ion. cross section with an rms deviation of 0.1%. The maximum deviation is 0.1% at 3.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$$He + He \rightarrow He + He^{+} + e$$
$$He + He \rightarrow He + He^{2+} + 2e$$



Ionization Cross Sections for

$He^+ + H \rightarrow He^+ + H^+ + e$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
6.3E+03	1.10E+08	1.28E-17
7.0E+03	1.16E+08	1.47E-17
8.0E+03	1.24E+08	1.75E-17
9.0E+03	1.32E+08	2.07E-17
1.0E+04	1.39E+08	2.45E-17
1.5E+04	1.70E+08	4.80E-17
2.0E+04	1.96E+08	8.07E-17
3.0E+04	2.41E+08	1.46E-16
4.0E+04	2.78E+08	1.98E-16
5.0E+04	3.11E+08	2.16E-16
6.0E+04	3.40E+08	2.18E-16
7.0E+04	3.68E+08	2.10E-16
8.0E+04	3.93E+08	2.00E-16
9.0E+04	4.17E+08	1.91E-16
1.0E+05	4.39E+08	1.78E-16
1.5E+05	5.38E+08	1.36E-16
2.0E+05	6.21E+08	1.10E-16
3.0E+05	7.61E+08	8.04E-17
4.0E+05	8.78E+08	6.41E-17
5.0E+05	9.82E+08	5.36E-17
6.0E+05	1.08E+09	4.56E-17
7.0E+05	1.16E+09	4.04E-17
8.0E+05	1.24E+09	3.61E-17
9.0E+05	1.32E+09	3.26E-17
1.0E+06	1.39E+09	2.94E-17
2.0E+06	1.96E+09	1.59E-17
3.0E+06	2.40E+09	1.09E-17
4.0E+06	2.77E+09	8.33E-18
5.0E+06	3.09E+09	6.74E-18
6.0E+06	3.39E+09	5.76E-18
7.0E+06	3.65E+09	4.97E-18
8.0E+06	3.90E+09	4.39E-18

References: 564, 565, 566

Accuracy: Unknown

Notes: (1) As far as we know no published experimental data exist. The data plotted are theoretical results. (2) For energies less than 10^5 eV/amu the data of Ref. 566 has been plotted. The classical Monte Carlo calculations have been quite accurate in predicting similar collisional processes where experimental data is available for comparison. (3) For energies greater than 10^5 eV/amu data of Ref. 565 is plotted. These calculations are Born approximations which should be good at these high energies. At lower energies the cross sections are much higher than those given in Ref. 566.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 6.3E+03 eV/amu, E_{max} = 8.0E+06 eV/amu

A0	A1	A2	A3	A4	A5	A6
-76.1808	-.896422	-1.40016	.494799	-.0137803	-.135113	.110603

The fit represents the above cross section with an rms deviation of 3.2%. The maximum deviation is 6.7% at 1.5E+04 eV/amu.

See appendix for Chebyshev fit details.

$$He^+ + H \rightarrow He^+ + H^+ + e$$



Ionization Rate Coefficients for

$H + He^+ \rightarrow H^+ + He^+ + e$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^+ Temp. (eV)	H Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	3.41E-09	1.58E-08	5.49E-08	7.72E-08	7.82E-08	6.83E-08	5.26E-08
2.0E+00	3.41E-09	1.58E-08	5.49E-08	7.71E-08	7.82E-08	6.83E-08	5.26E-08
4.0E+00	3.41E-09	1.58E-08	5.48E-08	7.71E-08	7.82E-08	6.83E-08	5.26E-08
7.0E+00	3.42E-09	1.58E-08	5.48E-08	7.71E-08	7.82E-08	6.83E-08	5.26E-08
1.0E+01	3.43E-09	1.58E-08	5.47E-08	7.71E-08	7.82E-08	6.83E-08	5.26E-08
2.0E+01	3.45E-09	1.58E-08	5.46E-08	7.71E-08	7.82E-08	6.83E-08	5.26E-08
4.0E+01	3.49E-09	1.59E-08	5.45E-08	7.71E-08	7.82E-08	6.83E-08	5.26E-08
7.0E+01	3.55E-09	1.60E-08	5.43E-08	7.70E-08	7.82E-08	6.83E-08	5.26E-08
1.0E+02	3.61E-09	1.61E-08	5.43E-08	7.70E-08	7.82E-08	6.83E-08	5.26E-08
2.0E+02	3.81E-09	1.64E-08	5.41E-08	7.69E-08	7.81E-08	6.83E-08	5.26E-08
4.0E+02	4.20E-09	1.72E-08	5.40E-08	7.67E-08	7.81E-08	6.82E-08	5.26E-08
7.0E+02	4.84E-09	1.83E-08	5.39E-08	7.65E-08	7.79E-08	6.82E-08	5.25E-08
1.0E+03	5.52E-09	1.93E-08	5.39E-08	7.62E-08	7.77E-08	6.82E-08	5.25E-08
2.0E+03	7.93E-09	2.25E-08	5.41E-08	7.52E-08	7.72E-08	6.81E-08	5.25E-08
4.0E+03	1.29E-08	2.76E-08	5.46E-08	7.34E-08	7.62E-08	6.80E-08	5.24E-08
7.0E+03	1.97E-08	3.34E-08	5.55E-08	7.14E-08	7.48E-08	6.78E-08	5.23E-08
1.0E+04	2.56E-08	3.78E-08	5.63E-08	6.99E-08	7.35E-08	6.75E-08	5.23E-08
2.0E+04	3.99E-08	4.76E-08	5.86E-08	6.73E-08	7.03E-08	6.66E-08	5.21E-08

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

H Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-37.6259	1.12300	.601815	.164371	-.0334027	-.0580916	-.0243672
20000.	-35.3341	.489409	.271922	.0807229	-.00815166	-.0204835	-.00815012
40000.	-33.4270	.0164455	.02281157	.0149114	.00585597	.00147262	-3.367767E-06
70000.	-32.8194	-.0548059	-.0346997	-.0154134	-.00335201	.00137249	.00185552
100000.	-32.7709	-.0377570	-.0264510	-.0142784	-.00575226	-.00155112	-3.572900E-05
200000.	-33.0075	-.00828099	-.00567988	-.00349931	-.00201522	-.00104778	-4.638534E-04
500000.	-33.5256	-.00412902	-.00205167	-6.187466E-04	-2.150041E-04	-7.937795E-05	-1.271717E-04

See appendix for Chebyshev fit details.

$$H + He^+ \rightarrow He^+ + H^+ + e$$



Cross Sections for the Production of Slow Positive Ions and Electrons for He^+ Passing Through H_2

$$He^+ + H_2 \rightarrow \Sigma \text{ Slow Ions } \sigma^+$$
$$\rightarrow \Sigma \text{ Electrons } \sigma^-$$

Slow Ions σ^+		Electrons σ^-	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
7.5E+02	1.28E-16	2.5E+03	4.19E-17
8.0E+02	1.24E-16	3.0E+03	4.59E-17
1.0E+03	1.15E-16	4.0E+03	5.36E-17
1.3E+03	1.11E-16	7.0E+03	7.38E-17
1.5E+03	1.14E-16	1.0E+04	9.89E-17
2.0E+03	1.34E-16	1.5E+04	1.52E-16
4.0E+03	2.66E-16	2.0E+04	2.01E-16
7.0E+03	3.72E-16	4.0E+04	3.30E-16
1.0E+04	4.20E-16	6.0E+04	3.61E-16
1.5E+04	4.65E-16	7.0E+04	3.63E-16
2.0E+04	4.92E-16	1.0E+05	3.36E-16
3.0E+04	5.08E-16	1.5E+05	2.84E-16
4.0E+04	5.03E-16	2.0E+05	2.46E-16
7.0E+04	4.11E-16	4.0E+05	1.67E-16
1.0E+05	3.41E-16	5.0E+05	1.46E-16
1.5E+05	2.70E-16		
2.0E+05	2.29E-16		
4.0E+05	1.46E-16		
5.0E+05	1.24E-16		

<u>References:</u> 96, 104, 105, 353, 355, 364, 462, 463, 478, 479, 480, 481

<u>Accuracy:</u> 50%

<u>Notes:</u> (1) The reactions that produce slow ions and electrons as reaction products are: (a) $He^+ + H_2 \rightarrow He^+ + H_2^+ + e$, single ionization; (b) $He^+ + H_2 \rightarrow He + H_2^+$, single capture; (c) $He^+ + H_2 \rightarrow He^+ + H^+ + H + e$, dissociative ionization; (d) $He^+ + H_2 \rightarrow He^+ + H^+ + H^+ + 2e$, double ionization; (e) $He^+ + H_2 \rightarrow He + H^+ + H$, dissociative electron capture; (f) $He^+ + H_2 \rightarrow He + H^+ + H^+ + e$, electron capture with ionization. σ^+ is equal to the sum $\sigma(a) + \sigma(b) + \sigma(c) + 2\sigma(d) + \sigma(e) + 2\sigma(f)$. σ^- is equal to $\sigma(a) + \sigma(c) + 2\sigma(d) + \sigma(f)$. (2) For energies less than 10^5 eV/amu the electron capture cross sections to form H_2^+ dominate the σ^+ cross sections. (3) the σ^- cross sections of Ref. 355 are not plotted below 2.5×10^3 eV/amu due to the small energy dependence of the cross sections.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

Slow Ions σ^+ E_{min} = 7.5E+02 eV/amu, E_{max} = 5.0E+05 eV/amu
Electrons σ^- E_{min} = 2.5E+03 eV/amu, E_{max} = 5.0E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
Slow Ions σ^+	-72.1635	.175055	-.713665	-.133412	.172753	-.0619895	.0345550
Electrons σ^-	-73.0711	.770468	-.649678	-.176586	.134550	.0300410	-.0402778

The fit represents the σ^+ cross section with an rms deviation of 3.2%. The maximum deviation is 5.5% at 2.0E+03 eV/amu.

The fit represents the σ^- cross section with an rms deviation of 0.7%. The maximum deviation is 1.6% at 1.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + H_2 \rightarrow \Sigma$Slow Ions ($\sigma^+$)
$He^+ + H_2 \rightarrow \Sigma$Electrons ($\sigma^-$)



Cross Sections for the Formation of H_2^+, H^+, and for Double Ionization for He^+ in H_2

$$He^+ + H_2 \rightarrow He + H_2^+$$
$$\rightarrow He^+ + H_2^+ + e \quad [\sigma(H_2^+)]$$

$$He^+ + H_2 \rightarrow \Sigma H^+ \quad [\sigma(H^+)]$$

$$He^+ + H_2 \rightarrow He^+ + H^+ + H^+ + 2e \quad \text{(Double Ionization)}$$

H_2^+ Production		H^+ Production		Double Ionization	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
7.3E+02	3.08E-17	7.4E+02	9.39E-17	2.5E+04	4.77E-18
8.0E+02	3.23E-17	8.0E+02	8.99E-17	3.0E+04	4.53E-18
1.0E+03	3.55E-17	9.0E+02	8.45E-17	4.0E+04	3.68E-18
1.5E+03	4.45E-17	1.0E+03	8.00E-17	5.0E+04	2.86E-18
2.0E+03	5.45E-17	1.5E+03	6.92E-17	6.0E+04	2.31E-18
4.0E+03	1.04E-16	2.0E+03	6.71E-17	7.0E+04	1.92E-18
7.0E+03	1.84E-16	2.5E+03	6.65E-17	8.0E+04	1.62E-18
1.0E+04	2.83E-16	3.0E+03	6.70E-17	9.0E+04	1.40E-18
1.5E+04	3.96E-16	4.0E+03	6.93E-17	1.0E+05	1.22E-18
2.0E+04	4.21E-16	5.0E+03	7.50E-17	1.5E+05	6.79E-19
4.0E+04	3.93E-16	6.0E+03	8.29E-17	2.0E+05	4.23E-19
7.0E+04	3.37E-16	7.0E+03	9.05E-17	3.0E+05	2.26E-19
1.0E+05	2.83E-16	8.0E+03	1.02E-16	4.0E+05	1.42E-19
1.5E+05	2.22E-16	8.6E+03	1.05E-16	5.0E+05	1.00E-19
2.0E+05	1.86E-16	9.0E+03	1.04E-16	6.0E+05	7.31E-20
4.0E+05	1.25E-16	1.0E+04	9.64E-17	7.0E+05	5.83E-20
7.0E+05	9.17E-17	1.2E+04	8.49E-17	8.0E+05	4.73E-20
8.0E+05	8.51E-17			9.0E+05	4.09E-20

References: 360, 365, 462, 463, 481, 482

Accuracy: $\sigma(H_2^+)$ - 20%
$\sigma(H^+)$ - Unknown
σ(Double Ion.) - 20%

Notes: (1) H_2^+ is formed from ionization of H_2 by He^+ and from electron capture collisions. (2) H^+ is formed from dissociative ionization, $He^+ + H_2 \rightarrow He^+ + H^+ + H + e$; from double ionization, $He^+ + H_2 \rightarrow He^+ + H^+ + H^+ + 2e$; and from transfer ionization, $He^+ + H_2 \rightarrow He + H^+ + H^+ + e$.

Chebyshev Fitting Parameters for Cross Sections

H_2^+ Prod. E_{min} = 7.3E+02 eV/amu, E_{max} = 8.0E+05 eV/amu
H^+ Prod. E_{min} = 7.4E+02 eV/amu, E_{max} = 1.2E+04 eV/amu
Double Ion. E_{min} = 2.5E+04 eV/amu, E_{max} = 9.0E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H_2^+ Prod.	-73.3806	.601066	-1.00967	-.0679372	.247157	-.0416476	-.0640838
H^+ Prod.	-74.0876	.0639707	.162049	-.0778454	-.0615354	-.0437939	-.0172413
Double Ion.	-84.0918	-2.48636	-.206129	.0854195	-.00909082	.0148658	-.00608471

The fit represents the H_2^+ Prod. cross section with an rms deviation of 3.7%. The maximum deviation is 9.0% at 4.0E+04 eV/amu.

The fit represents the H^+ Prod. cross section with an rms deviation of 1.8%. The maximum deviation is 5.7% at 1.0E+04 eV/amu.

The fit represents the Double Ion. cross section with an rms deviation of 1.1%. The maximum deviation is 2.5% at 2.0E+05 eV/amu.

See appendix for Chebyshev fit details.


He+ + H2 -> σ(H2+)
He+ + H2 -> σ(H+)
He+ + H2 -> He+ + 2H+ + 2e
Cross Section vs. Energy
Cross Section (cm2)
Energy (eV/amu)
Δ = σ(H2+)
× = σ(H+)
∇ = He+ + 2H+ + 2e
Recommended Data
Chebyshev Fit

Cross Sections for Production of Slow Positive Charges (σ^+) and Slow Electrons (σ^-) for He^+ Passing Through He.

$He^+ + He \rightarrow \sigma^+, \sigma^-$

$He^+ + He$ (σ^+) Energy (eV/amu)	Cross Section (cm^2)	$He^+ + He$ (σ^-) Energy (eV/amu)	Cross Section (cm^2)
2.5E+03	6.13E-16	2.5E+03	1.40E-17
4.0E+03	5.83E-16	4.0E+03	2.15E-17
7.0E+03	5.05E-16	7.0E+03	3.37E-17
1.0E+04	4.49E-16	1.0E+04	4.46E-17
2.0E+04	3.49E-16	2.0E+04	7.56E-17
4.0E+04	2.68E-16	4.0E+04	1.19E-16
7.0E+04	2.12E-16	7.0E+04	1.59E-16
1.0E+05	1.84E-16	1.0E+05	1.81E-16
2.0E+05	1.30E-16	1.1E+05	1.83E-16
4.0E+05	7.58E-17	2.0E+05	1.51E-16
7.0E+05	4.72E-17	4.0E+05	9.77E-17
1.0E+06	3.53E-17	5.0E+05	8.39E-17
2.0E+06	1.94E-17		
2.8E+06	1.45E-17		

<u>References:</u> 96, 104, 105, 355, 364, 388, 463, 472, 478, 479, 480, 487, 493, 497, 498, 499, 500, 579

<u>Accuracy:</u> $E > 3\text{-}4\times10^4$ eV/amu - 20%
$E < 3\text{-}4\times10^4$ eV/amu Unknown

Notes: (1) σ^+ and σ^- are the sum of the cross sections for processes producing slow ions and electrons, respectively. (2) The reaction channels contributing to these processes are (a) single electron capture; (b) single and double ionization of the target atom; (c) single capture plus ionization of the target; (d) capture into the continuum; (e) electron loss; (f) electron loss plus ionization of the target. For He^+ + He collisions, proceses (c, d and f) are believed to contribute a factor of ten less than the other processes especially at low energies. (3) For He^+ energies less than 10^5 eV/amu, the low-energy σ^- data of Refs. 104, 105, 355, 364, 463, 480, 487, 498, and 500 are not included in the present compilation. These data are questionable since the cross sections are only slightly dependent on energy. (4) For low energies the σ^+ cross sections are dominated by single electron capture collisions.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

$He^+ + He$ (σ^+) E_{min} = 2.5E+03 eV/amu, E_{max} = 2.8E+06 eV/amu
$He^+ + He$ (σ^-) E_{min} = 2.5E+03 eV/amu, E_{max} = 5.0E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
$He^+ + He$ (σ^+)	-73.0794	-1.87553	-.368452	-.0292343	.0164797	.0394079	-.00613696
$He^+ + He$ (σ^-)	-74.6255	1.02842	-.609151	-.175413	-.00862801	.0447902	.0224536

The fit represents the $He^+ + He$ (σ^+) cross section with an rms deviation of 1.5%. The maximum deviation is 2.1% at 7.0E+05 eV/amu.

The fit represents the $He^+ + He$ (σ^-) cross section with an rms deviation of 1.5%. The maximum deviation is 2.5% at 2.0E+05 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + He \rightarrow \sigma^+$
$He^+ + He \rightarrow \sigma^-$



Ionization Rate Coefficients for

$He^+ + He \rightarrow \sigma^+$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

He^+ Temp. (eV)	Equal Temp.	He Temp. (eV) 5000.	6000.	8000.	10000.	12000.	15000.	20000.
1.0E+03	8.49E-10#	1.73E-08#	2.12E-08*	2.78E-08*	3.30E-08*	3.71E-08*	4.20E-08*	4.76E-08*
1.5E+03	3.93E-09#	1.93E-08*	2.30E-08*	2.92E-08*	3.41E-08*	3.81E-08*	4.26E-08*	4.81E-08*
2.0E+03	8.30E-09#	2.12E-08*	2.47E-08*	3.06E-08*	3.52E-08*	3.89E-08*	4.33E-08*	4.85E-08*
2.5E+03	1.29E-08#	2.30E-08*	2.63E-08*	3.18E-08*	3.62E-08*	3.97E-08*	4.39E-08*	4.90E-08*
3.0E+03	1.73E-08#	2.47E-08*	2.78E-08*	3.30E-08*	3.71E-08*	4.05E-08*	4.45E-08*	4.94E-08*
3.5E+03	2.12E-08*	2.63E-08*	2.92E-08*	3.41E-08*	3.81E-08*	4.13E-08*	4.51E-08*	4.98E-08*
4.0E+03	2.47E-08*	2.78E-08*	3.06E-08*	3.52E-08*	3.89E-08*	4.20E-08*	4.56E-08*	5.02E-08
4.5E+03	2.78E-08*	2.92E-08*	3.18E-08*	3.62E-08*	3.97E-08*	4.26E-08*	4.62E-08*	5.06E-08
5.0E+03	3.06E-08*	3.06E-08*	3.30E-08*	3.71E-08*	4.05E-08*	4.33E-08*	4.67E-08*	5.09E-08
5.5E+03	3.30E-08*	3.18E-08*	3.41E-08*	3.81E-08*	4.13E-08*	4.39E-08*	4.72E-08*	5.13E-08
6.0E+03	3.52E-08*	3.30E-08*	3.52E-08*	3.89E-08*	4.20E-08*	4.45E-08*	4.76E-08*	5.16E-08
8.0E+03	4.20E-08*	3.71E-08*	3.89E-08*	4.20E-08*	4.45E-08*	4.67E-08*	4.94E-08*	5.29E-08
1.0E+04	4.67E-08*	4.05E-08*	4.20E-08*	4.45E-08*	4.67E-08*	4.85E-08*	5.09E-08	5.40E-08
1.2E+04	5.02E-08	4.33E-08*	4.45E-08*	4.67E-08*	4.85E-08*	5.02E-08	5.23E-08	5.51E-08
1.4E+04	5.29E-08	4.56E-08*	4.67E-08*	4.85E-08*	5.02E-08	5.16E-08	5.35E-08	5.60E-08
1.6E+04	5.51E-08	4.76E-08*	4.85E-08*	5.02E-08	5.16E-08	5.29E-08	5.46E-08	5.69E-08
1.8E+04	5.69E-08	4.94E-08*	5.02E-08	5.16E-08	5.29E-08	5.40E-08	5.56E-08	5.77E-08
2.0E+04	5.84E-08	5.09E-08	5.16E-08	5.29E-08	5.40E-08	5.51E-08	5.65E-08	5.84E-08

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+03 eV, E_{max} = 2.0E+04 eV

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
5000.	-34.6840	.561425	.00846562	-.0219365	-2.039967E-04	.00120158	-1.325802E-05
6000.	-34.4871	.460985	.0203888	-.0170479	-.00123903	8.950819E-04	7.412802E-05
8000.	-34.2085	.331032	.0296262	-.0101056	-.00187431	4.167700E-04	1.213301E-04
10000.	-34.0188	.252148	.0310288	-.00582488	-.00180584	1.394312E-04	1.072709E-04
12000.	-33.8797	.200089	.0298230	-.00316385	-.00155179	-8.216817E-06	8.115796E-05
15000.	-33.7274	.149363	.0266925	-8.769851E-04	-.00114857	-1.026825E-04	4.607838E-05
20000.	-33.5570	.101222	.0215477	7.809161E-04	-6.525756E-04	-1.278772E-04	1.216316E-05
Equal Temp.	-36.1071	1.93272	-.683257	.177855	-.0338898	.00493571	-5.499191E-04

See appendix for Chebyshev fit details.

$He^+ + He \rightarrow \sigma^+$



Ionization Rate Coefficients for

$He + He^+ \rightarrow \sigma^+$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^+ Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	6.24E-08	6.86E-08	7.45E-08	7.79E-08	8.08E-08	8.07E-08	6.16E-08
2.0E+00	6.24E-08	6.86E-08	7.44E-08	7.79E-08	8.08E-08	8.07E-08	6.16E-08
4.0E+00	6.24E-08	6.86E-08	7.44E-08	7.79E-08	8.08E-08	8.07E-08	6.16E-08
7.0E+00	6.24E-08	6.86E-08	7.44E-08	7.79E-08	8.08E-08	8.07E-08	6.16E-08
1.0E+01	6.24E-08	6.86E-08	7.44E-08	7.79E-08	8.08E-08	8.07E-08	6.16E-08
2.0E+01	6.23E-08	6.86E-08	7.44E-08	7.79E-08	8.08E-08	8.07E-08	6.16E-08
4.0E+01	6.23E-08	6.85E-08	7.44E-08	7.79E-08	8.08E-08	8.07E-08	6.16E-08
7.0E+01	6.23E-08	6.85E-08	7.44E-08	7.79E-08	8.08E-08	8.06E-08	6.16E-08
1.0E+02	6.23E-08	6.85E-08	7.44E-08	7.79E-08	8.07E-08	8.06E-08	6.16E-08
2.0E+02	6.23E-08	6.85E-08	7.44E-08	7.79E-08	8.07E-08	8.05E-08	6.16E-08
4.0E+02	6.23E-08	6.85E-08	7.44E-08	7.79E-08	8.07E-08	8.05E-08	6.16E-08
7.0E+02	6.23E-08	6.85E-08	7.44E-08	7.80E-08	8.06E-08	8.04E-08	6.16E-08
1.0E+03	6.23E-08	6.86E-08	7.43E-08	7.80E-08	8.06E-08	8.03E-08	6.16E-08
2.0E+03	6.23E-08	6.86E-08	7.43E-08	7.80E-08	8.05E-08	8.01E-08	6.16E-08
4.0E+03	6.21E-08	6.87E-08	7.43E-08	7.81E-08	8.04E-08	7.98E-08	6.16E-08
7.0E+03	6.19E-08	6.88E-08	7.43E-08	7.81E-08	8.02E-08	7.95E-08	6.16E-08
1.0E+04	6.19E-08	6.88E-08	7.43E-08	7.81E-08	8.01E-08	7.93E-08	6.16E-08
2.0E+04	6.28E-08	6.90E-08	7.44E-08	7.82E-08	7.99E-08	7.87E-08	6.16E-08

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-33.1826	-8.447088E-04	4.379986E-04	.00153621	.00196721	.00191297	.00165579
20000.	-32.9886	.00246657	.00200537	.00110670	3.142096E-04	-7.069038E-05	-9.257167E-05
40000.	-32.8276	-7.087943E-04	1.614074E-04	4.588880E-04	3.529485E-04	1.836710E-04	9.755263E-05
70000.	-32.7333	.00154471	7.127662E-04	1.336512E-04	-9.932082E-05	-1.253244E-04	-7.595796E-05
100000.	-32.6686	-.00490652	-.00228784	-7.224299E-04	-1.313023E-04	-2.878208E-05	1.350276E-05
200000.	-32.6778	-.0107031	-.00545996	-.00203612	-5.571363E-04	-8.996061E-05	1.739583E-05
500000.	-33.2050	-2.649854E-04	-1.908228E-04	-1.196480E-04	-6.719115E-05	-3.770261E-05	-2.551690E-05

See appendix for Chebyshev fit details.

$He + He^+ \rightarrow \sigma^+$



Ionization Rate Coefficients for

$He^+ + He \rightarrow \sigma^-$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

He^+ Temp. (eV)	Equal Temp.	He Temp. (eV) 5000.	6000.	8000.	10000.	12000.	15000.	20000.
1.0E+03	2.40E-11#	7.15E-10*	9.55E-10*	1.46E-09*	1.99E-09*	2.54E-09	3.39E-09	4.87E-09
1.5E+03	1.23E-10#	8.33E-10*	1.08E-09*	1.59E-09*	2.13E-09*	2.68E-09	3.54E-09	5.02E-09
2.0E+03	2.86E-10#	9.55E-10*	1.20E-09*	1.73E-09*	2.27E-09*	2.82E-09	3.68E-09	5.17E-09
2.5E+03	4.89E-10*	1.08E-09*	1.33E-09*	1.86E-09*	2.40E-09	2.97E-09	3.83E-09	5.32E-09
3.0E+03	7.15E-10*	1.20E-09*	1.46E-09*	1.99E-09*	2.54E-09	3.11E-09	3.98E-09	5.48E-09
3.5E+03	9.55E-10*	1.33E-09*	1.59E-09*	2.13E-09*	2.68E-09	3.25E-09	4.12E-09	5.63E-09
4.0E+03	1.20E-09*	1.46E-09*	1.73E-09*	2.27E-09*	2.82E-09	3.39E-09	4.27E-09	5.78E-09
4.5E+03	1.46E-09*	1.59E-09*	1.86E-09*	2.40E-09	2.97E-09	3.54E-09	4.42E-09	5.93E-09
5.0E+03	1.73E-09*	1.73E-09*	1.99E-09*	2.54E-09	3.11E-09	3.68E-09	4.57E-09	6.09E-09
5.5E+03	1.99E-09*	1.86E-09*	2.13E-09*	2.68E-09	3.25E-09	3.83E-09	4.72E-09	6.24E-09
6.0E+03	2.27E-09*	1.99E-09*	2.27E-09*	2.82E-09	3.39E-09	3.98E-09	4.87E-09	6.40E-09
8.0E+03	3.39E-09	2.54E-09	2.82E-09	3.39E-09	3.98E-09	4.57E-09	5.48E-09	7.02E-09
1.0E+04	4.57E-09	3.11E-09	3.39E-09	3.98E-09	4.57E-09	5.17E-09	6.09E-09	7.64E-09
1.2E+04	5.78E-09	3.68E-09	3.98E-09	4.57E-09	5.17E-09	5.78E-09	6.71E-09	8.27E-09
1.4E+04	7.02E-09	4.27E-09	4.57E-09	5.17E-09	5.78E-09	6.40E-09	7.33E-09	8.91E-09
1.6E+04	8.27E-09	4.87E-09	5.17E-09	5.78E-09	6.40E-09	7.02E-09	7.96E-09	9.54E-09
1.8E+04	9.54E-09	5.48E-09	5.78E-09	6.40E-09	7.02E-09	7.64E-09	8.59E-09	1.02E-08
2.0E+04	1.08E-08	6.09E-09	6.40E-09	7.02E-09	7.64E-09	8.27E-09	9.22E-09	1.08E-08

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+03 eV, E_{max} = 2.0E+04 eV

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
5000.	-40.2457	1.08016	.137224	-.00960777	-.00225681	.00426467	-3.707387E-05
6000.	-39.9198	.955399	.144445	-.00462863	-.00297896	1.656917E-04	3.638382E-05
8000.	-39.4021	.782041	.145020	.00249323	-.00313214	-2.249516E-04	6.938838E-05
10000.	-38.9951	.665607	.138788	.00682158	-.00268049	-4.285723E-04	4.558085E-05
12000.	-38.6575	.580938	.130790	.00942419	-.00211989	-5.139889E-04	1.312445E-05
15000.	-38.2374	.488850	.118758	.0114760	-.00136912	-5.313647E-04	-2.876352E-05
20000.	-37.6842	.386897	.101410	.0124848	-5.096408E-04	-4.658120E-04	-7.014682E-05
Equal Temp.	-41.6708	2.88394	-.522363	.167306	-.0405614	.00499358	-2.402961E-04

See appendix for Chebyshev fit details.

$He^+ + He \rightarrow \sigma^-$



Ionization Rate Coefficients for

$He + He^{+} \rightarrow \sigma^{-}$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^+ Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	400000.
1.0E+00	8.99E-09	3.51E-09	2.12E-08	5.88E-08	8.73E-08	9.38E-08	8.59E-08
2.0E+00	7.46E-09	9.85E-09	2.88E-08	5.91E-08	8.45E-08	9.38E-08	8.58E-08
4.0E+00	6.44E-09	1.40E-08	3.24E-08	5.84E-08	8.03E-08	9.38E-08	8.58E-08
7.0E+00	6.22E-09	1.48E-08	3.30E-08	5.84E-08	7.95E-08	9.37E-08	8.58E-08
1.0E+01	6.20E-09	1.48E-08	3.30E-08	5.84E-08	7.94E-08	9.37E-08	8.58E-08
2.0E+01	6.20E-09	1.48E-08	3.30E-08	5.84E-08	7.94E-08	9.37E-08	8.58E-08
4.0E+01	6.21E-09	1.48E-08	3.30E-08	5.84E-08	7.94E-08	9.37E-08	8.58E-08
7.0E+01	6.22E-09	1.49E-08	3.30E-08	5.84E-08	7.94E-08	9.36E-08	8.58E-08
1.0E+02	6.23E-09	1.49E-08	3.30E-08	5.84E-08	7.93E-08	9.36E-08	8.58E-08
2.0E+02	6.26E-09	1.49E-08	3.30E-08	5.84E-08	7.93E-08	9.35E-08	8.58E-08
4.0E+02	6.33E-09	1.49E-08	3.31E-08	5.84E-08	7.92E-08	9.34E-08	8.58E-08
7.0E+02	6.43E-09	1.50E-08	3.31E-08	5.84E-08	7.91E-08	9.33E-08	8.58E-08
1.0E+03	6.53E-09	1.51E-08	3.32E-08	5.84E-08	7.90E-08	9.32E-08	8.57E-08
2.0E+03	6.87E-09	1.55E-08	3.35E-08	5.86E-08	7.87E-08	9.30E-08	8.57E-08
4.0E+03	7.56E-09	1.62E-08	3.41E-08	5.89E-08	7.83E-08	9.27E-08	8.56E-08
7.0E+03	8.57E-09	1.72E-08	3.50E-08	5.94E-08	7.78E-08	9.22E-08	8.51E-08
1.0E+04	9.57E-09	1.82E-08	3.58E-08	5.99E-08	7.74E-08	9.20E-08	8.39E-08
2.0E+04	1.29E-08	2.16E-08	3.87E-08	6.11E-08	7.66E-08	9.13E-08	7.91E-08

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-37.3767	.152586	.271251	.0286627	.0720550	-.00417889	1.885647E-04
20000.	-36.2921	.498746	-.223651	.296885	-.142577	.103860	-.0465049
40000.	-34.5146	.168467	-.0600577	.0991481	-.0398571	.0325305	-.0123581
70000.	-33.2915	.0119357	.0143042	.00553446	.00283829	.00124894	-.00136315
100000.	-32.6810	-.0457164	.0165590	-.0198832	.00750636	-.00146373	-.00200726
200000.	-32.3792	-.0113032	-.00566531	-.00204888	-5.329708E-04	-7.815125E-05	1.640098E-05
400000.	-32.5635	-.0211742	-.0170066	-.0140215	-.00943770	-.00589724	-.00293363

See appendix for Chebyshev fit details.

$He + He^{+} \rightarrow \sigma^{-}$



Single and Double Ionization Cross Sections for He^+ in He

$$He^+ + He \rightarrow He^+ + He^+ + e$$
$$\rightarrow He^+ + He^{2+} + 2e$$

$He^+ + He^+ + e$		$He^+ + He^{2+} + 2e$	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
1.2E+04	4.09E-16	1.2E+04	2.16E-18
1.5E+04	3.57E-16	1.5E+04	2.87E-18
2.0E+04	3.07E-16	2.0E+04	3.88E-18
3.0E+04	2.52E-16	3.0E+04	5.82E-18
4.0E+04	2.19E-16	4.0E+04	7.76E-18
6.0E+04	1.80E-16	6.0E+04	9.74E-18
8.0E+04	1.54E-16	8.0E+04	9.08E-18
1.0E+05	1.36E-16	1.0E+05	7.69E-18
1.5E+05	1.08E-16	1.5E+05	4.95E-18
2.0E+05	9.12E-17	2.0E+05	3.32E-18
3.0E+05	7.02E-17	3.0E+05	1.81E-18
4.0E+05	5.85E-17	4.0E+05	1.19E-18
6.0E+05	4.53E-17	6.0E+05	7.14E-19
7.8E+05	3.82E-17	7.5E+05	5.60E-19

<u>References:</u> 388, 462, 463, 494, 495, 496

<u>Accuracy:</u> 20%

<u>Note:</u> At lower energies the formation of He^+ is dominated by single electron capture.

Chebyshev Fitting Parameters for Cross Sections

$He^+ + He^+ + e$ E_{min} = 1.2E+04 eV/amu, E_{max} = 7.8E+05 eV/amu
$He^+ + He^{2+} + 2e$ E_{min} = 1.2E+04 eV/amu, E_{max} = 7.5E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
$He^+ + He^+ + e$	-73.1564	-1.17268	-.0534400	-.00896703	.0130892	-.00303440	.000546481
$He^+ + He^{2+} + 2e$	-80.9940	-.809711	-.972278	.157307	.149794	-.0339390	-.0320888

The fit represents the $He^+ + He^+ + e$ cross section with an rms deviation of 0.2%. The maximum deviation is 0.3% at 1.0E+05 eV/amu.

The fit represents the $He^+ + He^{2+} + 2e$ cross section with an rms deviation of 2.0%. The maximum deviation is 3.3% at 3.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + He \rightarrow He^+ + He^+ + e \quad \sigma(He^+)$
$He^+ + He \rightarrow He^+ + He^{2+} + 2e \quad \sigma(He^{2+})$



Single Ionization Rate Coefficients for

$He + He^+ \rightarrow He^+ + He^+ + e$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^+ Temp. (eV)	He Energy (eV/amu) 15000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	6.07E-08	6.03E-08	6.08E-08	6.08E-08	5.97E-08	5.66E-08	4.99E-08
2.0E+00	6.08E-08	6.03E-08	6.08E-08	6.08E-08	5.97E-08	5.66E-08	4.99E-08
4.0E+00	6.08E-08	6.03E-08	6.08E-08	6.08E-08	5.97E-08	5.66E-08	4.99E-08
7.0E+00	6.08E-08	6.03E-08	6.08E-08	6.08E-08	5.97E-08	5.66E-08	4.99E-08
1.0E+01	6.08E-08	6.03E-08	6.08E-08	6.08E-08	5.97E-08	5.66E-08	4.99E-08
2.0E+01	6.08E-08	6.03E-08	6.08E-08	6.08E-08	5.97E-08	5.66E-08	4.99E-08
4.0E+01	6.08E-08	6.03E-08	6.08E-08	6.08E-08	5.97E-08	5.66E-08	4.99E-08
7.0E+01	6.08E-08	6.03E-08	6.08E-08	6.08E-08	5.97E-08	5.66E-08	4.99E-08
1.0E+02	6.08E-08	6.04E-08	6.08E-08	6.08E-08	5.97E-08	5.66E-08	4.99E-08
2.0E+02	6.06E-08	6.04E-08	6.08E-08	6.08E-08	5.97E-08	5.66E-08	4.99E-08
4.0E+02	5.91E-08	6.04E-08	6.09E-08	6.08E-08	5.97E-08	5.66E-08	4.99E-08
7.0E+02	5.65E-08	6.04E-08	6.09E-08	6.08E-08	5.97E-08	5.66E-08	4.99E-08
1.0E+03	5.45E-08*	6.04E-08	6.08E-08	6.08E-08	5.97E-08	5.66E-08	4.99E-08
2.0E+03	5.05E-08*	5.96E-08	6.09E-08	6.08E-08	5.97E-08	5.65E-08	4.99E-08
4.0E+03	4.73E-08*	5.73E-08	6.09E-08	6.07E-08	5.97E-08	5.65E-08	4.99E-08
7.0E+03	4.59E-08*	5.49E-08	6.08E-08	6.07E-08	5.96E-08	5.64E-08	4.99E-08
1.0E+04	4.55E-08*	5.36E-08*	6.07E-08	6.06E-08	5.96E-08	5.64E-08	4.99E-08
2.0E+04	4.61E-08*	5.18E-08*	5.98E-08	6.03E-08	5.95E-08	5.63E-08	4.99E-08

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
15000.	-33.4287	-.158108	-.0677688	.0633240	.0303214	.0168471	-.00235247
20000.	-33.3167	-.0613195	-.0440891	-.0209720	-.00324850	.00492917	.00599209
40000.	-33.2337	-.00382566	-.00347023	-.00301869	-.00227878	-.00145508	-8.331803E-04
70000.	-33.2330	-.00284092	-.00208988	-.00125435	-6.089006E-04	-2.360207E-04	-6.070840E-05
100000.	-33.2684	-.00150867	-9.658744E-04	-5.35351E-04	-2.92897E-04	-1.57790E-04	-6.49004E-05
200000.	-33.3763	-.00257819	-.00151167	-4.93080E-04	-2.11497E-04	5.23848E-05	-7.73477E-06
500000.	-33.6268	-2.54059E-04	-1.83372E-04	-1.19077E-04	-7.06237E-05	-4.19110E-05	-3.05576E-05

See appendix for Chebyshev fit details

$He + He^+ \rightarrow He^+ + He^+ + e$



Double Ionization Rate Coefficients for

$$He + He^{+} \rightarrow He^{+} + He^{2+} + e$$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^{+} Temp. (eV)	He Energy (eV/amu) 15000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	4.90E-10	7.62E-10	2.15E-09	3.45E-09	3.38E-09	2.06E-09	8.82E-10
2.0E+00	4.88E-10	7.62E-10	2.15E-09	3.45E-09	3.38E-09	2.06E-09	8.82E-10
4.0E+00	4.88E-10	7.62E-10	2.15E-09	3.45E-09	3.38E-09	2.06E-09	8.82E-10
7.0E+00	4.88E-10	7.62E-10	2.15E-09	3.45E-09	3.38E-09	2.06E-09	8.82E-10
1.0E+01	4.88E-10	7.62E-10	2.15E-09	3.45E-09	3.37E-09	2.06E-09	8.82E-10
2.0E+01	4.88E-10	7.62E-10	2.15E-09	3.45E-09	3.37E-09	2.06E-09	8.82E-10
4.0E+01	4.88E-10	7.63E-10	2.15E-09	3.45E-09	3.37E-09	2.06E-09	8.82E-10
7.0E+01	4.88E-10	7.64E-10	2.15E-09	3.45E-09	3.37E-09	2.06E-09	8.82E-10
1.0E+02	4.89E-10	7.64E-10	2.15E-09	3.45E-09	3.37E-09	2.06E-09	8.82E-10
2.0E+02	4.89E-10	7.67E-10	2.14E-09	3.45E-09	3.36E-09	2.06E-09	8.82E-10
4.0E+02	4.86E-10	7.72E-10	2.14E-09	3.45E-09	3.36E-09	2.06E-09	8.82E-10
7.0E+02	4.81E-10	7.81E-10	2.14E-09	3.45E-09	3.35E-09	2.06E-09	8.82E-10
1.0E+03	4.80E-10	7.89E-10	2.14E-09	3.44E-09	3.35E-09	2.06E-09	8.82E-10
2.0E+03	4.91E-10*	8.12E-10	2.16E-09	3.42E-09	3.33E-09	2.06E-09	8.82E-10
4.0E+03	5.35E-10*	8.60E-10	2.19E-09	3.38E-09	3.30E-09	2.06E-09	8.82E-10
7.0E+03	6.13E-10*	9.36E-10	2.22E-09	3.32E-09	3.27E-09	2.06E-09	8.83E-10
1.0E+04	6.93E-10	1.01E-09	2.25E-09	3.26E-09	3.23E-09	2.06E-09	8.83E-10
2.0E+04	9.42E-10	1.25E-09	2.32E-09	3.12E-09	3.12E-09	2.05E-09	8.85E-10

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
15000.	-42.6506	.210697	.170191	.107504	.0517361	.0113452	-.00599787
20000.	-41.7921	.173873	.119280	.0639172	.0267485	.00833917	.00105484
40000.	-39.8862	.0257832	.0218619	.0122377	.00362207	-2.35724E-04	-9.66877E-04
70000.	-39.0084	-.0336772	-.0251731	-.0147490	-.00625255	-.00143355	2.795328E-04
100000.	-39.0462	-.0279022	-.0171564	-.00916918	-.00454775	-.00203657	-7.514507E-04
200000.	-40.0019	-.00189800	-8.33043E-04	-1.62792E-04	-2.09259E-04	-7.95723E-05	-1.42798E-04
500000.	-41.6965	9.43319E-04	7.29142E-04	4.97094E-04	2.96459E-04	1.50204E-04	6.26450E-05

See appendix for Chebyshev fit details.

$$He + He^{+} \rightarrow He^{+} + He^{2+} + 2e$$



Ionization Cross Sections for

$He^{2+} + H \rightarrow He^{2+} + H^{+} + e$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
3.1E+04	2.45E+08	2.31E-16
4.0E+04	2.78E+08	3.39E-16
5.0E+04	3.11E+08	4.05E-16
6.0E+04	3.40E+08	4.29E-16
7.0E+04	3.68E+08	4.22E-16
8.0E+04	3.93E+08	4.07E-16
9.0E+04	4.17E+08	3.90E-16
1.0E+05	4.39E+08	3.77E-16
1.5E+05	5.38E+08	3.04E-16
2.0E+05	6.21E+08	2.53E-16
3.0E+05	7.61E+08	1.90E-16
4.0E+05	8.78E+08	1.53E-16
5.0E+05	9.82E+08	1.28E-16
5.5E+05	1.03E+09	1.18E-16

References: 506, 507, 508, 509, 510, 522, 512, 653

Accuracy: 30%

Notes: (1) As far as we are aware only one experimental measurement has been reported (Ref. 510). Above 100 keV/amu the theoretical results of Crothers and McCann (Ref. 507) are within 10% of the experimental results. At energies less than 100 keV/amu the theoretical data (Refs. 506, 507, 508, 509, 511, 512) vary by as much as an order of magnitude.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 3.1E+04 eV/amu, E_{max} = 5.5E+05 eV/amu

A0	A1	A2	A3	A4	A5	A6
-71.9027	-.471110	-.344985	.128260	-.0459423	.00932704	.00279820

The fit represents the above cross section with an rms deviation of 0.5%. The maximum deviation is 0.8% at 9.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$He^{2+} + H \rightarrow He^{2+} + H^{+} + e$



Ionization Rate Coefficients for

$$H + He^{2+} \rightarrow H^{+} + He^{2+} + e$$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	H Energy (eV/amu) 35000.	40000.	60000.	80000.	100000.	200000.	500000.
1.0E+00	7.21E-08	9.41E-08	1.46E-07	1.60E-07	1.66E-07	1.57E-07	1.26E-07
2.0E+00	7.21E-08	9.40E-08	1.46E-07	1.60E-07	1.66E-07	1.57E-07	1.26E-07
4.0E+00	7.21E-08	9.40E-08	1.46E-07	1.60E-07	1.66E-07	1.57E-07	1.26E-07
7.0E+00	7.21E-08	9.39E-08	1.46E-07	1.60E-07	1.66E-07	1.57E-07	1.26E-07
1.0E+01	7.21E-08	9.39E-08	1.46E-07	1.60E-07	1.65E-07	1.57E-07	1.26E-07
2.0E+01	7.21E-08	9.38E-08	1.46E-07	1.60E-07	1.65E-07	1.57E-07	1.26E-07
4.0E+01	7.22E-08	9.37E-08	1.46E-07	1.60E-07	1.65E-07	1.57E-07	1.26E-07
7.0E+01	7.22E-08	9.35E-08	1.45E-07	1.60E-07	1.65E-07	1.57E-07	1.26E-07
1.0E+02	7.23E-08	9.35E-08	1.45E-07	1.60E-07	1.65E-07	1.57E-07	1.26E-07
2.0E+02	7.19E-08	9.33E-08	1.45E-07	1.60E-07	1.65E-07	1.57E-07	1.26E-07
4.0E+02	7.01E-08	9.31E-08	1.45E-07	1.60E-07	1.65E-07	1.57E-07	1.26E-07
7.0E+02	6.80E-08	9.29E-08	1.44E-07	1.60E-07	1.65E-07	1.57E-07	1.26E-07
1.0E+03	6.67E-08*	9.25E-08	1.44E-07	1.59E-07	1.65E-07	1.57E-07	1.25E-07
2.0E+03	6.54E-08*	9.10E-08	1.43E-07	1.59E-07	1.64E-07	1.57E-07	1.24E-07
4.0E+03	6.66E-08*	8.96E-08	1.42E-07	1.59E-07	1.64E-07	1.57E-07	1.18E-07
7.0E+03	6.96E-08*	8.96E-08	1.40E-07	1.58E-07	1.63E-07	1.56E-07	1.10E-07*
1.0E+04	7.26E-08*	9.04E-08*	1.38E-07	1.56E-07	1.62E-07	1.56E-07	1.04E-07*
2.0E+04	8.06E-08*	9.43E-08*	1.34E-07	1.53E-07	1.60E-07	1.56E-07	9.36E-08*

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

H Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
35000.	-32.8985	.00549401	.0364412	.0474517	.0313047	.00563515	-.00840841
40000.	-32.3844	-.0158075	6.87820E-04	.00939742	.0108456	.00801907	.00326444
60000.	-31.5217	-.0331828	-.0190037	-.00885517	-.00343877	-9.09107E-04	9.58875E-05
80000.	-31.3145	-.0148657	-.0105289	-.00656864	-.00357109	-.00159023	-5.08065E-04
100000.	-31.2432	-.0124344	-.00696173	-.00342119	-.00172670	-9.03335E-04	-4.39816E-04
200000.	-31.3365	-.00345476	-.00195539	-9.10531E-04	-3.44995E-04	-1.21204E-04	-2.21966E-05
500000.	-31.8962	-.105965	-.0779255	-.0443668	-.0153954	-4.90580E-04	.00568340

See appendix for Chebyshev fit details.

$$H + He^{2+} \rightarrow H^{+} + He^{2+} + e$$



Cross Sections for the Production of Slow Positive Ions and Electrons for He^{2+} Colliding with H_2 Molecules

$He^{2+} + H_2 \rightarrow \Sigma\ (H_2^+ \text{ and } H^+ \text{ Prod.})\ \sigma^+$
$\rightarrow \Sigma$ Electrons σ^-

$He^{2+} + H_2$ (σ^+)		$He^{2+} + H_2$ (σ^-)	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
2.0E+03	3.82E-16	2.5E+03	9.15E-17
4.0E+03	6.17E-16	4.0E+03	7.80E-17
7.0E+03	8.79E-16	5.0E+03	7.52E-17
1.0E+04	1.11E-15	7.0E+03	8.35E-17
1.5E+04	1.37E-15	1.0E+04	1.18E-16
2.0E+04	1.47E-15	1.5E+04	2.12E-16
2.3E+04	1.49E-15	2.0E+04	3.17E-16
4.0E+04	1.28E-15	4.0E+04	6.19E-16
7.0E+04	9.83E-16	7.0E+04	7.00E-16
1.0E+05	7.90E-16	1.0E+05	6.97E-16
1.5E+05	6.21E-16	1.5E+05	6.01E-16
2.0E+05	5.15E-16	2.0E+05	4.94E-16
3.0E+05	4.01E-16	4.0E+05	2.51E-16
		5.0E+05	1.90E-16

References: 95, 143, 144, 346, 479, 532, 533, 534

Accuracy: 25%

Notes: (1) The seven reactions producing slow ions and electrons are: (a) $He^{2+} + H_2 \rightarrow He^{2+} + H_2^+ + e$ (single ionization), (b) $He^{2+} + H_2 \rightarrow He^{2+} + H^+ + H + e$ (dissociative ionization), (c) $He^{2+} + H_2 \rightarrow He^{2+} + H^+ + H^+ + 2e$ (double ionization), (d) $He^{2+} + H_2 \rightarrow He^+ + H_2^+$ (single electron capture), (e) $He^{2+} + H_2 \rightarrow He^+ + H^+ + H$ (dissociative electron capture), (f) $He^{2+} + H_2 \rightarrow He^+ + H^+ + H^+ + e$ (electron capture with ionization), (g) $He^{2+} + H_2 \rightarrow He + H^+ + H^+$ (double electron capture). (2) Afrosimov et al. (Ref. 95) have made a detailed study of these reactions in the energy region 2.3×10^3 to 2.5×10^4 eV/amu. (3) The cross section for slow electron production (σ^-) is the sum of $\sigma(a) + \sigma(b) + 2\sigma(c) + \sigma(f)$. (4) The cross section for slow positive ion production is the sum of $\sigma(a) + \sigma(b) + 2\sigma(c) + \sigma(d) + \sigma(e) + 2\sigma(f) + 2\sigma(g)$.

Chebyshev Fitting Parameters for Cross Sections

$He^{2+} + H_2$ (σ^+) E_{min} = 2.0E+03 eV/amu, E_{max} = 3.0E+05 eV/amu
$He^{2+} + H_2$ (σ^-) E_{min} = 2.5E+03 eV/amu, E_{max} = 5.0E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
$He^{2+} + H_2$ (σ^+)	-69.8186	.0111387	-.631977	.0179570	.0939313	-.00776735	-.0310932
$He^{2+} + H_2$ (σ^-)	-72.0755	.794419	-.668306	-.447465	.196243	-.000576099	-.0629005

The fit represents the (σ^+) cross section with an rms deviation of 1.3%. The maximum deviation is 3.0% at 4.0E+04 eV/amu.

The fit represents the (σ^-) cross section with an rms deviation of 3.2%. The maximum deviation is 5.8% at 7.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$$He^{2+} + H_2 \rightarrow \Sigma(H_2^{+} + H^{+}) \quad \sigma^{+}$$
$$He^{2+} + H_2 \rightarrow \Sigma\, e \quad \sigma^{-}$$



Cross Sections for the Production of H^+ and H_2^+ Ions for He^{2+} Incident On H_2

$$He^{2+} + H_2 \rightarrow \sigma(H^+)$$
$$\rightarrow \sigma(H_2^+)$$

(Curve A) Total H_2^+ Prod.		(Curve B) Single Ion. H_2^+ Prod.		(Curve C) Total H^+ Prod.		(Curve D) Ion. H^+ Prod.	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
2.1E+03	9.68E-17	2.5E+03	9.97E-18	2.3E+03	2.17E-16	2.3E+03	1.66E-16
7.0E+03	6.30E-16	1.0E+04	5.30E-17	7.0E+03	1.91E-16	7.0E+03	1.12E-16
1.0E+04	7.92E-16	1.5E+04	1.36E-16	1.0E+04	2.15E-16	1.0E+04	1.29E-16
1.5E+04	9.00E-16	2.0E+04	2.90E-16	1.5E+04	2.51E-16	1.5E+04	1.60E-16
2.0E+04	9.95E-16	3.0E+04	1.50E-15	2.0E+04	2.86E-16	2.0E+04	1.79E-16
3.0E+04	1.48E-15	4.0E+04	3.20E-15	4.0E+04	4.53E-16	2.5E+04	1.90E-16
4.0E+04	3.18E-15	8.4E+04	5.19E-15	7.0E+04	6.35E-16	4.0E+04	1.76E-16
7.0E+04	5.07E-15	1.0E+05	5.18E-15	8.0E+04	6.43E-16	7.0E+04	1.36E-16
8.4E+04	5.21E-15	1.5E+05	4.77E-15	1.0E+05	6.32E-16	1.0E+05	1.06E-16
1.0E+05	5.18E-15	2.0E+05	4.26E-15	1.5E+05	5.59E-16	1.5E+05	7.51E-17
1.5E+05	4.80E-15	4.0E+05	2.97E-15	2.0E+05	4.87E-16	2.0E+05	5.46E-17
2.0E+05	4.29E-15	5.4E+05	2.50E-15	4.0E+05	3.20E-16	4.0E+05	2.33E-17
4.0E+05	2.97E-15			4.6E+05	2.92E-16	6.0E+05	1.50E-17
5.4E+05	2.50E-15						

References: 95, 346, 553

Accuracy: 25%

Notes: (1) The seven reactions producing reaction products H^+ and H_2^+ are: (a) $He^{2+} + H_2 \rightarrow He^{2+} + H_2^+ + e$, single ionization; (b) $He^{2+} + H_2 \rightarrow He^{2+} + H^+ + H + e$, dissociative ionization; (c) $He^{2+} + H_2 \rightarrow He^{2+} + H^+ + H^+ + 2e$, double ionization; (d) $He^{2+} + H_2 \rightarrow He^+ + H_2^+$, electron capture; (e) $He^{2+} + H_2 \rightarrow He^+ + H^+ + H$, dissociative electron capture; (f) $He^{2+} + H_2 \rightarrow He^+ + H^+ + H^+ + e$, electron capture with ionization; (g) $He^{2+} + H_2 \rightarrow He + H^+ + H^+$, double electron capture. (2) Afrosimov et al. (Ref. 96) have made detailed studies of these reactions. (3) Shown as curve A is the total cross section for producing H_2^+ and is the sum of $\sigma(a)$ and $\sigma(d)$. (4) Curve B is the single ionization cross section for producing H_2^+, $\sigma(a)$. (5) Curve C is the total cross section for producing H^+ and is the sum of $\sigma(b) + 2\sigma(c) + \sigma(e) + 2\sigma(f)$. (6) Curve D is the cross section for producing H^+ by ionization collisions, $\sigma(b) + 2\sigma(c)$.

Chebyshev Fitting Parameters for Cross Sections

Curve A	E_{min} =	2.1E+03	eV/amu,	E_{max} =	5.4E+05	eV/amu
Curve B	E_{min} =	2.5E+03	eV/amu,	E_{max} =	5.4E+05	eV/amu
Curve C	E_{min} =	2.3E+03	eV/amu,	E_{max} =	4.6E+05	eV/amu
Curve D	E_{min} =	2.3E+03	eV/amu,	E_{max} =	6.0E+05	eV/amu

	A0	A1	A2	A3	A4	A5	A6
Curve A	-68.8191	1.63697	-.768285	-.163048	-.111162	.206844	.0545283
Curve B	-70.9500	3.23148	-1.12038	-.650292	.397513	.189249	-.208238
Curve C	-71.4299	.407648	-.257736	-.323208	.0419822	.0784881	.0252954
Curve D	-74.1448	-.984730	-.626797	-.217335	.186783	-.0134584	-.0263402

The fit represents the Curve A cross section with an rms deviation of 12.0%.
The maximum deviation is 23.3% at 3.0E+04 eV/amu.
The fit represents the Curve B cross section with an rms deviation of 12.7%.
The maximum deviation is 31.9% at 2.0E+04 eV/amu.
The fit represents the Curve C cross section with an rms deviation of 2.3%.
The maximum deviation is 3.3% at 1.5E+05 eV/amu.
The fit represents the Curve D cross section with an rms deviation of 3.2%.
The maximum deviation is 4.4% at 7.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$He^{2+} + H_2 \rightarrow \sigma(H^+), \sigma(H_2^+)$



Cross Sections for the Production of Slow Positive Ions and Electrons for He^{2+} Ions Interacting with He.

He^{2+} + He -> σ^+
He^{2+} + He -> σ^-

He^{2+} + He (σ^+)		He^{2+} + He (σ^-)	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
9.0E+03	3.72E-16	5.0E+03	1.59E-18
1.0E+04	4.03E-16	7.0E+03	4.68E-18
1.5E+04	5.30E-16	1.0E+04	1.29E-17
2.0E+04	5.87E-16	1.5E+04	3.07E-17
3.0E+04	6.03E-16	2.0E+04	5.11E-17
4.0E+04	5.91E-16	4.0E+04	1.30E-16
7.0E+04	4.98E-16	7.0E+04	2.20E-16
1.0E+05	4.19E-16	1.0E+05	2.74E-16
1.5E+05	3.29E-16	1.3E+05	2.88E-16
2.0E+05	2.73E-16	1.5E+05	2.84E-16
4.0E+05	1.70E-16	2.0E+05	2.51E-16
7.0E+05	1.14E-16	4.0E+05	1.62E-16
1.0E+06	8.54E-17	7.0E+05	1.06E-16
1.5E+06	6.10E-17	1.0E+06	7.96E-17
2.0E+06	4.74E-17	1.5E+06	5.57E-17
2.9E+06	3.41E-17	2.0E+06	4.31E-17
		2.8E+06	3.29E-17

<u>References:</u> 143, 144, 148, 375, 381, 382, 479, 495, 532, 534, 537, 538, 539, 540

<u>Accuracy:</u> σ^+ - 30%
σ^- - 50%

<u>Notes:</u> (1) The reactions occurring when He^{2+} reacts with He are: (a) He^{2+} + He -> He + He^{2+}, double electron capture; (b) He^{2+} + He -> He^+ + He^+, single electron capture; (c) He^{2+} + He -> He^{2+} + He^{2+} + 2e, double ionization; (d) He^{2+} + He -> He^{2+} + He^+ + e, single ionization; (e) He^{2+} + He -> He^+ + He^{2+} + e, single capture with single ionization. The cross sections for slow electron production σ^- is $2\sigma(c) + \sigma(d) + \sigma(e)$. The cross sections for slow ion production σ^+ is $\sigma(a) + \sigma(b) + \sigma(c) + \sigma(d) + \sigma(e)$. (2) For energies less than 10^5 eV/amu electron capture dominates overionization.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

σ^+ E_{min} = 9.0E+03 eV/amu, E_{max} = 2.9E+06 eV/amu
σ^- E_{min} = 5.0E+03 eV/amu, E_{max} = 2.8E+06 eV/amu

	A0	A1	A2	A3	A4	A5	A6
σ^+	-72.3312	-1.34390	-.519938	.150932	-.0442699	-.00102101	.00498217
σ^-	-75.4418	1.18618	-1.77854	.310230	.0670546	.0236438	-.0500356

The fit represents the σ^+ cross section with an rms deviation of 0.6%. The maximum deviation is 1.3% at 3.0E+04 eV/amu.

The fit represents the σ^- cross section with an rms deviation of 2.9%. The maximum deviation is 5.1% at 7.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$He^{2+} + He \rightarrow \sigma^-$
$He^{2+} + He \rightarrow \sigma^+$



Slow Positive Ion Production Rate Coefficients for

$He^{2+} + He \rightarrow \sigma^{+}$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	Equal Temp.	He Temp. (eV) 5000.	6000.	8000.	10000.	12000.	15000.	20000.
3.5E+03	1.01E-09#	2.40E-09#	3.68E-09#	6.95E-09#	1.09E-08#	1.53E-08#	2.22E-08#	3.37E-08*
4.0E+03	1.86E-09#	3.01E-09#	4.42E-09#	7.89E-09#	1.20E-08#	1.64E-08#	2.33E-08*	3.48E-08*
4.5E+03	3.01E-09#	3.68E-09#	5.22E-09#	8.86E-09#	1.31E-08#	1.75E-08#	2.45E-08*	3.59E-08*
5.0E+03	4.42E-09#	4.42E-09#	6.06E-09#	9.86E-09#	1.42E-08#	1.87E-08#	2.57E-08*	3.70E-08*
5.5E+03	6.06E-09#	5.22E-09#	6.95E-09#	1.09E-08#	1.53E-08#	1.99E-08#	2.68E-08*	3.81E-08*
6.0E+03	7.89E-09#	6.06E-09#	7.89E-09#	1.20E-08#	1.64E-08#	2.10E-08#	2.80E-08*	3.92E-08*
8.0E+03	1.64E-08#	9.86E-09#	1.20E-08#	1.64E-08#	2.10E-08#	2.57E-08*	3.26E-08*	4.35E-08*
1.0E+04	2.57E-08*	1.42E-08#	1.64E-08#	2.10E-08#	2.57E-08*	3.03E-08*	3.70E-08*	4.76E-08*
1.2E+04	3.48E-08*	1.87E-08#	2.10E-08#	2.57E-08*	3.03E-08*	3.48E-08*	4.14E-08*	5.15E-08*
1.4E+04	4.35E-08*	2.33E-08*	2.57E-08*	3.03E-08*	3.48E-08*	3.92E-08*	4.55E-08*	5.53E-08*
1.6E+04	5.15E-08*	2.80E-08*	3.03E-08*	3.48E-08*	3.92E-08*	4.35E-08*	4.95E-08*	5.89E-08*
1.8E+04	5.89E-08*	3.26E-08*	3.48E-08*	3.92E-08*	4.35E-08*	4.76E-08*	5.34E-08*	6.24E-08*
2.0E+04	6.57E-08*	3.70E-08*	3.92E-08*	4.35E-08*	4.76E-08*	5.15E-08*	5.71E-08*	6.57E-08*

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 3.5E+03 eV, E_{max} = 2.0E+04 eV

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
5000.	-36.8434	1.38333	-.0598328	-.0157120	.00186832	1.83603E-04	6.05353E-05
6000.	-36.4161	1.19746	-.0301418	-.0151798	9.78258E-04	1.15739E-04	-3.58181E-05
8000.	-35.7440	.928769	.00411079	-.0126285	1.55066E-04	2.07816E-04	1.37906E-04
10000.	-35.2350	.745842	.0204872	-.00979258	-3.75639E-04	2.38560E-04	-3.33682E-05
12000.	-34.8337	.615060	.0278881	-.00727889	-6.44358E-04	-3.66627E-05	4.09092E-05
15000.	-34.3648	.477864	.0321621	-.00474753	-6.70690E-04	1.26091E-04	-4.23094E-05
20000.	-33.8047	.336327	.0310835E-02	-.00214049	-6.33145E-04	-5.04160E-05	-1.09705E-04
Equal Temp.	-36.4417	2.02992	-.399477	.0593303	-.00723537	5.77470E-04	-3.03527E-05

See appendix for Chebyshev fit details.

$He^{2+} + He \rightarrow \sigma^{+}$



Slow Positive Ion Production Rate Coefficients for

$He + He^{2+} \rightarrow \sigma^{+}$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	5.60E-08	1.15E-07	1.64E-07	1.83E-07	1.84E-07	1.69E-07	1.42E-07
2.0E+00	5.60E-08	1.15E-07	1.64E-07	1.83E-07	1.84E-07	1.70E-07	1.42E-07
4.0E+00	5.60E-08	1.15E-07	1.64E-07	1.83E-07	1.84E-07	1.70E-07	1.42E-07
7.0E+00	5.60E-08	1.15E-07	1.64E-07	1.83E-07	1.84E-07	1.70E-07	1.42E-07
1.0E+01	5.60E-08	1.15E-07	1.64E-07	1.83E-07	1.84E-07	1.70E-07	1.42E-07
2.0E+01	5.60E-08	1.15E-07	1.64E-07	1.83E-07	1.84E-07	1.70E-07	1.42E-07
4.0E+01	5.55E-08	1.15E-07	1.64E-07	1.83E-07	1.84E-07	1.70E-07	1.42E-07
7.0E+01	5.42E-08	1.15E-07	1.64E-07	1.83E-07	1.84E-07	1.70E-07	1.42E-07
1.0E+02	5.29E-08	1.15E-07	1.64E-07	1.83E-07	1.84E-07	1.70E-07	1.42E-07
2.0E+02	4.99E-08*	1.15E-07	1.63E-07	1.83E-07	1.84E-07	1.70E-07	1.42E-07
4.0E+02	4.75E-08*	1.14E-07	1.63E-07	1.82E-07	1.84E-07	1.69E-07	1.42E-07
7.0E+02	4.67E-08*	1.14E-07	1.63E-07	1.82E-07	1.83E-07	1.69E-07	1.42E-07
1.0E+03	4.68E-08*	1.14E-07	1.63E-07	1.82E-07	1.83E-07	1.69E-07	1.42E-07
2.0E+03	4.90E-08*	1.14E-07	1.62E-07	1.82E-07	1.83E-07	1.69E-07	1.42E-07
4.0E+03	5.40E-08*	1.15E-07	1.62E-07	1.81E-07	1.83E-07	1.69E-07	1.42E-07
7.0E+03	6.06E-08*	1.15E-07	1.61E-07	1.80E-07	1.82E-07	1.69E-07	1.42E-07
1.0E+04	6.62E-08*	1.16E-07	1.60E-07	1.80E-07	1.82E-07	1.69E-07	1.42E-07
2.0E+04	8.14E-08*	1.20E-07	1.59E-07	1.78E-07	1.80E-07	1.69E-07	1.42E-07

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-33.3528	.0888838	.145456	.115375	.0287771	-.0197504	-.00975561
20000.	-31.9486	.00733446	.0104567	.00858353	.00508869	.00265079	.00155039
40000.	-31.2633	-.0136989	-.00614936	-.00195506	-5.61861E-04	-2.03312E-04	-6.59434E-05
70000.	-31.0420	-.0112913	-.00592950	-.00253976	-.00106355	- 4.84457E-04	-2.29325E-04
100000.	-31.0255	-.00737792	-.00415937	-.00198048	-9.12662E-04	-4.31500E-04	-1.81183E-04
200000.	-31.1826	-.00193461	-.00125764	-4.52646E-04	-2.53650E-04	2.30595E-05	-3.75081E-05
500000.	-31.5305	-2.56726E-04	-1.78746E-04	-1.04195E-04	-5.27594E-05	-2.69854E-05	-1.83293E-05

See appendix for Chebyshev fit details

$He + He^{2+} \rightarrow \sigma^{+}$



Slow Electron Production Rate Coefficients for

$He^{2+} + He \rightarrow \sigma^-$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	Equal Temp.	He Temp. (eV) 5000.	6000.	8000.	10000.	12000.	15000.	20000.
1.5E+03	1.17E-12#	6.45E-11#	1.11E-10*	2.47E-10*	4.44E-10*	6.99E-10	1.18E-09	2.24E-09
2.0E+03	6.87E-12#	8.57E-11*	1.39E-10*	2.91E-10*	5.02E-10*	7.72E-10	1.28E-09	2.36E-09
2.5E+03	2.11E-11#	1.11E-10*	1.71E-10*	3.38E-10*	5.64E-10*	8.48E-10	1.37E-09	2.48E-09
3.0E+03	4.68E-11#	1.39E-10*	2.07E-10*	3.89E-10*	6.30E-10*	9.27E-10	1.47E-09	2.61E-09
3.5E+03	8.57E-11*	1.71E-10*	2.47E-10*	4.44E-10*	6.99E-10	1.01E-09	1.57E-09	2.73E-09
4.0E+03	1.39E-10*	2.07E-10*	2.91E-10*	5.02E-10*	7.72E-10	1.10E-09	1.68E-09	2.86E-09
4.5E+03	2.07E-10*	2.47E-10*	3.38E-10*	5.64E-10*	8.48E-10	1.18E-09	1.78E-09	3.00E-09
5.0E+03	2.91E-10*	2.91E-10*	3.89E-10*	6.30E-10*	9.27E-10	1.28E-09	1.89E-09	3.13E-09
5.5E+03	3.89E-10*	3.38E-10*	4.44E-10*	6.99E-10	1.01E-09	1.37E-09	2.00E-09	3.27E-09
6.0E+03	5.02E-10*	3.89E-10*	5.02E-10*	7.72E-10	1.10E-09	1.47E-09	2.12E-09	3.41E-09
8.0E+03	1.10E-09	6.30E-10*	7.72E-10	1.10E-09	1.47E-09	1.89E-09	2.61E-09	3.99E-09
1.0E+04	1.89E-09	9.27E-10	1.10E-09	1.47E-09	1.89E-09	2.36E-09	3.13E-09	4.60E-09
1.2E+04	2.86E-09	1.28E-09	1.47E-09	1.89E-09	2.36E-09	2.86E-09	3.69E-09	5.24E-09
1.4E+04	3.99E-09	1.68E-09	1.89E-09	2.36E-09	2.86E-09	3.41E-09	4.29E-09	5.91E-09
1.6E+04	5.24E-09	2.12E-09	2.36E-09	2.86E-09	3.41E-09	3.99E-09	4.92E-09	6.60E-09
1.8E+04	6.60E-09	2.61E-09	2.86E-09	3.41E-09	3.99E-09	4.60E-09	5.57E-09	7.32E-09
2.0E+04	8.06E-09	3.13E-09	3.41E-09	3.99E-09	4.60E-09	5.24E-09	6.25E-09	8.06E-09

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.5E+03 eV, E_{max} = 2.0E+04 eV

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
5000.	43.3317	1.97188	.145899	-.0314858	-.00292157	9.87310E-04	6.59233E-05
6000.	-42.7445	1.73787	.165098	-.0242892	-.00406700	7.29676E-04	1.18519E-04
8000.	-41.8072	1.40305	.177579	-.0125933	-.00472104	2.21088E-04	1.43982E-04
10000.	-41.0752	1.17389	.175396	-.00453799	-.00443211	-1.26246E-04	1.20206E-04
12000.	-40.4766	1.00680	.167979	8.55383E-04	-.00385784	-3.27458E-04	8.43893E-05
15000.	-39.7464	.826618	.154121	.00577291	-.00294670	-4.58815E-04	3.46633E-05
20000.	-38.8140	.631796	.131790	.00946407	-.00174398	-4.77146E-04	-1.78125E-05
Equal Temp.	-44.6706	4.27088	-.685712	.139921	-.0311302	.00607534	-8.51332E-04

See appendix for Chebyshev fit details.

$He^{2+} + He \rightarrow \sigma^-$

Maxwellian – Maxwellian



Slow Electron Production Rate Coefficients for

$He + He^{2+} \rightarrow \sigma^-$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	1.06E-09	1.04E-08	6.65E-08	3.41E-08	5.31E-08	1.56E-07	1.34E-07
2.0E+00	1.60E-09	1.00E-08	4.88E-08	6.09E-08	9.79E-08	1.56E-07	1.34E-07
4.0E+00	1.78E-09	1.00E-08	3.86E-08	7.70E-08	1.18E-07	1.56E-07	1.34E-07
7.0E+00	1.79E-09	1.00E-08	3.63E-08	8.04E-08	1.20E-07	1.56E-07	1.34E-07
1.0E+01	1.78E-09	1.00E-08	3.61E-08	8.07E-08	1.20E-07	1.56E-07	1.34E-07
2.0E+01	1.78E-09	1.00E-08	3.60E-08	8.07E-08	1.20E-07	1.56E-07	1.34E-07
4.0E+01	1.79E-09	1.00E-08	3.60E-08	8.07E-08	1.20E-07	1.56E-07	1.34E-07
7.0E+01	1.79E-09	1.00E-08	3.60E-08	8.07E-08	1.20E-07	1.56E-07	1.34E-07
1.0E+02	1.80E-09	1.00E-08	3.60E-08	8.06E-08	1.20E-07	1.56E-07	1.34E-07
2.0E+02	1.83E-09	1.00E-08	3.60E-08	8.06E-08	1.20E-07	1.56E-07	1.34E-07
4.0E+02	1.90E-09	1.01E-08	3.60E-08	8.06E-08	1.20E-07	1.56E-07	1.34E-07
7.0E+02	2.01E-09	1.03E-08	3.61E-08	8.06E-08	1.19E-07	1.55E-07	1.34E-07
1.0E+03	2.13E-09	1.04E-08	3.62E-08	8.06E-08	1.19E-07	1.55E-07	1.34E-07
2.0E+03	2.51E-09	1.09E-08	3.67E-08	8.09E-08	1.19E-07	1.55E-07	1.34E-07
4.0E+03	3.27E-09	1.20E-08	3.79E-08	8.16E-08	1.19E-07	1.55E-07	1.34E-07
7.0E+03	4.44E-09	1.37E-08	3.97E-08	8.28E-08	1.19E-07	1.54E-07	1.34E-07
1.0E+04	5.63E-09	1.54E-08	4.14E-08	8.38E-08	1.19E-07	1.54E-07	1.34E-07
2.0E+04	9.85E-09	2.08E-08	4.70E-08	8.69E-08	1.19E-07	1.52E-07	1.34E-07

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-39.6384	.794564	.318341	.275468	-.0163980	.0386510	-.0302765
20000.	-36.5385	.251533	.188617	.0914015	.0423640	.00564777	.00225398
40000.	-33.9830	-.0965962	.217818	-.0655465	.0766555	-.0163273	5.77974E-04
70000.	-32.8790	.252448	-.174234	.159512	-.0904273	.0541651	-.0228943
100000.	-32.0888	.196302	-.176198	.140458	-.0957365	.0592874	-.0318816
200000.	-31.3593	-.00872687	-.00490003	-.00214295	-9.76224E-04	-3.39591E-04	-1.98244E-04
500000.	-31.6466	-3.56481E-04	-2.64796E-04	-1.77709E-04	-1.07797E-04	-6.25087E-05	-4.02042E-05

See appendix for Chebyshev fit details.

$He + He^{2+} \rightarrow \sigma^-$

Beam – Maxwellian



Single and Double Ionization Cross Sections for He^{2+} + He Collisions

$$He^{2+} + He \rightarrow He^{2+} + He^{+} + e$$
$$He^{2+} + He \rightarrow He^{2+} + He^{2+} + 2e$$

Single Ionization		Double Ionization	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
1.5E+03	2.72E-18	1.4E+04	4.43E-19
2.0E+03	2.85E-18	1.5E+04	4.70E-19
4.0E+03	3.41E-18	2.0E+04	6.21E-19
7.0E+03	4.73E-18	4.0E+04	1.32E-18
1.0E+04	5.96E-18	7.0E+04	2.76E-18
1.5E+04	8.11E-18	1.0E+05	6.06E-18
2.0E+04	1.10E-17	1.4E+05	7.60E-18
4.0E+04	2.92E-17	1.5E+05	7.50E-18
7.0E+04	1.28E-16	2.0E+05	6.65E-18
1.0E+05	1.85E-16	4.0E+05	3.01E-18
1.5E+05	2.09E-16	7.0E+05	1.27E-18
2.0E+05	2.05E-16	1.0E+06	7.13E-19
4.0E+05	1.50E-16	1.5E+06	3.51E-19
7.0E+05	1.03E-16	2.0E+06	2.53E-19
1.0E+06	8.17E-17	2.7E+06	2.06E-19
1.5E+06	6.10E-17		
2.0E+06	4.97E-17		
2.4E+06	4.46E-17		

<u>References:</u> 148, 282, 381, 382, 383, 384, 387, 537, 538, 540

<u>Accuracy:</u> $\sigma(He^{+})$ - 30%
$\sigma(He^{2+})$ - 50%

<u>Note:</u> The data of DuBois (Ref. 538) supercede earlier results (Ref. 282) due to recalibration of the slow recoil ion detector. These corrected data are consistent with those of Afrosimov et al. (Ref. 148), which define the low-energy behavior of the cross section.

Chebyshev Fitting Parameters for Cross Sections

$\sigma(He^{+})$ E_{min} = 1.5E+03 eV/amu, E_{max} = 2.4E+06 eV/amu
$\sigma(He^{2+})$ E_{min} = 1.4E+04 eV/amu, E_{max} = 2.7E+06 eV/amu

	A0	A1	A2	A3	A4	A5	A6
$\sigma(He^{+})$	-76.5996	1.88840	-.913101	-.705245	.237700	.295143	-.101724
$\sigma(He^{2+})$	-82.9537	-.441854	-1.49781	.0576939	.413751	-.0249295	-.0956323

The fit represents the $\sigma(He^{+})$ cross section with an rms deviation of 15.4%. The maximum deviation is 44.5% at 4.0E+04 eV/amu.

The fit represents the $\sigma(He^{2+})$ cross section with an rms deviation of 8.4%. The maximum deviation is 18.4% at 7.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$He^{2+} + He \rightarrow He^{2+} + He^{+} + e \quad \sigma(He^{+})$
$He^{2+} + He \rightarrow He^{2+} + He^{2+} + 2e \quad \sigma(He^{2+})$



Single Ionization Rate Coefficients for

$He^{2+} + He \rightarrow He^{2+} + He^{+} + e$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	Equal Temp.	He Temp. (eV) 1000.	2000.	4000.	7000.	10000.	15000.	20000.
1.0E+03	2.02E-11#	2.02E-11#	5.26E-11#	1.21E-10*	2.21E-10*	3.23E-10	5.15E-10	7.50E-10
1.5E+03	5.26E-11#	3.57E-11#	6.99E-11*	1.38E-10*	2.38E-10*	3.41E-10	5.36E-10	7.76E-10
2.0E+03	8.72E-11*	5.26E-11#	8.72E-11*	1.55E-10*	2.54E-10*	3.59E-10	5.58E-10	8.04E-10
2.5E+03	1.21E-10*	6.99E-11*	1.04E-10*	1.71E-10*	2.71E-10*	3.77E-10	5.80E-10	8.31E-10
3.0E+03	1.55E-10*	8.72E-11*	1.21E-10*	1.88E-10*	2.88E-10	3.96E-10	6.03E-10	8.60E-10
3.5E+03	1.88E-10*	1.04E-10*	1.38E-10*	2.04E-10*	3.06E-10	4.15E-10	6.26E-10	8.89E-10
4.0E+03	2.21E-10*	1.21E-10*	1.55E-10*	2.21E-10*	3.23E-10	4.34E-10	6.50E-10	9.19E-10
4.5E+03	2.54E-10*	1.38E-10*	1.71E-10*	2.38E-10*	3.41E-10	4.54E-10	6.74E-10	9.50E-10
5.0E+03	2.88E-10	1.55E-10*	1.88E-10*	2.54E-10*	3.59E-10	4.74E-10	6.99E-10	9.81E-10
5.5E+03	3.23E-10	1.71E-10*	2.04E-10*	2.71E-10*	3.77E-10	4.94E-10	7.24E-10	1.01E-09
6.0E+03	3.59E-10	1.88E-10*	2.21E-10*	2.88E-10	3.96E-10	5.15E-10	7.50E-10	1.05E-09
8.0E+03	5.15E-10	2.54E-10*	2.88E-10	3.59E-10	4.74E-10	6.03E-10	8.60E-10	1.19E-09
1.0E+04	6.99E-10	3.23E-10	3.59E-10	4.34E-10	5.58E-10	6.99E-10	9.81E-10	1.34E-09
1.2E+04	9.19E-10	3.96E-10	4.34E-10	5.15E-10	6.50E-10	8.04E-10	1.11E-09	1.51E-09
1.4E+04	1.19E-09	4.74E-10	5.15E-10	6.03E-10	7.50E-10	9.19E-10	1.26E-09	1.69E-09
1.6E+04	1.51E-09	5.58E-10	6.03E-10	6.99E-10	8.60E-10	1.05E-09	1.42E-09	1.90E-09
1.8E+04	1.90E-09	6.50E-10	6.99E-10	8.04E-10	9.81E-10	1.19E-09	1.60E-09	2.12E-09
2.0E+04	2.36E-09	7.50E-10	8.04E-10	9.19E-10	1.11E-09	1.34E-09	1.79E-09	2.36E-09

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+03 eV, E_{max} = 2.0E+04 eV

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
1000.	-45.5517	1.76637	-.0538507	.0411721	.0121873	-.00162780	4.71335E-04
2000.	-44.8184	1.34639	.0940426	.0160336	.0100615	7.61162E-04	-5.25996E-06
4000.	-43.9954	.993020	.171815	.0181703	.00532378	.00150693	1.33413E-04
7000.	-43.2344	.780722	.188817	.0272646	.00407424	.00107015	2.33342E-04
10000.	-42.6662	.678316	.186986	.0320032	.00435983	7.84218E-04	1.42277E-04
15000.	-41.8920	.588941	.178679	.0346782	.00461614	3.94516E-04	-9.31127E-05
20000.	-41.2236	.538493	.169263	.0338729	.00410893	1.87221E-05	-2.35709E-04
Equal Temp.	-44.3188	2.21934	-.0710730	.154144	-.0123650	.00522272	-.00149471

See appendix for Chebyshev fit details.

$He^{2+} + He \rightarrow He^{2+} + He^{+} + e$

Maxwellian – Maxwellian



Single Ionization Rate Coefficients for

$He + He^{2+} \rightarrow He^{+} + He^{2+} + e$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	7.33E-11	1.69E-10	9.57E-09	1.61E-08	7.23E-08	1.27E-07	1.27E-07
2.0E+00	3.50E-10	9.12E-10	9.31E-09	3.37E-08	8.37E-08	1.27E-07	1.27E-07
4.0E+00	6.77E-10	1.78E-09	8.43E-09	4.44E-08	8.29E-08	1.27E-07	1.27E-07
7.0E+00	8.03E-10	2.10E-09	8.17E-09	4.66E-08	8.17E-08	1.27E-07	1.27E-07
1.0E+01	8.25E-10	2.16E-09	8.16E-09	4.68E-08	8.13E-08	1.27E-07	1.27E-07
2.0E+01	8.30E-10	2.17E-09	8.18E-09	4.67E-08	8.10E-08	1.27E-07	1.27E-07
4.0E+01	8.31E-10	2.18E-09	8.22E-09	4.66E-08	8.09E-08	1.27E-07	1.27E-07
7.0E+01	8.33E-10	2.18E-09	8.26E-09	4.64E-08	8.08E-08	1.27E-07	1.27E-07
1.0E+02	8.35E-10	2.19E-09	8.30E-09	4.63E-08	8.08E-08	1.27E-07	1.27E-07
2.0E+02	8.40E-10	2.20E-09	8.40E-09	4.61E-08	8.06E-08	1.27E-07	1.27E-07
4.0E+02	8.50E-10	2.23E-09	8.57E-09	4.59E-08	8.04E-08	1.27E-07	1.27E-07
7.0E+02	8.65E-10	2.28E-09	8.79E-09	4.57E-08	8.01E-08	1.27E-07	1.27E-07
1.0E+03	8.80E-10	2.32E-09	9.01E-09	4.56E-08	8.00E-08	1.26E-07	1.27E-07
2.0E+03	9.34E-10	2.46E-09	9.68E-09	4.56E-08	7.97E-08	1.26E-07	1.27E-07
4.0E+03	1.05E-09	2.73E-09	1.10E-08	4.61E-08	7.95E-08	1.26E-07	1.27E-07
7.0E+03	1.26E-09	3.20E-09	1.30E-08	4.71E-08	7.94E-08	1.25E-07	1.27E-07
1.0E+04	1.50E-09	3.78E-09	1.48E-08	4.81E-08	7.94E-08	1.24E-07	1.27E-07
2.0E+04	2.65E-09	6.33E-09	2.04E-08	5.15E-08	7.99E-08	1.23E-07	1.27E-07

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-42.0879	1.05416	-.278168	.571603	-.178692	.163417	-.0427310
20000.	-40.2183	1.05217	-.325687	.576861	-.203701	.175474	-.0514200
40000.	-36.7712	.292755	.267684	.0744621	.0454617	.00691932	-.0124274
70000.	-34.0338	.300282	-.207456	.207032	-.108795	.0715453	-.0312105
100000.	-32.6844	.00402251	-.0177396	.0232129	-.0175932	.0185648	-.0137519
200000.	-31.7707	-.0135888	-.00673208	-.00257568	-7.31948E-04	-2.50136E-04	-2.83646E-05
500000.	-31.7621	-4.22604E-04	-3.26343E-04	-2.33365E-04	-1.51845E-04	-9.21185E-05	-5.83670E-05

See appendix for Chebyshev fit details.

$He + He^{2+} \rightarrow He^{+} + He^{2+} + e$

Beam – Maxwellian



Double Ionization Rate Coefficients for

$He + He^{2+} \rightarrow He^{2+} + He^{2+} + e$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	He Energy (eV/amu) 15000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	9.35E-11	1.13E-10	1.86E-10	1.74E-09	1.73E-09	4.13E-09	2.09E-09
2.0E+00	8.19E-11	1.21E-10	3.04E-10	1.26E-09	2.49E-09	4.13E-09	2.09E-09
4.0E+00	8.00E-11	1.22E-10	3.59E-10	1.05E-09	2.65E-09	4.13E-09	2.09E-09
7.0E+00	8.00E-11	1.22E-10	3.67E-10	1.02E-09	2.65E-09	4.13E-09	2.09E-09
1.0E+01	8.01E-11	1.22E-10	3.67E-10	1.02E-09	2.65E-09	4.12E-09	2.09E-09
2.0E+01	7.98E-11	1.22E-10	3.67E-10	1.02E-09	2.65E-09	4.12E-09	2.09E-09
4.0E+01	7.81E-11	1.22E-10	3.68E-10	1.02E-09	2.64E-09	4.12E-09	2.09E-09
7.0E+01	7.50E-11	1.23E-10	3.68E-10	1.02E-09	2.63E-09	4.11E-09	2.09E-09
1.0E+02	7.25E-11*	1.23E-10	3.69E-10	1.03E-09	2.63E-09	4.11E-09	2.09E-09
2.0E+02	6.79E-11*	1.23E-10	3.70E-10	1.03E-09	2.62E-09	4.10E-09	2.10E-09
4.0E+02	6.47E-11*	1.24E-10	3.73E-10	1.04E-09	2.60E-09	4.09E-09	2.10E-09
7.0E+02	6.39E-11*	1.26E-10	3.76E-10	1.06E-09	2.59E-09	4.08E-09	2.10E-09
1.0E+03	6.45E-11*	1.27E-10	3.79E-10	1.07E-09	2.58E-09	4.07E-09	2.10E-09
2.0E+03	6.87E-11*	1.30E-10	3.90E-10	1.11E-09	2.56E-09	4.05E-09	2.10E-09
4.0E+03	7.87E-11*	1.37E-10	4.10E-10	1.18E-09	2.56E-09	4.01E-09	2.10E-09
7.0E+03	9.41E-11*	1.52E-10	4.43E-10	1.26E-09	2.56E-09	3.97E-09	2.10E-09
1.0E+04	1.10E-10*	1.69E-10	4.78E-10	1.34E-09	2.58E-09	3.94E-09	2.10E-09
2.0E+04	1.66E-10*	2.34E-10	6.12E-10	1.54E-09	2.62E-09	3.84E-09	2.10E-09

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
15000.	-46.3312	.153224	.279715	.156829	.0725151	.0242560	.00705852
20000.	-45.4317	.231432	.136168	.105908	.0379291	.0289022	.00177492
40000.	-43.4356	.345707	-.0276535	.185072	-.0448587	.0604026	-.0183190
70000.	-41.0648	.0136654	.223879	-.0488164	.0714513	-.0272212	.00699819
100000.	-39.6220	.0822767	-.0834305	.0811386	-.0492265	.0386945	-.0239545
200000.	-38.6471	-.0288702	-.0153708	-.00634634	-.00248827	-8.91443E-04	-4.33357E-04
500000.	-39.9659	.00135586	8.97190E-04	4.16825E-04	1.05292E-04	-2.86625E-05	-6.55320E-05

See appendix for Chebyshev fit details.

$$He + He^{2+} \rightarrow He^{2+} + He^{2+} + e$$

Beam – Maxwellian



Cross Sections for the Ionization of Li by Proton Impact

$$H^+ + Li \rightarrow H^+ + Li^+ + e$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.5E+04	2.20E+08	1.20E-15
4.0E+04	2.78E+08	7.56E-16
7.0E+04	3.68E+08	4.49E-16
1.0E+05	4.39E+08	3.22E-16
1.5E+05	5.38E+08	2.24E-16
2.0E+05	6.21E+08	1.73E-16
4.0E+05	8.78E+08	9.48E-17
7.0E+05	1.16E+09	5.79E-17
1.0E+06	1.39E+09	4.21E-17
1.5E+06	1.70E+09	2.98E-17
2.0E+06	1.96E+09	2.36E-17
2.3E+06	2.10E+09	2.12E-17

References: 232, 234, 737, 738

Accuracy: 30%

Note: Cross sections for double ionization and electron transfer ionization may be found in Ref. 234.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 2.5E+04 eV/amu, E_{max} = 2.3E+06 eV/amu

A0	A1	A2	A3	A4	A5	A6
-72.8380	-2.01716	.0360083	-.00414961	.00683897	.00418712	.00222666

The fit represents the above cross section with an rms deviation of 0.2%. The maximum deviation is 0.4% at 1.0E+06 eV/amu.

See appendix for Chebyshev fit details.

$$H^+ + Li \rightarrow H^+ + Li^+ + e$$



Ionization Rate Coefficients for

$Li + H^+ \rightarrow Li^+ + H^+ + e$

Beam - Maxwellian Rate Coefficients (cm^3/s)

H^+ Temp. (eV)	Li Energy (eV/amu) 25000.	30000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	1.32E-07*	2.41E-07	2.10E-07	1.65E-07	1.41E-07	1.07E-07	7.65E-08
2.0E+00	1.32E-07*	2.41E-07	2.10E-07	1.65E-07	1.41E-07	1.07E-07	7.65E-08
4.0E+00	1.32E-07*	2.41E-07	2.10E-07	1.65E-07	1.41E-07	1.07E-07	7.65E-08
7.0E+00	1.32E-07*	2.41E-07	2.10E-07	1.65E-07	1.41E-07	1.07E-07	7.65E-08
1.0E+01	1.32E-07*	2.41E-07	2.10E-07	1.65E-07	1.41E-07	1.07E-07	7.65E-08
2.0E+01	1.32E-07*	2.41E-07	2.10E-07	1.65E-07	1.41E-07	1.07E-07	7.65E-08
4.0E+01	1.32E-07*	2.41E-07	2.10E-07	1.65E-07	1.41E-07	1.07E-07	7.65E-08
7.0E+01	1.32E-07*	2.40E-07	2.10E-07	1.65E-07	1.42E-07	1.07E-07	7.65E-08
1.0E+02	1.32E-07*	2.37E-07	2.10E-07	1.65E-07	1.42E-07	1.07E-07	7.65E-08
2.0E+02	1.32E-07*	2.26E-07	2.10E-07	1.65E-07	1.41E-07	1.08E-07	7.65E-08
4.0E+02	1.32E-07*	2.07E-07*	2.10E-07	1.65E-07	1.42E-07	1.07E-07	7.65E-08
7.0E+02	1.32E-07*	1.91E-07*	2.08E-07	1.65E-07	1.42E-07	1.08E-07	7.64E-08
1.0E+03	1.32E-07*	1.81E-07*	2.05E-07	1.65E-07	1.42E-07	1.08E-07	7.65E-08
2.0E+03	1.33E-07*	1.66E-07*	1.92E-07	1.65E-07	1.42E-07	1.08E-07	7.65E-08
4.0E+03	1.33E-07*	1.54E-07*	1.75E-07*	1.64E-07	1.42E-07	1.08E-07	7.64E-08
7.0E+03	1.34E-07*	1.47E-07*	1.63E-07*	1.59E-07	1.41E-07	1.08E-07	7.64E-08
1.0E+04	1.35E-07*	1.44E-07*	1.55E-07*	1.55E-07	1.40E-07	1.08E-07	7.64E-08
2.0E+04	1.36E-07*	1.39E-07*	1.44E-07*	1.43E-07	1.35E-07	1.07E-07	7.63E-08

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Li Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
25000.	-31.6698	.0118975	.00739492	.00353790	.00119300	1.935930E-04	-1.149263E-04
30000.	-30.8686	-.300540	-.102787	.0265566	.03755728	4.406779E-04	-.0128332
40000.	-30.9410	-.164365	-.104816	-.0390710	.00300641	.0148191	.00812484
70000.	-31.2792	-.0415852	-.0337922	-.0234716	-.0135033	-.00588597	-.00106044
100000.	-31.5530	-.0108539	-.0102585	-.00887341	-.00653162	-.00400398	-.00213914
200000.	-32.0915	3.788109E-04	-1.517981E-04	-3.574076E-04	-3.435617E-04	-2.366833E-04	-1.024141E-04
500000.	-32.7738	-8.388920E-04	-5.417499E-04	-3.069308E-04	-2.075953E-04	-1.598618E-04	-6.145641E-05

See appendix for Chebyshev fit details

$$Li + H^+ \rightarrow Li^+ + H^+ + e$$

Beam – Maxwellian



Cross Sections for the Ionization of Li by He^+ Ions.

$He^+ + Li \rightarrow He^+ + Li^+ + e$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
8.2E+03	1.26E+08	1.67E-15
9.0E+03	1.32E+08	1.66E-15
1.0E+04	1.39E+08	1.65E-15
1.5E+04	1.70E+08	1.53E-15
2.0E+04	1.96E+08	1.38E-15
2.5E+04	2.20E+08	1.17E-15
3.0E+04	2.41E+08	9.31E-16
3.4E+04	2.56E+08	7.58E-16

References: 232, 739

Accuracy: Unknown

Notes: (1) The theoretical cross section of Tiwary et al. (Ref. 739) continues to increase at energies less than 10^4 eV/amu. (2) The experimental data of DuBois (Ref. 232) indicate a maximum in the cross section at 10^4 eV/amu.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 8.2E+03 eV/amu, E_{max} = 3.4E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-68.5495	-.358317	-.141101	-.0391361	-.00786217	.00262636	.00278367

The fit represents the above cross section with an rms deviation of 0.0%. The maximum deviation is 0.1% at 2.5E+04 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + Li \rightarrow He^+ + Li^+ + e$



Cross Sections for Ionization of Li by He^{2+} Ions

$$He^{2+} + Li \rightarrow He^{2+} + Li^{+} + e$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
6.6E+03	1.13E+08	5.62E-16
7.0E+03	1.16E+08	6.85E-16
1.0E+04	1.39E+08	1.42E-15
1.5E+04	1.70E+08	2.77E-15
2.0E+04	1.96E+08	3.80E-15
2.4E+04	2.15E+08	4.06E-15
4.0E+04	2.78E+08	3.02E-15
7.0E+04	3.68E+08	1.86E-15
1.0E+05	4.39E+08	1.35E-15
1.5E+05	5.38E+08	9.43E-16
2.0E+05	6.21E+08	7.36E-16
4.0E+05	8.78E+08	4.01E-16
7.0E+05	1.16E+09	2.51E-16
1.0E+06	1.39E+09	1.84E-16
1.5E+06	1.70E+09	1.28E-16

<u>References:</u> 232, 234, 264, 731, 735, 737

<u>Accuracy:</u> $E > 2x10^4$ eV/amu - 30%
$E < 2x10^4$ eV/amu - Unknown

<u>Notes:</u> (1) The theoretical data of Shipsey, et al. (Ref. 264) have been used to extend the experimental data below $2x10^4$ eV/amu. At higher energies the theoretical data of this paper are approximately 50% below the experimental data. (2) See Shah, et al. (Ref. 234) for double ionization and transfer ionization.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

E_{min} = 6.6E+03 eV/amu, E_{max} = 1.5E+06 eV/amu

A0	A1	A2	A3	A4	A5	A6
-69.7410	-1.20403	-.864397	.480069	-.171618	-.0181494	.0678290

The fit represents the above cross sections with an rms deviation of 3.8%. The maximum deviation is 6.5% at 4.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$He^{2+} + Li \rightarrow He^{2+} + Li^{+} + e$



Ionization Rate Coefficients for

$Li + He^{2+} \rightarrow Li^{+} + He^{2+} + e$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^{2+} Temp. (eV)	Li Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	1.97E-07	7.45E-07	8.39E-07	6.84E-07	5.93E-07	4.57E-07	3.27E-07
2.0E+00	1.97E-07	7.45E-07	8.39E-07	6.84E-07	5.93E-07	4.57E-07	3.27E-07
4.0E+00	1.97E-07	7.44E-07	8.38E-07	6.84E-07	5.93E-07	4.57E-07	3.27E-07
7.0E+00	1.97E-07	7.44E-07	8.38E-07	6.84E-07	5.93E-07	4.57E-07	3.27E-07
1.0E+01	1.97E-07	7.43E-07	8.38E-07	6.83E-07	5.93E-07	4.57E-07	3.27E-07
2.0E+01	1.97E-07	7.42E-07	8.37E-07	6.83E-07	5.93E-07	4.57E-07	3.27E-07
4.0E+01	1.97E-07	7.40E-07	8.37E-07	6.83E-07	5.93E-07	4.57E-07	3.27E-07
7.0E+01	1.98E-07	7.39E-07	8.36E-07	6.83E-07	5.93E-07	4.57E-07	3.27E-07
1.0E+02	1.98E-07	7.38E-07	8.36E-07	6.83E-07	5.93E-07	4.57E-07	3.27E-07
2.0E+02	2.01E-07	7.35E-07	8.34E-07	6.83E-07	5.93E-07	4.57E-07	3.27E-07
4.0E+02	2.06E-07	7.32E-07	8.32E-07	6.83E-07	5.93E-07	4.57E-07	3.27E-07
7.0E+02	2.13E-07	7.28E-07	8.30E-07	6.83E-07	5.93E-07	4.57E-07	3.27E-07
1.0E+03	2.20E-07	7.23E-07	8.28E-07	6.83E-07	5.93E-07	4.57E-07	3.27E-07
2.0E+03	2.44E-07	7.12E-07	8.23E-07	6.82E-07	5.93E-07	4.57E-07	3.27E-07
4.0E+03	2.87E-07	6.95E-07	8.15E-07	6.82E-07	5.93E-07	4.57E-07	3.27E-07
7.0E+03	3.38E-07	6.81E-07	8.03E-07	6.81E-07	5.93E-07	4.56E-07	3.27E-07
1.0E+04	3.77E-07	6.72E-07	7.89E-07	6.79E-07	5.93E-07	4.56E-07	3.27E-07
2.0E+04	4.63E-07	6.57E-07	7.52E-07	6.72E-07	5.92E-07	4.56E-07	3.27E-07

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

Li Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-30.4517	.363146	.218568	.0797980	.00504386	-.0143333	-.0104786
20000.	-28.2941	-.0576610	-.0273357	-.00812937	3.616003E-04	.00260866	.00171889
40000.	-28.0291	-.0386370	-.0241107	-.0131556	-.00664902	-.00299352	-.00102071
70000.	-28.3978	-.00517514	-.00356459	-.00231958	-.00145198	-8.487303E-04	-4.870846E-04
100000.	-28.6764	-3.661189E-04	-4.106187E-04	-3.730240E-04	-2.606518E-04	-1.714793E-04	-7.710836E-05
200000.	-29.1978	-8.814731E-04	-6.105531E-04	-1.205126E-04	-7.285731E-05	1.063444E-04	7.216290E-06
500000.	-29.8691	-1.603658E-04	-9.779805E-05	-3.895455E-05	-5.564509E-06	2.962654E-06	-4.029762E-07

See appendix for Chebyshev fit details.

$$Li + He^{2+} \rightarrow Li^{+} + He^{2+} + e$$



E. Projectile Electron Loss or Stripping Cross Sections

(* denotes rate coefficient data also)

	Page
$H + H \rightarrow H^+ + H + e$	E-2
$H(2s) + H(1s) \rightarrow H^+ + H + e$	E-4
$H + H_2 \rightarrow H^+ + H_2 + e$	E-6
$H(2s) + H_2 \rightarrow H^+ + H_2 + e$	E-8
$H + He \rightarrow H^+ + He + e$	E-10*
$H(2s) + He \rightarrow H^+ + He + e$	E-14
$He + H \rightarrow He^+ + H + e$	E-16
$He + H_2 \rightarrow He^+ + H_2 + e$	E-18
$He + H_2 \rightarrow He^{2+} + H_2 + 2e$	E-18
$He^* + H_2 \rightarrow He$ or He^+	E-20
$He^* + H_2 \rightarrow He^+ + H_2 + e$	E-20
$He^* + H_2 \rightarrow He + H_2$	E-20
$He + He \rightarrow He^+ + He + e$	E-22*
$He + He \rightarrow He^{2+} + He + 2e$	E-22
$He^+ + H \rightarrow He^{2+} + H + e$	E-28
$He^+ + H_2 \rightarrow He^{2+} + H_2 + e$	E-30
$He^+ + He \rightarrow He^{2+} + He + e$	E-32*

Projectile Electron Loss or Stripping Cross Sections for

$H + H \rightarrow H^{+} + H + e$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.2E+03	4.81E+07	2.89E-18
2.0E+03	6.21E+07	1.50E-17
4.0E+03	8.79E+07	3.65E-17
6.0E+03	1.08E+08	5.22E-17
7.0E+03	1.16E+08	5.87E-17
1.0E+04	1.39E+08	7.68E-17
1.5E+04	1.70E+08	1.07E-16
2.0E+04	1.96E+08	1.26E-16
4.0E+04	2.78E+08	9.64E-17
7.0E+04	3.68E+08	7.35E-17
1.0E+05	4.39E+08	5.98E-17
1.5E+05	5.38E+08	4.49E-17
2.0E+05	6.21E+08	3.57E-17
4.0E+05	8.78E+08	1.96E-17
7.0E+05	1.16E+09	1.27E-17
1.0E+06	1.39E+09	9.67E-18
1.5E+06	1.70E+09	7.08E-18
2.0E+06	1.96E+09	5.76E-18
3.5E+06	2.59E+09	3.88E-18

References: 33, 34, 209, 296, 297, 298, 299, 300, 301

Accuracy: 20%

Note: To our knowledge there are no known experimental ionization cross sections for H interacting with H atoms.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.2E+03 eV/amu, E_{max} = 3.5E+06 eV/amu

A0	A1	A2	A3	A4	A5	A6
-77.1794	-.451408	-1.51760	.543975	-.0355154	.00258954	-.0750900

The fit represents the above cross section with an rms deviation of 8.0%. The maximum deviation is 10.3% at 7.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$$H + H \rightarrow H^{+} + H + e$$



Projectile Electron Loss or Stripping Cross Sections for

$H(2s) + H(1s) \rightarrow H^+ + H + e$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
5.0E+03	9.82E+07	2.24E-16
6.0E+03	1.08E+08	2.39E-16
7.0E+03	1.16E+08	2.51E-16
8.0E+03	1.24E+08	2.62E-16
9.0E+03	1.32E+08	2.69E-16
1.0E+04	1.39E+08	2.75E-16
1.5E+04	1.70E+08	2.95E-16
2.0E+04	1.96E+08	2.94E-16
3.0E+04	2.41E+08	2.60E-16
4.0E+04	2.78E+08	2.29E-16
5.0E+04	3.11E+08	2.10E-16
6.0E+04	3.40E+08	1.99E-16
7.0E+04	3.68E+08	1.91E-16
8.0E+04	3.93E+08	1.84E-16
9.0E+04	4.17E+08	1.79E-16
1.0E+05	4.39E+08	1.74E-16

References: 33, 285, 302

Accuracy: Unknown

Note: The experimental data of Hill, et al. (Ref. 33) have been extrapolated from 20 keV/amu to 100 keV/amu using the theoretical data of Abrines and Percival (Ref. 285), and Prasad and Unnikrishnan (Ref. 302). In the energy range of overlap the experimental and theoretical data agree to within 25%.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 5.0E+03 eV/amu, E_{max} = 1.0E+05 eV/amu

A0	A1	A2	A3	A4	A5	A6
-72.0121	-.164818	-.174972	.0460650	.0293401	-.00879104	-.0113468

The fit represents the above cross section with an rms deviation of 0.4%. The maximum deviation is 0.5% at 4.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$$H(2s) + H \rightarrow H^{+} + H + e$$



Projectile Electron Loss or Stripping Cross Sections for

$H + H_2 \rightarrow H^+ + H_2 + e$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
6.0E+01	1.08E+07	1.17E-19
7.0E+01	1.16E+07	1.91E-19
1.0E+02	1.39E+07	5.24E-19
1.5E+02	1.70E+07	1.37E-18
2.0E+02	1.96E+07	2.39E-18
4.0E+02	2.78E+07	8.32E-18
7.0E+02	3.68E+07	2.09E-17
1.0E+03	4.39E+07	3.37E-17
1.5E+03	5.38E+07	5.08E-17
2.0E+03	6.21E+07	6.36E-17
4.0E+03	8.79E+07	8.63E-17
7.0E+03	1.16E+08	9.09E-17
1.0E+04	1.39E+08	9.21E-17
1.5E+04	1.70E+08	1.06E-16
2.0E+04	1.96E+08	1.36E-16
4.0E+04	2.78E+08	1.54E-16
7.3E+04	3.75E+08	1.34E-16
1.0E+05	4.39E+08	1.10E-16
1.5E+05	5.38E+08	8.61E-17
2.0E+05	6.21E+08	7.04E-17
4.0E+05	8.78E+08	4.20E-17
7.0E+05	1.16E+09	2.63E-17
1.0E+06	1.39E+09	1.91E-17
1.5E+06	1.70E+09	1.30E-17
2.0E+06	1.96E+09	9.74E-18
4.0E+06	2.77E+09	4.60E-18
7.0E+06	3.65E+09	2.46E-18
1.0E+07	4.36E+09	1.65E-18
1.5E+07	5.32E+09	1.02E-18
2.0E+07	6.11E+09	7.34E-19

<u>References:</u> 26, 27, 28, 29, 31, 33, 37, 45, 47, 49, 51, 143, 269, 270, 296, 298, 299, 301, 303, 304, 305, 306, 307, 308, 310, 313, 315, 316, 321

<u>Accuracy:</u> 30%

<u>Note:</u> Many Investigators have been involved in these measurements. Consequently there is large scatter in the data, but most of the measurements are within the quoted accuracy of 30%.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

E_{min} = 6.0E+01 eV/amu, E_{max} = 2.0E+07 eV/amu

A0	A1	A2	A3	A4	A5	A6
-79.0892	.346403	-3.04496	.508490	-.0758661	.108006	.00593450

The fit represents the above cross section with an rms deviation of 10.7%. The maximum deviation is 29.8% at 1.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$$H + H_2 \rightarrow H^+ + H_2 + e$$



Projectile Electron Loss or Stripping Cross Sections for

$H(2s) + H_2 \rightarrow H^+ + H_2 + e$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
5.0E+02	3.11E+07	6.14E-17
7.0E+02	3.68E+07	1.09E-16
1.0E+03	4.39E+07	1.90E-16
1.5E+03	5.38E+07	2.89E-16
2.0E+03	6.21E+07	3.33E-16
4.0E+03	8.79E+07	4.05E-16
7.0E+03	1.16E+08	4.36E-16
1.0E+04	1.39E+08	4.50E-16
1.5E+04	1.70E+08	4.63E-16
2.0E+04	1.96E+08	4.76E-16
4.0E+04	2.78E+08	4.00E-16
7.0E+04	3.68E+08	3.00E-16
1.0E+05	4.39E+08	2.48E-16
1.5E+05	5.38E+08	1.95E-16
2.0E+05	6.21E+08	1.59E-16
4.0E+05	8.78E+08	9.56E-17
5.0E+05	9.82E+08	8.13E-17

References: 27, 33, 306, 326, 327, 328, 329, 331, 332

Accuracy: 50%

Note: Due to the uncertainties in measuring the H(2s) metastable fraction, the errors in measuring the cross sections may be large.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 5.0E+02 eV/amu, E_{max} = 5.0E+05 eV/amu

A0	A1	A2	A3	A4	A5	A6
-72.3651	-.0686522	-.939077	.160670	-.0746703	.0714502	-.00310738

The fit represents the above cross section with an rms deviation of 3.2%. The maximum deviation is 7.0% at 7.0E+02 eV/amu.

See appendix for Chebyshev fit details.

$$H(2s) + H_2 \rightarrow H^+ + H_2 + e$$



Projectile Electron Loss or Stripping Cross Sections for

$H + He \rightarrow H^+ + He + e$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
5.0E+01	9.82E+06	2.25E-19
7.0E+01	1.16E+07	5.27E-19
1.0E+02	1.39E+07	1.08E-18
2.0E+02	1.96E+07	3.40E-18
4.0E+02	2.78E+07	9.54E-18
7.0E+02	3.68E+07	2.06E-17
1.0E+03	4.39E+07	3.31E-17
2.0E+03	6.21E+07	6.83E-17
4.0E+03	8.79E+07	1.07E-16
7.0E+03	1.16E+08	1.28E-16
1.0E+04	1.39E+08	1.33E-16
2.0E+04	1.96E+08	1.30E-16
4.0E+04	2.78E+08	1.21E-16
7.0E+04	3.68E+08	1.05E-16
1.0E+05	4.39E+08	8.77E-17
2.0E+05	6.21E+08	5.41E-17
4.0E+05	8.78E+08	3.12E-17
7.0E+05	1.16E+09	1.91E-17
1.0E+06	1.39E+09	1.38E-17
2.0E+06	1.96E+09	7.20E-18
4.0E+06	2.77E+09	3.68E-18
7.0E+06	3.65E+09	2.13E-18
1.0E+07	4.36E+09	1.49E-18
2.0E+07	6.11E+09	7.31E-19

<u>References:</u> 26, 27, 28, 29, 47, 49, 51, 103, 268, 269, 270, 286, 287, 296, 299, 300, 303, 307, 309, 313, 314, 316, 317, 318, 319, 320, 321, 322

<u>Accuracy:</u> 15%

<u>Notes:</u> None

<u>Chebyshev Fitting Parameters for Cross Sections</u>

E_{min} = 5.0E+01 eV/amu, E_{max} = 2.0E+07 eV/amu

A0	A1	A2	A3	A4	A5	A6
-79.0638	.107432	-2.86460	.477451	.0989948	-.0319510	-.0136705

The fit represents the above cross section with an rms deviation of 4.7%. The maximum deviation is 9.9% at 2.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$$H + He \rightarrow H^{+} + He + e$$



Projectile Electron Loss or Stripping Rate Coefficients for

$$H + He \rightarrow H^{+} + He + e$$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

H Temp. (eV)	Equal Temp.	He Temp. (eV) 10.	100.	500.	1000.	5000.	10000.	20000.
1.0E+01	2.29E-13#	2.29E-13#	6.10E-12*	1.09E-10	3.69E-10	3.81E-09	7.66E-09	1.30E-08
2.0E+01	2.53E-12*	1.86E-12*	1.10E-11	1.25E-10	3.94E-10	3.84E-09	7.69E-09	1.30E-08
4.0E+01	1.40E-11	9.69E-12	2.48E-11	1.60E-10	4.47E-10	3.91E-09	7.74E-09	1.30E-08
6.0E+01	3.35E-11	2.28E-11	4.34E-11	1.98E-10	5.01E-10	3.98E-09	7.80E-09	1.31E-08
8.0E+01	6.03E-11	4.08E-11	6.64E-11	2.40E-10	5.58E-10	4.05E-09	7.85E-09	1.31E-08
1.0E+02	9.38E-11	6.33E-11	9.38E-11	2.84E-10	6.16E-10	4.12E-09	7.90E-09	1.31E-08
2.0E+02	3.44E-10	2.35E-10	2.84E-10	5.43E-10	9.27E-10	4.46E-09	8.16E-09	1.33E-08
4.0E+02	1.09E-09	7.76E-10	8.47E-10	1.18E-09	1.62E-09	5.12E-09	8.67E-09	1.36E-08
6.0E+02	1.97E-09	1.45E-09	1.53E-09	1.88E-09	2.34E-09	5.76E-09	9.15E-09	1.40E-08
8.0E+02	2.88E-09	2.16E-09	2.25E-09	2.61E-09	3.06E-09	6.36E-09	9.62E-09	1.43E-08
1.0E+03	3.77E-09	2.89E-09	2.97E-09	3.33E-09	3.77E-09	6.95E-09	1.01E-08	1.46E-08
2.0E+03	7.64E-09	6.22E-09	6.29E-09	6.58E-09	6.95E-09	9.50E-09	1.21E-08	1.60E-08
4.0E+03	1.30E-08	1.11E-08	1.12E-08	1.14E-08	1.16E-08	1.34E-08	1.53E-08	1.83E-08
6.0E+03	1.66E-08	1.46E-08	1.46E-08	1.48E-08	1.49E-08	1.63E-08	1.78E-08	2.03E-08
8.0E+03	1.93E-08	1.72E-08	1.72E-08	1.73E-08	1.75E-08	1.86E-08	1.98E-08	2.19E-08
1.0E+04	2.15E-08	1.93E-08	1.94E-08	1.95E-08	1.96E-08	2.05E-08	2.15E-08	2.33E-08
1.5E+04	2.55E-08	2.33E-08	2.33E-08	2.34E-08	2.35E-08	2.41E-08	2.48E-08	2.61E-08
2.0E+04	2.81E-08	2.61E-08	2.61E-08	2.61E-08	2.62E-08	2.66E-08	2.71E-08	2.81E-08

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+01 eV, E_{max} = 2.0E+04 eV

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
10.	-44.0257	5.73463	-1.21932	.0247503	-.0292957	.0619373	-.0219438
100.	-42.5585	4.48698	-.446257	-.330879	.0963671	.0257805	-.0137058
500.	-40.6373	3.03270	.121804	-.344684	.00515230	.0559208	-.00813670
1000.	-39.6384	2.34972	.269827	-.258791	-.0440918	.0450062	.00550612
5000.	-37.3658	1.00814	.315318	-.0289392	-.0543988	-.00704923	.00822057
10000.	-36.5490	.622335	.247753	.0221122	-.0286150	-.0130702	4.392519E-04
20000.	-35.8634	.357047	.168697	.0373094	-.00809424	-.00961802	-.00307060
Equal Temp.	-43.5995	5.66872	-1.35443	.109372	-.0570829	.0766441	-.0349995

See appendix for Chebyshev fit details.

$$H + He \rightarrow H^{+} + He + e$$

Maxwellian – Maxwellian



Projectile Electron Loss or Stripping Cross Sections for

$H(2s) + He \rightarrow H^+ + He + e$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
5.0E+02	3.11E+07	3.08E-17
7.0E+02	3.68E+07	8.29E-17
1.0E+03	4.39E+07	1.37E-16
1.5E+03	5.38E+07	1.94E-16
2.0E+03	6.21E+07	2.39E-16
4.0E+03	8.79E+07	3.28E-16
7.0E+03	1.16E+08	3.42E-16
1.0E+04	1.39E+08	3.42E-16
1.5E+04	1.70E+08	3.32E-16
2.0E+04	1.96E+08	3.07E-16
4.0E+04	2.78E+08	2.23E-16
7.0E+04	3.68E+08	1.61E-16
1.0E+05	4.39E+08	1.26E-16
1.5E+05	5.38E+08	9.97E-17
2.0E+05	6.21E+08	8.78E-17
4.0E+05	8.78E+08	6.97E-17
5.0E+05	9.82E+08	6.72E-17

References: 29, 324, 326, 327, 329, 332

Accuracy: 50%

Note: Due to the uncertainties in measuring the H(2s) metastable fraction the errors in measuring the cross section may be large. For energies less than 1 keV/amu the data should by used with caution.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 5.0E+02 eV/amu, E_{max} = 5.0E+05 eV/amu

A0	A1	A2	A3	A4	A5	A6
-73.2008	-.0752112	-.927667	.363861	-.0349147	.08236627	-.0565338

The fit represents the above cross section with an rms deviation of 2.6%. The maximum deviation is 4.8% at 1.5E+03 eV/amu.

See appendix for Chebyshev fit details.

$$H(2s) + He \rightarrow H^{+} + He + e$$



Projectile Electron Loss or Stripping Cross Sections for

$He + H \rightarrow He^{+} + H + e$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
5.0E+04	3.11E+08	1.56E-16
6.0E+04	3.40E+08	1.42E-16
7.0E+04	3.68E+08	1.31E-16
8.0E+04	3.93E+08	1.23E-16
9.0E+04	4.17E+08	1.15E-16
1.0E+05	4.39E+08	1.09E-16
1.5E+05	5.38E+08	8.49E-17
2.0E+05	6.21E+08	6.87E-17
3.0E+05	7.61E+08	4.80E-17
4.0E+05	8.78E+08	3.68E-17
5.0E+05	9.82E+08	2.93E-17
6.0E+05	1.08E+09	2.44E-17
7.0E+05	1.16E+09	2.07E-17
8.0E+05	1.24E+09	1.85E-17
9.0E+05	1.32E+09	1.65E-17
1.0E+06	1.39E+09	1.49E-17
1.2E+06	1.52E+09	1.23E-17

References: 301, 398

Accuracy: 15%

Notes: None

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 5.0E+04 eV/amu, E_{max} = 1.2E+06 eV/amu

A0	A1	A2	A3	A4	A5	A6
-75.0809	-1.28466	-.136342	.022920	.0144810	-.00700300	-.00377068

The fit represents the above cross section with an rms deviation of 0.4%. The maximum deviation is 1.3% at 7.0E+05 eV/amu.

See appendix for Chebyshev fit details.

$He + H \rightarrow He^{+} + H + e$



Single and Double Projectile Electron Loss or Stripping Cross Sections for

$$He + H_2 \rightarrow He^{+} + H_2 + e \quad \sigma_{01}$$
$$He + H_2 \rightarrow He^{2+} + H_2 + 2e \quad \sigma_{02}$$

(σ_{01})		(σ_{02})	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
5.0E+02	9.96E-19	5.0E+04	2.59E-18
7.0E+02	2.13E-18	6.0E+04	3.14E-18
1.0E+03	4.68E-18	7.0E+04	3.51E-18
1.5E+03	9.24E-18	8.0E+04	3.78E-18
2.0E+03	1.33E-17	9.0E+04	3.98E-18
4.0E+03	2.73E-17	1.0E+05	4.07E-18
7.0E+03	4.26E-17	1.2E+05	4.10E-18
1.0E+04	5.33E-17	1.5E+05	3.93E-18
1.5E+04	6.75E-17	2.0E+05	3.46E-18
2.0E+04	7.93E-17	3.0E+05	2.44E-18
4.0E+04	1.11E-16	4.0E+05	1.75E-18
7.0E+04	1.27E-16	5.0E+05	1.31E-18
1.0E+05	1.23E-16	6.0E+05	1.09E-18
1.5E+05	1.08E-16	7.0E+05	9.38E-19
2.0E+05	9.38E-17	8.0E+05	8.21E-19
4.0E+05	6.08E-17		
7.0E+05	4.01E-17		
1.0E+06	3.00E-17		

<u>References:</u> (Single Stripping) 94, 97, 99, 102, 103, 143, 198, 301, 316, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409
(Double Stripping) 400, 405

<u>Accuracy:</u> σ_{01} - 15%
σ_{02} - Unknown

<u>Notes:</u> (1) The stripping cross sections are measured in two ways: (a) by the attenuation of the He beam in the H_2 gas cell; (b) by the formation of He^{+}. The attenuation method gives the sum of the single and double stripping cross sections. Since the double-stripping cross section is at least a factor of 30 less than that for single-stripping, the two methods give the same cross sections within the measurement errors. (2) The data for double stripping do not permit an error evaluation.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

σ_{01}	E_{min}	=	5.0E+02	eV/amu,	E_{max}	=	1.0E+06 eV/amu
σ_{02}	E_{min}	=	5.0E+04	eV/amu,	E_{max}	=	8.0E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
σ_{01}	-76.5793	1.58478	-1.39956	.0531448	-.0845579	.0891012	.0231543
σ_{02}	-81.3267	-.669570	-.425168	.0911177	.0266527	.00757867	-.00546826

The fit represents the σ_{01} cross section with an rms deviation of 2.5%. The maximum deviation is 5.4% at 7.0E+02 eV/amu.

The fit represents the σ_{02} cross section with an rms deviation of 0.7%. The maximum deviation is 1.8% at 5.0E+05 eV/amu.

See appendix for Chebyshev fit details.

$$He + H_2 \rightarrow He^{+} + H_2 + e \quad \sigma_{01}$$
$$He + H_2 \rightarrow He^{2+} + H_2 + 2e \quad \sigma_{02}$$



Total Metastable Destruction, Electron Stripping, and De-excitation Cross Sections for

$He^* + H_2 \rightarrow He$ or He^+ (σ_T)
$He^* + H_2 \rightarrow He^+ + H_2 + e$ (σ_{0*1})
$He^* + H_2 \rightarrow He + H_2$ (σ_{0*0})

σ_T		σ_{0*1}		σ_{0*0}	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
1.8E+03	3.72E-16	2.5E+03	2.34E-16	6.3E+03	2.09E-16
2.0E+03	3.94E-16	4.0E+03	3.81E-16	7.0E+03	2.09E-16
4.0E+03	6.16E-16	7.0E+03	5.41E-16	1.0E+04	2.01E-16
7.0E+03	7.49E-16	8.6E+03	5.53E-16	1.5E+04	1.77E-16
8.4E+03	7.63E-16	1.0E+04	5.48E-16	2.0E+04	1.52E-16
1.0E+04	7.54E-16	1.5E+04	4.88E-16	4.0E+04	8.20E-17
1.5E+04	6.69E-16	2.0E+04	4.43E-16	7.0E+04	3.66E-17
2.0E+04	5.98E-16	4.0E+04	3.38E-16	1.0E+05	1.84E-17
4.0E+04	4.32E-16	7.0E+04	2.66E-16	1.2E+05	1.21E-17
7.0E+04	3.13E-16	1.0E+05	2.23E-16		
1.0E+05	2.49E-16	1.3E+05	1.90E-16		
1.2E+05	2.26E-16				

References: 102, 198, 402, 408, 410, 411, 412, 413, 414, 415, 416

Accuracy: 50%

Notes: (1) Tawara (Ref. 102) has listed the excited state lifetimes for He as: (a) 2^1P - 5.6×10^{-10} sec; (b) 2^3P - 1.05×10^{-7} sec; (c) 2^1S - 0.14 sec; (d) 2^3S - 10^5 sec. (2) $\sigma_T = \sigma_{0*1} + \sigma_{0*0}$. (3) Benton, et al. (Ref. 412) found the total destruction cross sections of the triplet metastable state to be $6.0\times10^{-16} cm^2$ at thermal energies. For electron stripping cross sections of other excited states see Pedersen (Ref. 414).

Chebyshev Fitting Parameters for Cross Sections

σ_T	E_{min}	=	1.8E+03	eV/amu,	E_{max}	=	1.2E+05 eV/amu
σ_{0*1}	E_{min}	=	2.5E+03	eV/amu,	E_{max}	=	1.3E+05 eV/amu
σ_{0*0}	E_{min}	=	6.3E+03	eV/amu,	E_{max}	=	1.2E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
σ_T	-70.7481	-.310880	-.437383	.0856563	.0221775	-.0286798	.0139816
σ_{0*1}	-71.3457	-.218099	-.407518	.142300	-.0331160	-.0270893	.0189680
σ_{0*0}	-74.1810	-1.38382	-.434191	-.0390903	-.00312428	-.00143317	-.000970525

The fit represents the σ_T cross section with an rms deviation of 0.6%. The maximum deviation is 0.6% at 1.8E+03 eV/amu.

The fit represents the σ_{0*1} cross section with an rms deviation of 0.5%. The maximum deviation is 0.9% at 1.5E+04 eV/amu.

The fit represents the σ_{0*0} cross section with an rms deviation of 0.0%. The maximum deviation is 0.1% at 7.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$He^* + H_2 \rightarrow He^+ + H_2 + e \; (\sigma_{0'1})$
$He^* + H_2 \rightarrow He \quad + H_2 \qquad (\sigma_{0'0})$



Single and Double Projectile Electron Loss or Stripping Cross Sections for

$$He + He \rightarrow He^{+} + He + e \quad (\sigma_{01})$$
$$He + He \rightarrow He^{2+} + He + 2e \quad (\sigma_{02})$$

σ_{01} Energy (eV/amu)	Cross Section (cm^2)	σ_{02} Energy (eV/amu)	Cross Section (cm^2)
5.0E+01	1.19E-19	5.0E+04	3.92E-18
7.0E+01	2.30E-19	7.0E+04	5.00E-18
1.0E+02	3.40E-19	1.0E+05	5.85E-18
1.5E+02	4.19E-19	1.5E+05	5.04E-18
2.0E+02	4.37E-19	2.0E+05	3.83E-18
4.0E+02	1.37E-18	4.0E+05	1.72E-18
7.0E+02	3.36E-18	7.0E+05	8.71E-19
1.0E+03	5.79E-18	1.0E+06	5.50E-19
1.5E+03	1.07E-17		
2.0E+03	1.64E-17		
4.0E+03	3.77E-17		
7.0E+03	5.37E-17		
1.0E+04	6.74E-17		
1.5E+04	8.16E-17		
2.0E+04	9.21E-17		
4.0E+04	1.08E-16		
7.0E+04	1.00E-16		
1.0E+05	9.31E-17		
1.5E+05	8.37E-17		
2.0E+05	7.40E-17		
4.0E+05	4.89E-17		
5.0E+05	4.18E-17		

<u>References:</u> σ_{01} - 97, 99, 103, 198, 304, 316, 399, 400, 401, 404, 405, 407, 408, 410, 465, 466, 467, 468, 469, 470, 471
σ_{02} - 94, 400, 401, 405, 470, 471

<u>Accuracy:</u> σ_{01} $E > 1x10^3$ eV/amu - 20%
$E < 1x10^3$ eV/amu - 100%
σ_{02} 100%

<u>Note:</u> Most of the data have been taken for a mixed beam of ground state and metastable states of incident He.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

σ_{01} E_{min} = 5.0E+01 eV/amu, E_{max} = 5.0E+05 eV/amu
σ_{02} E_{min} = 5.0E+04 eV/amu, E_{max} = 1.0E+06 eV/amu

	A0	A1	A2	A3	A4	A5	A6
σ_{01}	-78.6347	3.20546	-1.37570	-.359168	.188553	-.0115646	-.0973845
σ_{02}	-81.2673	-1.09626	-.438191	.146687	-.0102958	-.0324691	.0197680

The fit represents the σ_{01} cross section with an rms deviation of 10.2%. The maximum deviation is 32.7% at 2.0E+02 eV/amu.

The fit represents the σ_{02} cross section with an rms deviation of 0.1%. The maximum deviation is 0.2% at 2.0E+05 eV/amu.

See appendix for Chebyshev fit details.

$He + He \rightarrow He^{+} + He + e\ (\sigma_{01})$
$He + He \rightarrow He^{2+} + He + 2e\ (\sigma_{02})$



Single Projectile Electron Loss or Stripping Rate Coefficients for

$He + He \rightarrow He^{+} + He + e$

Maxwellian - Maxwellian Rate Coefficients (cm^3/s)

He Temp. (eV)	Equal Temp.	He Temp. (eV) 10.	100.	500.	1000.	5000.	10000.	20000.
1.0E+01	2.56E-16#	2.56E-16#	9.95E-13*	1.53E-11	5.98E-11	1.13E-09	2.96E-09	6.52E-09
2.0E+01	3.55E-14#	6.78E-15#	1.19E-12*	1.58E-11	6.10E-11	1.14E-09	2.96E-09	6.52E-09
4.0E+01	4.65E-13#	9.72E-14#	1.59E-12*	1.70E-11	6.34E-11	1.14E-09	2.97E-09	6.53E-09
6.0E+01	1.19E-12*	3.17E-13#	2.02E-12*	1.83E-11	6.59E-11	1.15E-09	2.98E-09	6.54E-09
8.0E+01	2.02E-12*	6.30E-13*	2.46E-12*	1.96E-11	6.85E-11	1.16E-09	2.98E-09	6.54E-09
1.0E+02	2.93E-12*	9.95E-13*	2.93E-12*	2.10E-11	7.10E-11	1.17E-09	2.99E-09	6.55E-09
2.0E+02	9.61E-12	3.17E-12*	5.75E-12	2.85E-11	8.46E-11	1.20E-09	3.03E-09	6.58E-09
4.0E+02	3.73E-11	1.01E-11	1.47E-11	4.74E-11	1.15E-10	1.27E-09	3.10E-09	6.65E-09
6.0E+02	8.46E-11	2.17E-11	2.85E-11	7.10E-11	1.50E-10	1.34E-09	3.18E-09	6.72E-09
8.0E+02	1.50E-10	3.83E-11	4.74E-11	9.93E-11	1.88E-10	1.41E-09	3.25E-09	6.78E-09
1.0E+03	2.31E-10	5.98E-11	7.10E-11	1.32E-10	2.31E-10	1.48E-09	3.32E-09	6.85E-09
2.0E+03	7.94E-10	2.33E-10	2.53E-10	3.50E-10	4.87E-10	1.85E-09	3.69E-09	7.18E-09
4.0E+03	2.21E-09	7.97E-10	8.27E-10	9.60E-10	1.13E-09	2.58E-09	4.42E-09	7.83E-09
6.0E+03	3.69E-09	1.49E-09	1.52E-09	1.67E-09	1.85E-09	3.32E-09	5.13E-09	8.47E-09
8.0E+03	5.13E-09	2.22E-09	2.25E-09	2.40E-09	2.58E-09	4.06E-09	5.83E-09	9.08E-09
1.0E+04	6.52E-09	2.96E-09	2.99E-09	3.14E-09	3.32E-09	4.77E-09	6.52E-09	9.69E-09
1.5E+04	9.69E-09	4.78E-09	4.81E-09	4.95E-09	5.13E-09	6.52E-09	8.15E-09	1.11E-08
2.0E+04	1.25E-08	6.52E-09	6.55E-09	6.68E-09	6.85E-09	8.15E-09	9.69E-09	1.25E-08

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+01 eV, E_{max} = 2.0E+04 eV

He Temp. (eV)	A0	A1	A2	A3	A4	A5	A6
10.	-51.7583	8.06294	-1.24558	.480900	-.345667	.00168045	.0989086
100.	-47.7268	4.71689	.727833	-.356780	-.134711	.0403620	.0326103
500.	-45.2262	3.19352	.872249	-.155633	-.153319	.00295441	.0304174
1000.	-43.7465	2.41883	.806189	-.0302189	-.122890	-.0208521	.0205933
5000.	-40.0684	.892003	.435779	.111572	-.00729485	-.0179737	-.00576743
10000.	-38.6577	.504603	.274535	.0948483	.0137433	-.00755328	-.00472901
20000.	-37.3817	.261777	.152417	.0618891	.0157868	5.99102E-04	-.00183208
Equal Temp.	-49.6803	7.93762	-1.58981	.635046	-.579197	.290394	-.0450460

See appendix for Chebyshev fit details.

$He + He \rightarrow He^{+} + He + e$

Maxwellian – Maxwellian



Single Projectile Electron Loss or Stripping Rate Coefficients for

$He + He \rightarrow He^{+} + He + e$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	8.89E-09	2.37E-08	3.00E-08	3.68E-08	4.09E-08	4.60E-08	2.05E-08#
2.0E+00	9.34E-09	1.90E-08	3.00E-08	3.67E-08	4.09E-08	4.59E-08	2.05E-08#
4.0E+00	9.36E-09	1.81E-08	3.00E-08	3.67E-08	4.09E-08	4.60E-08	2.05E-08#
7.0E+00	9.35E-09	1.81E-08	3.00E-08	3.67E-08	4.09E-08	4.60E-08	2.05E-08#
1.0E+01	9.35E-09	1.81E-08	3.00E-08	3.67E-08	4.09E-08	4.59E-08	2.05E-08#
2.0E+01	9.35E-09	1.81E-08	2.99E-08	3.67E-08	4.09E-08	4.59E-08	2.05E-08#
4.0E+01	9.35E-09	1.81E-08	2.99E-08	3.67E-08	4.09E-08	4.59E-08	2.05E-08#
7.0E+01	9.35E-09	1.81E-08	2.99E-08	3.67E-08	4.09E-08	4.59E-08	2.05E-08#
1.0E+02	9.36E-09	1.80E-08	2.99E-08	3.67E-08	4.09E-08	4.59E-08	2.05E-08#
2.0E+02	9.38E-09	1.80E-08	2.98E-08	3.67E-08	4.09E-08	4.59E-08	2.05E-08#
4.0E+02	9.43E-09	1.81E-08	2.97E-08	3.67E-08	4.09E-08	4.59E-08	2.04E-08#
7.0E+02	9.51E-09	1.81E-08	2.97E-08	3.67E-08	4.08E-08	4.58E-08	2.04E-08#
1.0E+03	9.60E-09	1.81E-08	2.96E-08	3.67E-08	4.08E-08	4.58E-08	2.04E-08#
2.0E+03	9.91E-09	1.83E-08	2.95E-08	3.67E-08	4.08E-08	4.57E-08	2.03E-08#
4.0E+03	1.05E-08	1.86E-08	2.95E-08	3.67E-08	4.08E-08	4.57E-08	2.02E-08#
7.0E+03	1.14E-08	1.91E-08	2.95E-08	3.68E-08	4.08E-08	4.55E-08	2.01E-08#
1.0E+04	1.22E-08	1.96E-08	2.95E-08	3.68E-08	4.08E-08	4.55E-08	2.00E-08#
2.0E+04	1.46E-08	2.11E-08	2.98E-08	3.69E-08	4.08E-08	4.52E-08	1.98E-08#

Accuracy: * - Possible Error Greater Than 10%
- Possible Error Greater Than 100%

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-36.8050	.170883	.101708	.0672106	.0156617	.0105796	-.00356384
20000.	-35.5374	-.0131879	.0937563	-.0235657	.0434438	-.0195503	.0142526
40000.	-34.6583	-.00778924	6.32056E-04	.00349830	.00265169	.00125371	4.62226E-04
70000.	-34.2378	9.38356E-04	.00134041	9.62664E-04	4.31072E-04	6.92929E-05	-2.31386E-05
100000.	-34.0260	-6.94332E-04	3.47489E-05	2.89415E-04	6.86603E-05	-2.08349E-05	-8.09043E-05
200000.	-33.7994	-.00708522	-.00341665	-.00168503	-3.16686E-04	-3.04255E-04	1.04496E-04
500000.	-35.4225	-.0144815	-.00746534	-.00291476	-9.54856E-04	-2.93045E-04	-9.76578E-05

See appendix for Chebyshev fit details.

$He + He \rightarrow He^{+} + He + e$



Electron Projectile Loss or Stripping Cross Sections for

$He^{+} + H \rightarrow He^{2+} + H + e$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.2E+04	1.52E+08	1.66E-19
1.5E+04	1.70E+08	4.71E-19
2.0E+04	1.96E+08	1.15E-18
4.0E+04	2.78E+08	7.03E-18
7.0E+04	3.68E+08	1.11E-17
1.0E+05	4.39E+08	1.04E-17
1.5E+05	5.38E+08	9.34E-18
2.0E+05	6.21E+08	8.17E-18
4.0E+05	8.78E+08	5.26E-18
7.0E+05	1.16E+09	3.63E-18
9.0E+05	1.32E+09	3.04E-18

References: 301, 476

Accuracy: 20%

Notes: None

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.2E+04 eV/amu, E_{max} = 9.0E+05 eV/amu

A0	A1	A2	A3	A4	A5	A6
-80.7839	1.04755	-1.36971	.441034	-.0292764	-.0527228	.0184479

The fit represents the above cross section with an rms deviation of 4.3%
The maximum devation is 8.7% at 2.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$He^{+} + H \rightarrow He^{2+} + H + e$



Projectile Electron Loss or Stripping Cross Sections for

$He^{+} + H_2 \rightarrow He^{2+} + H_2 + e$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
5.0E+03	9.82E+07	1.29E-19
7.0E+03	1.16E+08	1.99E-19
1.0E+04	1.39E+08	3.25E-19
1.5E+04	1.70E+08	6.18E-19
2.0E+04	1.96E+08	1.06E-18
4.0E+04	2.78E+08	5.77E-18
7.0E+04	3.68E+08	1.45E-17
1.0E+05	4.39E+08	2.02E-17
1.3E+05	5.01E+08	2.16E-17
1.5E+05	5.38E+08	2.16E-17
2.0E+05	6.21E+08	1.85E-17
4.0E+05	8.78E+08	1.13E-17
7.0E+05	1.16E+09	7.81E-18
9.5E+05	1.35E+09	6.29E-18

<u>References:</u> 94, 96, 97, 100, 101, 296, 301, 464, 476, 478

<u>Accuracy:</u> $E < 3x10^4$ eV/amu - 40%
$E > 3x10^4$ eV/amu - 20%

<u>Notes:</u> None

<u>Chebyshev Fitting Parameters for Cross Sections</u>

E_{min} = 5.0E+03 eV/amu, E_{max} = 9.5E+05 eV/amu

A0	A1	A2	A3	A4	A5	A6
-80.9631	2.17474	-1.29222	-.291348	.317779	.0620566	-.0993825

The fit represents the above cross section with an rms deviation of 2.3%. The maximum deviation is 5.1% at 2.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$$He^{+} + H_2 \rightarrow He^{2+} + H_2 + e$$



Cross Sections for Projectile Electron Loss or Stripping for He^+ Passing Through Helium

$$He^+ + He \rightarrow He^{2+} + He + e$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
3.5E+03	8.22E+07	2.63E-19
4.0E+03	8.79E+07	3.68E-19
7.0E+03	1.16E+08	1.02E-18
1.0E+04	1.39E+08	1.64E-18
1.5E+04	1.70E+08	2.74E-18
2.0E+04	1.96E+08	3.87E-18
4.0E+04	2.78E+08	8.89E-18
7.0E+04	3.68E+08	1.67E-17
1.0E+05	4.39E+08	2.14E-17
1.5E+05	5.38E+08	2.35E-17
1.7E+05	5.73E+08	2.36E-17
2.0E+05	6.21E+08	2.34E-17
4.0E+05	8.78E+08	1.69E-17
7.0E+05	1.16E+09	1.07E-17
1.0E+06	1.39E+09	7.73E-18
1.5E+06	1.70E+09	5.21E-18
2.0E+06	1.96E+09	3.91E-18
4.0E+06	2.77E+09	1.98E-18

<u>References:</u> 94, 96, 97, 101, 296, 375, 471, 477, 480, 481, 483, 485, 486, 487, 488, 489, 490, 491, 492, 493

<u>Accuracy:</u> 20%

<u>Notes:</u> None

<u>Chebyshev Fitting Parameters for Cross Sections</u>

E_{min} = 3.5E+03 eV/amu, E_{max} = 4.0E+06 eV/amu

A0	A1	A2	A3	A4	A5	A6
-80.4254	.937124	-1.62973	.00297670	.154841	.0696031	-.0892584

The fit represents the above cross section with an rms deviation of 1.6%. The maximum deviation is 3.5% at 4.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$He^{+} + He \rightarrow He^{2+} + He + e$



Projectile Electron Loss or Stripping Rate Coefficients for

$He + He^{+} \rightarrow He^{2+} + He + e$

Beam - Maxwellian Rate Coefficients (cm^3/s)

He^{+} Temp. (eV)	He Energy (eV/amu) 10000.	20000.	40000.	70000.	100000.	200000.	500000.
1.0E+00	2.93E-10	1.20E-09	8.75E-10	4.39E-09	1.06E-08	1.45E-08	1.38E-08
2.0E+00	2.38E-10	8.92E-10	1.82E-09	5.65E-09	9.90E-09	1.45E-08	1.38E-08
4.0E+00	2.28E-10	7.71E-10	2.38E-09	5.99E-09	9.37E-09	1.45E-08	1.38E-08
7.0E+00	2.28E-10	7.60E-10	2.47E-09	6.09E-09	9.37E-09	1.45E-08	1.38E-08
1.0E+01	2.28E-10	7.61E-10	2.47E-09	6.12E-09	9.38E-09	1.45E-08	1.38E-08
2.0E+01	2.28E-10	7.61E-10	2.47E-09	6.13E-09	9.38E-09	1.45E-08	1.38E-08
4.0E+01	2.28E-10	7.62E-10	2.47E-09	6.12E-09	9.38E-09	1.45E-08	1.38E-08
7.0E+01	2.29E-10	7.63E-10	2.47E-09	6.12E-09	9.37E-09	1.45E-08	1.38E-08
1.0E+02	2.30E-10	7.64E-10	2.47E-09	6.12E-09	9.36E-09	1.45E-08	1.38E-08
2.0E+02	2.32E-10	7.67E-10	2.48E-09	6.11E-09	9.35E-09	1.45E-08	1.38E-08
4.0E+02	2.36E-10	7.74E-10	2.49E-09	6.10E-09	9.34E-09	1.45E-08	1.38E-08
7.0E+02	2.43E-10	7.85E-10	2.50E-09	6.10E-09	9.32E-09	1.44E-08	1.38E-08
1.0E+03	2.50E-10	7.96E-10	2.52E-09	6.11E-09	9.31E-09	1.44E-08	1.38E-08
2.0E+03	2.72E-10	8.32E-10	2.57E-09	6.13E-09	9.29E-09	1.44E-08	1.38E-08
4.0E+03	3.19E-10	9.05E-10	2.68E-09	6.19E-09	9.29E-09	1.43E-08	1.38E-08
7.0E+03	3.92E-10	1.02E-09	2.83E-09	6.28E-09	9.30E-09	1.42E-08	1.38E-08
1.0E+04	4.70E-10	1.13E-09	2.99E-09	6.38E-09	9.32E-09	1.42E-08	1.38E-08
2.0E+04	7.54E-10	1.52E-09	3.46E-09	6.67E-09	9.41E-09	1.40E-08	1.38E-08

Chebyshev Fitting Parameters for Rate Coefficients

E_{min} = 1.0E+00 eV, E_{max} = 2.0E+04 eV

He Energy (eV/amu)	A0	A1	A2	A3	A4	A5	A6
10000.	-43.8442	.378813	.344057	.105569	.0867559	-.00821300	.0129056
20000.	-41.5896	.123902	.270872	.0115805	.0887436	-.0161519	.0126729
40000.	-39.7835	.385155	-.145042	.223297	-.0998296	.0753339	-.0326980
70000.	-37.8827	.110612	-.0493705	.0701363	-.0309008	.0252250	-.0129206
100000.	-36.9382	-.0374325	.0312796	-.0188116	.0166138	-.00628486	.00207752
200000.	-36.1132	-.0149848	-.00760145	-.00280029	-8.61479E-04	-1.69493E-04	-3.20142E-05
500000.	-36.1937	-5.70799E-04	-4.55366E-04	-3.44110E-04	-2.36175E-04	-1.47246E-04	-9.193766E-05

See appendix for Chebyshev fit details.

$$He + He^{+} \rightarrow He^{2+} + He + e$$

Beam – Maxwellian



F. Electron Detachment Cross Sections

	Page
$H^- + H \rightarrow H + H + e$	F-2
$\rightarrow (H + H + e)$ or $(H + H^-)$	F-2
$\rightarrow H^+ + H + 2e$	F-4
$H^- + H^+ \rightarrow H + H^+ + 2e$	F-6
$H^- + H_2 \rightarrow H + H_2 + e$	F-8
$\rightarrow H^+ + H_2 + 2e$	F-10
$H^- + He \rightarrow H + He + e$	F-12
$\rightarrow H^+ + He + 2e$	F-14
$He^- + H \rightarrow He + H + e$	F-16
$He^- + H_2 \rightarrow He + H_2 + e$	F-18
$\rightarrow He^+ + H_2 + 2e$	F-18
$He^- + He \rightarrow He + He + e$	F-20
$\rightarrow He^+ + He + 2e$	F-20
$Li^- + H_2 \rightarrow Li + H_2 + e$	F-22
$\rightarrow Li^+ + H_2 + 2e$	F-22
$Li^- + He \rightarrow Li + He + e$	F-24
$\rightarrow Li^+ + He + 2e$	F-24

Cross Section for Single Electron Detachment and Total H^0 Formation from H^- on H

$H^- + H \rightarrow H + H + e$
$H^- + H \rightarrow (H + H + e)$ or $(H + H^-)$ [Total H Prod.]

H + H + e		Total H Prod.	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
1.1E+02	8.65E-16	6.0E+01	6.64E-15
2.0E+02	1.28E-15	7.0E+01	6.75E-15
4.0E+02	1.44E-15	1.0E+02	6.79E-15
7.0E+02	1.39E-15	1.5E+02	6.48E-15
1.0E+03	1.32E-15	2.0E+02	6.11E-15
1.5E+03	1.19E-15	4.0E+02	4.92E-15
2.0E+03	1.11E-15	7.0E+02	3.83E-15
4.0E+03	1.01E-15	1.0E+03	3.13E-15
7.0E+03	9.75E-16	1.5E+03	2.45E-15
1.0E+04	8.72E-16	2.0E+03	2.03E-15
1.5E+04	7.18E-16	4.0E+03	1.26E-15
2.0E+04	6.26E-16	7.0E+03	1.01E-15
4.0E+04	4.25E-16	1.0E+04	8.74E-16
7.0E+04	2.90E-16	1.5E+04	7.18E-16
1.0E+05	2.21E-16	2.0E+04	6.26E-16
1.5E+05	1.61E-16	4.0E+04	4.25E-16
2.0E+05	1.27E-16	7.0E+04	2.90E-16
4.0E+05	7.04E-17	1.0E+05	2.21E-16
7.0E+05	4.24E-17	1.5E+05	1.61E-16
1.0E+06	2.99E-17	2.0E+05	1.27E-16
1.5E+06	2.09E-17	4.0E+05	7.04E-17
2.0E+06	1.59E-17	7.0E+05	4.24E-17
4.0E+06	8.23E-18	1.0E+06	2.99E-17
		1.5E+06	2.09E-17
		2.0E+06	1.59E-17
		4.0E+06	8.23E-18

<u>References:</u> 162, 281, 301, 395, 524, 525, 554, 555

<u>Accuracy:</u> 30%

<u>Note:</u> Data for total H^0 formation represent the sum of the cross sections for collisional detachment and charge exchange. The charge exchange contribution is negligible at energies above 10 keV/amu.

Chebyshev Fitting Parameters for Cross Sections

H + H + e	E_{min} = 1.1E+02 eV/amu,	E_{max} = 4.0E+06 eV/amu
Total H Prod.	E_{min} = 6.0E+01 eV/amu,	E_{max} = 4.0E+06 eV/amu

	A0	A1	A2	A3	A4	A5	A6
H + H + e	-72.0699	-2.47924	-.936325	.0886530	.0175392	.0545278	-.0645320
Total H Prod.	-70.7835	-3.41032	-.545568	.00370195	-.0606176	.0839778	.00626877

The fit represents H + H + e cross section with an rms deviation of 2.7%. The maximum deviation is 5.4% at 2.0E+03 eV/amu.

The fit repreents Total H Prod. cross section with an rms deviation of 3.7%. The maximum deviation is 12.4% at 4.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$H^- + H \rightarrow (H + H + e)$ or $(H + H^-)$
$H^- + H \rightarrow H + H + e$



Cross Sections for Double Electron Detachment of H^- in H

$$H^- + H \rightarrow H^+ + H + 2e$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.8E+03	5.89E+07	3.11E-18
2.0E+03	6.21E+07	3.57E-18
4.0E+03	8.79E+07	9.35E-18
7.0E+03	1.16E+08	1.50E-17
1.0E+04	1.39E+08	1.77E-17
1.5E+04	1.70E+08	1.87E-17
2.0E+04	1.96E+08	1.85E-17
4.0E+04	2.78E+08	1.43E-17
7.0E+04	3.68E+08	9.40E-18
1.0E+05	4.39E+08	6.79E-18
1.5E+05	5.38E+08	4.61E-18
2.0E+05	6.21E+08	3.51E-18
3.0E+05	7.61E+08	2.50E-18

References: 524, 525

Accuracy: 20%

Note: There is only one set of experimental measurements.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.8E+03 eV/amu, E_{max} = 3.0E+05 eV/amu

A0	A1	A2	A3	A4	A5	A6
-79.0633	-.259790	-.938476	.154218	.0364114	.00411484	.00724986

The fit represents the above cross section with an rms deviation of 0.9%. The maximum deviation is 1.8% at 2.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$$H^- + H \rightarrow H^+ + H + 2e$$



Cross Sections for Single Electron Detachment of H^- Colliding with H^+

$$H^- + H^+ \rightarrow H + H^+ + e$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.8E+03	7.35E+07	3.99E-16
3.0E+03	7.61E+07	6.57E-16
4.0E+03	8.79E+07	1.55E-15
5.0E+03	9.82E+07	2.24E-15
6.0E+03	1.08E+08	2.73E-15
7.0E+03	1.16E+08	3.11E-15
8.0E+03	1.24E+08	3.40E-15
9.0E+03	1.32E+08	3.64E-15
1.0E+04	1.39E+08	3.82E-15
1.5E+04	1.70E+08	4.23E-15
2.0E+04	1.96E+08	4.18E-15
3.0E+04	2.41E+08	3.88E-15
4.0E+04	2.78E+08	3.51E-15
5.0E+04	3.11E+08	3.19E-15
6.0E+04	3.40E+08	2.92E-15
7.0E+04	3.68E+08	2.66E-15

References: 69, 569, 572, 573

Accuracy: 40%

Note: Accuracy taken from the only experimental data available. Theoretical calculations presented by Fussen and Claeys (Ref. 572), and Ermolaev (Ref. 573) agree very well with experimental data.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 2.8E+03 eV/amu, E_{max} = 7.0E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-67.3879	.650676	-.658353	.211309	-.0972717	.0547639	-.0359904

The fit represents the above cross section with an rms deviation of 2.6%. The maximum deviation is 5.7% at 2.8E+03 eV/amu.

See appendix for Chebyshev fit details.

$$H^- + H^+ \rightarrow H + H^+ + e$$



Cross Sections for Single Electron Detachment of H^- in H_2

$$H^- + H_2 \rightarrow H + H_2 + e$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.3E+00	2.11E+06	8.91E-17
4.0E+00	2.78E+06	1.92E-16
7.0E+00	3.68E+06	3.18E-16
1.0E+01	4.39E+06	3.68E-16
1.5E+01	5.38E+06	4.14E-16
2.0E+01	6.21E+06	4.28E-16
4.0E+01	8.79E+06	4.15E-16
7.0E+01	1.16E+07	4.36E-16
1.0E+02	1.39E+07	4.80E-16
1.5E+02	1.70E+07	5.60E-16
2.0E+02	1.96E+07	6.29E-16
4.0E+02	2.78E+07	8.41E-16
7.0E+02	3.68E+07	1.00E-15
1.0E+03	4.39E+07	1.06E-15
1.4E+03	5.20E+07	1.10E-15
2.0E+03	6.21E+07	1.13E-15
4.0E+03	8.79E+07	1.14E-15
7.0E+03	1.16E+08	1.09E-15
1.0E+04	1.39E+08	1.02E-15
1.5E+04	1.70E+08	9.24E-16
2.0E+04	1.96E+08	8.36E-16
4.0E+04	2.78E+08	6.33E-16
7.0E+04	3.68E+08	4.82E-16
1.0E+05	4.39E+08	3.95E-16
1.5E+05	5.38E+08	3.06E-16
2.0E+05	6.21E+08	2.51E-16
4.0E+05	8.78E+08	1.43E-16
7.0E+05	1.16E+09	9.02E-17
1.0E+06	1.39E+09	6.43E-17
1.5E+06	1.70E+09	4.41E-17
2.0E+06	1.96E+09	3.29E-17
4.0E+06	2.77E+09	1.55E-17
7.0E+06	3.65E+09	8.13E-18
1.0E+07	4.36E+09	5.16E-18
1.5E+07	5.32E+09	3.17E-18
1.7E+07	5.65E+09	2.72E-18

<u>References:</u> 27, 32, 82, 269, 299, 301, 303, 307, 321, 395, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 526, 527, 541, 542, 544

<u>Accuracy:</u> 25%

<u>Note:</u> For total detachment cross sections ($\sigma_{-10} + \sigma_{-11}$) see: Muschlitz, Jr. et al., (Ref. 514, 516); Stedeford and Hasted (Ref. 82); Hasted and Smith (Ref. 515); Risley and Geballe (Ref. 520); Risley (Ref. 544); Jorgensen, Jr. et al. (Ref. 32); Lichtenbert et al. (Ref. 523)

<u>Chebyshev Fitting Parameters for Cross Sections</u>

E_{min} = 2.3E+00 eV/amu, E_{max} = 1.7E+07 eV/amu

A0	A1	A2	A3	A4	A5	A6
-73.1506	-1.75695	-2.00168	-.190280	.0171353	.127083	-.152313

The fit represents the above cross section with an rms deviation of 7.2%. The maximum deviation is 16.6% at 7.0E+01 eV/amu.

See appendix for Chebyshev fit details.

$$H^- + H_2 \rightarrow H + H_2 + e$$



Cross Sections for Double Electron Detachment of H^- in H_2

$$H^- + H_2 \rightarrow H^+ + H_2 + 2e$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.0E+03	4.39E+07	1.48E-17
2.0E+03	6.21E+07	2.31E-17
4.0E+03	8.79E+07	3.24E-17
7.0E+03	1.16E+08	3.79E-17
1.0E+04	1.39E+08	4.03E-17
1.5E+04	1.70E+08	4.16E-17
2.0E+04	1.96E+08	4.11E-17
4.0E+04	2.78E+08	3.97E-17
7.0E+04	3.68E+08	3.36E-17
1.0E+05	4.39E+08	2.84E-17
1.5E+05	5.38E+08	2.20E-17
2.0E+05	6.21E+08	1.76E-17
4.0E+05	8.78E+08	8.71E-18
7.0E+05	1.16E+09	4.98E-18
1.0E+06	1.39E+09	3.42E-18
1.5E+06	1.70E+09	2.23E-18
2.0E+06	1.96E+09	1.67E-18
4.0E+06	2.77E+09	8.12E-19
7.0E+06	3.65E+09	4.52E-19
1.0E+07	4.36E+09	3.13E-19

<u>References:</u> 269, 299, 303, 307, 321, 518, 521, 522, 524, 525, 526, 556, 563, 567

<u>Accuracy:</u> 30%

<u>Note:</u> There are no experimental or theoretical data for energies below 1 keV/amu.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.0E+03 eV/amu, E_{max} = 1.0E+07 eV/amu

A0	A1	A2	A3	A4	A5	A6
-79.0159	-2.10252	-1.24073	.174799	.106249	-.000434273	-0.0465674

The fit represents the above cross section with an rms deviation of 2.0%. The maximum deviation is 4.3% at 4.0E+05 eV/amu.

See appendix for Chebyshev fit details.

$$H^- + H_2 \rightarrow H^+ + H_2 + 2e$$



Cross Sections for Single Electron Detachment of H^- in He

$H^- + He \rightarrow H + He + e$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.0E+00	1.39E+06	6.13E-17
2.0E+00	1.96E+06	1.25E-16
4.0E+00	2.78E+06	2.33E-16
7.0E+00	3.68E+06	2.94E-16
1.0E+01	4.39E+06	3.03E-16
1.5E+01	5.38E+06	3.14E-16
2.0E+01	6.21E+06	3.22E-16
4.0E+01	8.79E+06	3.42E-16
7.0E+01	1.16E+07	3.70E-16
1.0E+02	1.39E+07	3.91E-16
1.5E+02	1.70E+07	4.21E-16
2.0E+02	1.96E+07	4.48E-16
4.0E+02	2.78E+07	5.10E-16
7.0E+02	3.68E+07	5.60E-16
1.0E+03	4.39E+07	6.02E-16
1.5E+03	5.38E+07	6.38E-16
2.0E+03	6.21E+07	6.53E-16
3.4E+03	8.10E+07	6.78E-16
4.0E+03	8.79E+07	6.73E-16
7.0E+03	1.16E+08	6.34E-16
1.0E+04	1.39E+08	6.08E-16
1.5E+04	1.70E+08	5.50E-16
2.0E+04	1.96E+08	5.01E-16
4.0E+04	2.78E+08	3.83E-16
7.0E+04	3.68E+08	2.80E-16
1.0E+05	4.39E+08	2.24E-16
1.5E+05	5.38E+08	1.72E-16
2.0E+05	6.21E+08	1.42E-16
4.0E+05	8.78E+08	8.33E-17
7.0E+05	1.16E+09	5.27E-17
1.0E+06	1.39E+09	4.00E-17
1.5E+06	1.70E+09	2.92E-17
2.0E+06	1.96E+09	2.27E-17
4.0E+06	2.77E+09	1.23E-17
7.0E+06	3.65E+09	6.98E-18
1.0E+07	4.36E+09	4.63E-18
1.5E+07	5.32E+09	2.82E-18

<u>References:</u> 27, 82, 269, 299, 300, 303, 307, 318, 321, 518, 519, 520, 521, 522, 523, 525, 527, 529, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 584

<u>Accuracy:</u> 25%

<u>Notes:</u> (1) For total detachment cross sections ($\sigma_{T0} + 2\sigma_{T1}$) see: Stedeford and Hasted (Ref. 82); Lichtenberg et al. (Ref. 523); Risley (Ref. 544); Hasted (Ref. 553); Bailey et al. (Ref. 543); Risley and Geballe (Ref. 520); Champion et al. (Ref. 545); Andersen et al. (Ref. 551); Anderson et al. (ref. 547); Huq et al. (Ref. 548); Risley and Olson (Ref. 549); Risley (Ref. 318).

(2) For $D^- + He \rightarrow D$ see: Risley (Ref. 544); Berkner et al. (Ref. 307); Champion et al. (Ref. 545); Anderson et al. (Ref. 547); Huq et al. (Ref. 548).

<u>Chebyshev Fitting Parameters for Cross Sections</u>

E_{min} = 1.0E+00 eV/amu, E_{max} = 1.5E+07 eV/amu

A0	A1	A2	A3	A4	A5	A6
-73.7795	-1.54839	-1.84526	-.140437	-.0282087	.183809	-.132562

The fit represents the above cross section with an rms deviation of 3.0%. The maximum deviation is 14.4% at 2.0E+00 eV/amu.

See appendix for Chebyshev fit details.

$H^- + He \rightarrow H + He + e$



Cross Sections for Double Electron Detachment of H^- on He

$$H^- + He \rightarrow H^+ + He + 2e$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
4.0E+02	2.78E+07	2.03E-18
7.0E+02	3.68E+07	7.10E-18
1.0E+03	4.39E+07	1.30E-17
1.5E+03	5.38E+07	2.32E-17
2.0E+03	6.21E+07	3.16E-17
4.0E+03	8.79E+07	4.96E-17
7.0E+03	1.16E+08	5.51E-17
1.0E+04	1.39E+08	5.46E-17
1.5E+04	1.70E+08	5.12E-17
2.0E+04	1.96E+08	4.66E-17
4.0E+04	2.78E+08	3.25E-17
7.0E+04	3.68E+08	2.01E-17
1.0E+05	4.39E+08	1.37E-17
1.5E+05	5.38E+08	8.92E-18
2.0E+05	6.21E+08	6.54E-18
4.0E+05	8.78E+08	3.16E-18
7.0E+05	1.16E+09	1.75E-18
1.0E+06	1.39E+09	1.19E-18
1.5E+06	1.70E+09	7.74E-19
2.0E+06	1.96E+09	5.59E-19
4.0E+06	2.77E+09	2.68E-19
7.0E+06	3.65E+09	1.48E-19
1.0E+07	4.36E+09	1.01E-19
1.5E+07	5.32E+09	6.53E-20
2.0E+07	6.11E+09	4.89E-20
2.3E+07	6.54E+09	4.16E-20

<u>References:</u> 268, 269, 299, 303, 307, 318, 321, 518, 521, 522, 523, 525, 552, 556, 561, 568, 584

<u>Accuracy:</u> 30%

<u>Note:</u> There are no experimental data for energies below 400 eV/amu. Theoretical calculations extend down to 100 eV/amu.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 4.0E+02 eV/amu, E_{max} = 2.3E+07 eV/amu

A0	A1	A2	A3	A4	A5	A6
-81.1984	-2.71121	-1.94690	.784427	-.131737	.0264605	-.00697915

The fit represents the above cross section with an rms deviation of 2.6%. The maximum deviation is 5.1% at 2.0E+05 eV/amu.

See appendix for Chebyshev fit details.

$H^- + He \rightarrow H^+ + He + 2e$



Cross Sections for Single Electron Detachment of He^- in H

$He^- + H \rightarrow He + H + e$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
3.7E+05	8.45E+08	6.39E-16
4.0E+05	8.78E+08	6.16E-16
7.0E+05	1.16E+09	4.29E-16
1.0E+06	1.39E+09	3.29E-16
1.5E+06	1.70E+09	2.39E-16
2.0E+06	1.96E+09	1.88E-16
4.0E+06	2.77E+09	1.04E-16
7.0E+06	3.65E+09	6.34E-17
9.0E+06	4.14E+09	4.99E-17

References: 301, 583

Accuracy: 40%

Note: There is only one set of experimental measurements.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 3.7E+05 eV/amu, E_{max} = 9.0E+06 eV/amu

A0	A1	A2	A3	A4	A5	A6
-72.3804	-1.28777	-.0622952	.0122474	-.00652473	.000145452	-.00224648

The fit represents the above cross section with an rms deviation of 0.1%. The maximum deviation is 0.2% at 2.0E+06 eV/amu.

See appendix for Chebyshev fit details.

$He^- + H \rightarrow He + H + e$



Cross Sections for Single and Double Electron Detachment of He^- in H_2

$$He^- + H_2 \rightarrow He + H_2 + e$$
$$\rightarrow He^+ + H_2 + 2e$$

$He + H_2 + e$		$He^+ + H_2 + 2e$	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
1.3E+02	2.15E-15	2.6E+04	3.86E-17
2.0E+02	2.17E-15	3.0E+04	3.89E-17
4.0E+02	2.14E-15	4.0E+04	3.92E-17
7.0E+02	2.09E-15	5.0E+04	3.94E-17
1.0E+03	2.06E-15	6.0E+04	3.91E-17
1.5E+03	2.02E-15	7.0E+04	3.91E-17
2.0E+03	1.97E-15	8.0E+04	3.86E-17
4.0E+03	1.84E-15	9.0E+04	3.82E-17
7.0E+03	1.69E-15	1.0E+05	3.73E-17
1.0E+04	1.56E-15	1.5E+05	3.31E-17
1.5E+04	1.41E-15	2.0E+05	2.96E-17
2.0E+04	1.30E-15	3.0E+05	2.42E-17
4.0E+04	9.76E-16	4.0E+05	2.01E-17
7.0E+04	7.15E-16	5.0E+05	1.72E-17
1.0E+05	5.67E-16	6.0E+05	1.50E-17
1.5E+05	4.27E-16	7.0E+05	1.33E-17
2.0E+05	3.44E-16	8.0E+05	1.19E-17
4.0E+05	1.97E-16		
7.0E+05	1.24E-16		
8.0E+05	1.11E-16		

References: 32, 301, 519, 521, 522, 525, 577, 580, 582, 583

Accuracy: 30%

Notes: (1) The total electron detachment cross section (single and double) has been measured by Jorgensen and Kuyatt (Ref. 32).

(2) For single detachment see: Ryding et al. (Ref. 577); Simpson and Gilbody (Ref. 519); Heinemeier et al. (Ref. 581); Heinemeier et al. (Ref. 521); Coggiola (Ref. 582); Hvelplund and Andersen (Ref. 301).

(3) For double detachment see: Ryding et al. (Ref. 577); Heinemeier et al. (Ref. 581); Heinemeier et al. (Ref. 521).

Chebyshev Fitting Parameters for Cross Sections

Single det. E_{min} = 1.3E+02 eV/amu, E_{max} = 8.0E+05 eV/amu
Double det. E_{min} = 2.6E+04 eV/amu, E_{max} = 8.0E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
Single det.	-69.4085	-1.41471	-.581942	-.0905120	.0321529	.0250996	.000363782
Double det.	-76.3306	-.578433	-.226903	-.00774557	.0112528	-.00347756	.000402148

The fit represents the single detachment cross section with an rms deviation of 0.4%. The maximum deviation is 0.8% at 1.0E+04 eV/amu.

The fit represents the double detachment cross section with an rms deviation of 0.4%. The maximum deviation is 0.7% at 6.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$He^- + H_2 \rightarrow He + H_2 + e$
$He^- + H_2 \rightarrow He^+ + H_2 + 2e$



Cross Sections for Single and Double Electron Detachment of He^- in He

$$He^- + He \rightarrow He + He + e$$
$$\rightarrow He^+ + He + 2e$$

He + He + e		He^+ + He + 2e	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
1.2E+02	1.27E-15	2.7E+04	4.18E-17
2.0E+02	1.29E-15	3.0E+04	4.16E-17
3.0E+02	1.32E-15	4.0E+04	4.05E-17
4.0E+02	1.33E-15	5.0E+04	3.90E-17
7.0E+02	1.35E-15	6.0E+04	3.82E-17
1.0E+03	1.35E-15	7.0E+04	3.71E-17
2.0E+03	1.32E-15	8.0E+04	3.58E-17
4.0E+03	1.23E-15	9.0E+04	3.47E-17
7.0E+03	1.12E-15	1.0E+05	3.37E-17
1.0E+04	1.04E-15	1.5E+05	2.98E-17
2.0E+04	8.07E-16	2.0E+05	2.66E-17
4.0E+04	5.72E-16	3.0E+05	2.23E-17
7.0E+04	4.07E-16	4.0E+05	1.93E-17
1.0E+05	3.22E-16	5.0E+05	1.72E-17
2.0E+05	1.94E-16	6.0E+05	1.58E-17
4.0E+05	1.11E-16	7.0E+05	1.47E-17
7.0E+05	6.98E-17	8.0E+05	1.36E-17

References: 519, 521, 522, 525, 575, 576, 577, 578, 581, 582, 583

Accuracy: 25%

Notes: (1) The total electron detachment cross section (single and double) has been measured by Windham et al. (Ref. 575); Nicholas et al. (Ref. 578); Simpson and Gilbody (Ref. 519).

(2) For single detachment see: Sweetman (Ref. 576); Ryding et al. (Ref. 577); Heinemeier et al. (Ref. 581); Heinemeier et al. (Ref. 521); Coggiola (Ref. 582).

(3) For double detachment see: Ryding et al. (Ref. 577); Heinemeier et al. (Ref. 521).

Chebyshev Fitting Parameters for Cross Sections

Single det. E_{min} = 1.2E+02 eV/amu, E_{max} = 7.0E+05 eV/amu
Double det. E_{min} = 2.7E+04 eV/amu, E_{max} = 8.0E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
Single det.	-70.3313	-1.40200	-.626470	-.0569330	.0458639	.0102379	-.00527677
Double det.	-76.3309	-.571055	-.112122	.0109146	.00565444	.000461742	-.00146987

The fit represents the single detachment cross section with an rms deviation of 0.5%. The maximum deviation is 1.1% at 4.0E+03 eV/amu.

The fit represents the double detachment cross section with an rms deviation of 0.4%. The maximum deviation is 0.6% at 5.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$He^- + He \rightarrow He + He + e$
$He^- + He \rightarrow He^+ + He + 2e$



Cross Sections for Single and Double Electron Detachment of Li^- in H_2

$$Li^- + H_2 \rightarrow Li + H_2 + e$$
$$\rightarrow Li^+ + H_2 + 2e$$

$Li + H_2 + e$

Energy (eV/amu)	Cross Section (cm^2)
1.5E+03	1.89E-15
2.0E+03	1.91E-15
4.0E+03	1.81E-15
7.0E+03	1.64E-15
1.0E+04	1.50E-15
1.5E+04	1.34E-15
2.0E+04	1.23E-15
4.0E+04	9.45E-16
7.0E+04	7.43E-16
1.0E+05	6.30E-16
1.5E+05	5.15E-16
2.0E+05	4.46E-16
4.0E+05	3.10E-16

$Li^+ + H_2 + 2e$

Energy (eV/amu)	Cross Section (cm^2)
1.5E+04	1.00E-16
2.0E+04	8.79E-17
3.0E+04	7.40E-17
4.0E+04	6.67E-17
5.0E+04	6.27E-17
6.0E+04	5.98E-17
7.0E+04	5.76E-17
8.0E+04	5.55E-17
9.0E+04	5.43E-17
1.0E+05	5.28E-17
1.5E+05	4.83E-17
1.8E+05	4.67E-17

References: 250, 522, 583, 588

Accuracy: 25%

Note: For total electron detachment (single and double) see: Allision et al. (Ref. 250). For both single and double detachment see: McCullough et al. (Ref. 588).

Chebyshev Fitting Parameters for Cross Sections

Single det. E_{min} = 1.5E+03 eV/amu, E_{max} = 4.0E+05 eV/amu
Double det. E_{min} = 1.5E+04 eV/amu, E_{max} = 1.8E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
Single det.	-69.2100	-.932154	-.198400	.0257913	-.00111248	.00258484	-.00164526
Double det.	-74.5383	-.376194	.0491243	-.00831956	-.00149259	.00388870	-.000530605

The fit represents the single detachment cross section with an rms deviation of 0.2%. The maximum deviation is 0.3% at 1.0E+04 eV/amu.

The fit represents the double detachment cross section with an rms deviation of 0.2%. The maximum deviation is 0.4% at 8.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$$Li^- + H_2 \rightarrow Li + H_2 + e$$
$$Li^- + H_2 \rightarrow Li^+ + H_2 + 2e$$



Cross Sections for Total Electron Detachment of Li^- on He

$Li^- + He \rightarrow Li, Li^+$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.8E+02	1.86E+07	4.22E-16
2.0E+02	1.96E+07	4.56E-16
4.0E+02	2.78E+07	6.65E-16
7.0E+02	3.68E+07	8.07E-16
1.0E+03	4.39E+07	8.83E-16
1.5E+03	5.38E+07	9.57E-16
2.0E+03	6.21E+07	9.93E-16
4.0E+03	8.79E+07	1.05E-15
7.0E+03	1.16E+08	1.04E-15
1.0E+04	1.39E+08	1.00E-15
1.5E+04	1.70E+08	9.42E-16
2.0E+04	1.96E+08	8.87E-16
4.0E+04	2.78E+08	7.43E-16
6.0E+04	3.40E+08	6.46E-16

References: 250, 522, 551, 583, 587

Accuracy: 25%

Note: Reported values for cross sections correspond to the sum of those for single and double detachment.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.8E+02 eV/amu, E_{max} = 6.0E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-69.6625	.18135	-.342099	.0241112	-.0117377	.00823347	-.00361753

The fit represents the above cross section with an rms deviation of 0.2%. The maximum deviation is 0.4% at 2.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$Li^- + He \rightarrow Li, Li^+$



G. Cross Sections for Dissociation of Molecules and Molecular Ions

	Page
$H + H_2 \rightarrow H^- + (H^+ + H^+) + e$	G-2
$\rightarrow H + (H^+ + H^+) + 2e$	G-2
$\rightarrow H^+ + (H^+ + H^+) + 3e$	G-2
$\rightarrow H + (H + H^+) + e$	G-4
$\rightarrow H^+ + (H + H^+) + 2e$	G-6
$\rightarrow H^- + (H^+ + H)$	G-8
$H^+ + H_2 \rightarrow H^+ + H^+ + H^+ + 2e$	G-10
$He^+ + H_2 \rightarrow He^+ + H^+ + H^+ + 2e$	G-12
$He^{2+} + H_2 \rightarrow He^{2+} + (H^+ + H) + e$ (via $1s_{\sigma g}$)	G-14
$\rightarrow He^{2+} + (H^+ + H) + e$ (via $2p_{\sigma u}$)	G-16
$\rightarrow He^+ + (H^+ + H)$	G-18
$\rightarrow He^+ + (H^+ + H^+) + e$	G-20
$\rightarrow He^{2+} + (H^+ + H^+) + 2e$	G-22
$\rightarrow He + (H^+ + H^+)$	G-24
$H_2^+ + H_2 \rightarrow H^+_{Proj.}$	G-26
$H_2^+ + H \rightarrow H^+_{Proj.}$	G-28
$H_2^+ + H_2 \rightarrow H_{Proj.}$	G-30
$\rightarrow (H^+ + H) + H_2$	G-32
$H_2^+ + He \rightarrow$ Total H^+	G-34
$\rightarrow (H^+ + H) + He$	G-36
$H_2 + H_2 \rightarrow H^+$(Fast,Total)	G-38
$\rightarrow$ H(Fast,Total)	G-40
$\rightarrow$ Total Destruction of Fast H_2 Proj.	G-42
$H_3^+ + H_2 \rightarrow$ Total Destruction of H_3^+	G-44
$H_2^+ + H_2 \rightarrow$ Total Destruction of H_2^+	G-46
$H_3^+ + H_2 \rightarrow H^+$(Total,Projectile)	G-48
$\rightarrow H_2^+$(Total,Projectile)	G-50
$\rightarrow$ H(Total,Projectile)	G-52
$\rightarrow H_2$(Total,Projectile)	G-54

G. (Cont'd)

H_3^+ + He -> H^+(Total) G-56
-> H_2^+ G-58

HeH^+ + H_2 -> Total HeH^+ Dissociation Prod. . G-60

HeH^+ + He -> Total HeH^+ Dissociation Prod. . G-62

Cross Sections for Double (Dissociative) Ionization, Double Ionization with Stripping, and Electron Capture with Ionization for H-Atom Impact on H_2

$$H + H_2 \rightarrow H + (H^+ + H^+) + 2e$$
$$H + H_2 \rightarrow H^+ + (H^+ + H^+) + 3e$$
$$H + H_2 \rightarrow H^- + (H^+ + H^+) + e$$

$(H + 2H^+ + 2e)$		$(H^+ + 2H^+ + 3e)$		$(H^- + 2H^+ + e)$	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
5.0E+03	1.62E-19	5.7E+03	1.06E-20	4.7E+03	2.18E-20
6.0E+03	2.59E-19	7.0E+03	1.65E-20	7.0E+03	6.65E-20
7.0E+03	3.90E-19	1.0E+04	3.87E-20	8.5E+03	1.00E-19
8.0E+03	5.40E-19	1.3E+04	9.90E-20	9.0E+03	1.17E-19
9.0E+03	7.04E-19	1.5E+04	1.38E-19	1.0E+04	1.51E-19
1.0E+04	8.88E-19	2.0E+04	2.79E-19	1.5E+04	3.22E-19
1.5E+04	1.91E-18	2.5E+04	4.68E-19	2.0E+04	3.69E-19
2.0E+04	2.58E-18	3.0E+04	6.32E-19	2.5E+04	3.47E-19
3.0E+04	3.22E-18	3.5E+04	7.82E-19	3.0E+04	3.05E-19
4.0E+04	3.35E-18	4.0E+04	9.19E-19	3.5E+04	2.70E-19
5.0E+04	3.11E-18	4.5E+04	1.00E-18	4.0E+04	2.43E-19
		5.0E+04	1.14E-18	4.5E+04	2.22E-19
		5.2E+04	1.17E-18	5.0E+04	2.05E-19

Reference: 30

Accuracy: Unknown

Note: Kinetic energy of the slow H^+ ions is approximately 9 eV.

Chebyshev Fitting Parameters for Cross Sections

$(H + 2H^+ + 2e)$	E_{min} =	5.0E+03	eV/amu,	E_{max} =	5.0E+04	eV/amu
$(H^+ + 2H^+ + 3e)$	E_{min} =	5.7E+03	eV/amu,	E_{max} =	5.2E+04	eV/amu
$(H^- + 2H^+ + e)$	E_{min} =	4.7E+03	eV/amu,	E_{max} =	5.0E+04	eV/amu

	A0	A1	A2	A3	A4	A5	A6
$(H + 2H^+ + 2e)$	-82.5844	1.50087	-.514162	-.0138263	.0274858	-.00853833	-.0116637
$(H^+ + 2H^+ + 3e)$	-86.7735	2.46227	-.301192	-.105842	.0379990	-.00417294	.0122290
$(H^- + 2H^+ + e)$	-86.8910	1.09636	-.754871	-.00932842	.0855477	.0300822	-.0393792

The fit represents the $(H + 2H^+ + 2e)$ cross section with an rms deviation of 0.8%. The maximum deviation is 1.1% at 6.0E+03 eV/amu.

The fit represents the $(H^+ + 2H^+ + 3e)$ cross section with an rms deviation of 1.8%. The maximum deviation is 2.9% at 1.0E+04 eV/amu.

The fit represents the $(H^- + 2H^+ + e)$ cross section with an rms deviation of 1.3%. The maximum deviation is 3.0% at 8.5E+03 eV/amu.

See appendix for Chebyshev fit details.

$H + H_2 \rightarrow H + (H^+ + H^+) + 2e$
$H + H_2 \rightarrow H^+ + (H^+ + H^+) + 3e$
$H + H_2 \rightarrow H^- + (H^+ + H^+) + e$

Cross Section vs. Energy



Cross Sections for Dissociative Ionization of H_2 in Collisions with Hydrogen Atoms.

$$H + H_2 \rightarrow H + (H + H^+) + e$$

via $1s\sigma_g$ State (Curve 1)		via $2p\sigma_u$ State (Curve 2)	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
5.0E+03	6.40E-19	5.0E+03	2.25E-18
6.0E+03	8.21E-19	6.0E+03	3.23E-18
7.0E+03	9.67E-19	7.0E+03	4.12E-18
8.0E+03	1.09E-18	8.0E+03	4.97E-18
9.0E+03	1.18E-18	9.0E+03	5.76E-18
1.0E+04	1.25E-18	1.0E+04	6.47E-18
1.5E+04	1.40E-18	1.5E+04	8.07E-18
2.0E+04	1.38E-18	2.0E+04	7.49E-18
2.5E+04	1.31E-18	2.5E+04	6.24E-18
3.0E+04	1.23E-18	3.0E+04	5.13E-18
3.5E+04	1.17E-18	3.5E+04	4.11E-18
4.0E+04	1.12E-18	4.0E+04	3.24E-18
4.5E+04	1.06E-18	4.5E+04	2.51E-18
5.0E+04	1.00E-18	5.0E+04	1.97E-18

Reference: 30

Accuracy: Unknown

Notes: (1) Cross section values plotted in curve 1 refer to the dissociation of the H_2^+ ion from the $1s\sigma_g$ state. Energy of the slow ion is Approximately zero. (2) Cross section values plotted in curve 2 refer to the electron transitions to the $2p\sigma_u$ state and also to all the excited electron states of the H_2^+ ion. Energy of the slow ion is approximately 7 eV.

Chebyshev Fitting Parameters for Cross Sections

(via $1s\sigma_g$ state) E_{min} = 5.0E+03 eV/amu, E_{max} = 5.0E+04 eV/amu
(via $2p\sigma_u$ state) E_{min} = 5.0E+03 eV/amu, E_{max} = 5.0E+04 eV/amu

	A0	A1	A2	A3	A4	A5	A6
(via $1s\sigma_g$ state)	-82.7739	.176787	-.277606	.0497140	-.000803387	-.00277426	-.00425677
(via $2p\sigma_u$ state)	-80.0822	-.050777	-.663490	-.0140905	.00850407	-.00531411	-.00795596

The fit represents the cross section (via $1s\sigma_g$ state) with an rms deviation of 0.2%. The maximum deviation is 0.4% at 3.0E+04 eV/amu.

The fit represents the cross section (via $2p\sigma_u$ state) with an rms deviation of 0.6%. The maximum deviation is 1.2% at 4.5E+04 eV/amu.

See appendix for Chebyshev fit details.

$$H + H_2 \rightarrow H + (H + H^+) + e$$



Cross Sections for Dissociative Ionization of H_2 by H-Atom Impact Accompanied by Stripping of the H Projectile

$$H + H_2 \rightarrow H^+ + (H + H^+) + 2e$$

via $1s\sigma_g$ state (Curve 1)		via $2p\sigma_u$ state (Curve 2)	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
5.0E+03	2.49E-20	5.0E+03	2.38E-20
6.0E+03	3.27E-20	6.0E+03	5.83E-20
7.0E+03	4.07E-20	7.0E+03	1.07E-19
8.0E+03	4.91E-20	8.0E+03	1.66E-19
9.0E+03	5.77E-20	9.0E+03	2.34E-19
1.0E+04	6.68E-20	1.0E+04	3.07E-19
1.5E+04	1.15E-19	1.5E+04	6.88E-19
2.0E+04	1.66E-19	2.0E+04	1.04E-18
3.0E+04	2.73E-19	2.5E+04	1.26E-18
4.0E+04	3.65E-19	3.0E+04	1.37E-18
5.0E+04	4.10E-19	3.5E+04	1.37E-18
		4.0E+04	1.32E-18
		4.5E+04	1.23E-18
		5.0E+04	1.13E-18

Reference: 30

Accuracy: Unknown

Notes: (1) Cross section values plotted in curve 1 refer to the dissociation of the H_2^+ ion via the $1s\sigma_g$ state of H_2^+. Energy of the slow ion is approximately zero. (2) Cross section values plotted in curve 2 refer to the dissociation of the H_2^+ ion via the $2p\sigma_u$ state and also all higher excited electronic states of the H_2^+ ion. Energy of the slow ion is approximately 7 eV.

Chebyshev Fitting Parameters for Cross Sections

(via $1s\sigma_g$ state) E_{min} = 5.0E+03 eV/amu, E_{max} = 5.0E+04 eV/amu
(via $2p\sigma_u$ state) E_{min} = 5.0E+03 eV/amu, E_{max} = 5.0E+04 eV/amu

	A0	A1	A2	A3	A4	A5	A6
(via $1s\sigma_g$ state)	-87.2426	1.43882	-.0948243	-.0308580	-.0190257	-.00744929	-.00364551
(via $2p\sigma_u$ state)	-84.8972	1.86566	-.764017	.0497586	-.0434187	.0151155	.00206675

The fit represents cross section (via $1s\sigma_g$ state) with an rms deviation of 0.1%. The maximum deviation is 0.1% at 2.0E+04 eV/amu.

The fit represents cross section (via $2p\sigma_u$ state) with an rms deviation of 0.3%. The maximum deviation is 0.6% at 1.5E+04 eV/amu.

See appendix for Chebyshev fit details.

$$H + H_2 \rightarrow H^+ + (H + H^+) + 2e$$



Cross Sections for Dissociative Electron Capture from H_2 by H-Atoms

$$H + H_2 \rightarrow H^- + (H^+ + H)$$

via $1s\sigma_g$ state (Curve 1)		via $2p\sigma_u$ state (Curve 2)	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
5.0E+03	8.20E-20	5.0E+03	2.91E-19
6.0E+03	1.05E-19	6.0E+03	4.72E-19
7.0E+03	1.26E-19	7.0E+03	6.57E-19
8.0E+03	1.45E-19	8.0E+03	8.34E-19
9.0E+03	1.63E-19	9.0E+03	9.97E-19
1.0E+04	1.79E-19	1.0E+04	1.15E-18
1.5E+04	2.18E-19	1.5E+04	1.45E-18
2.0E+04	1.81E-19	2.0E+04	1.13E-18
2.5E+04	1.31E-19	2.5E+04	7.75E-19
3.0E+04	1.02E-19	3.0E+04	5.33E-19
3.5E+04	8.51E-20	3.5E+04	3.61E-19
4.0E+04	7.39E-20	4.0E+04	2.48E-19
4.5E+04	6.68E-20	4.5E+04	1.78E-19
5.0E+04	6.17E-20	5.0E+04	1.27E-19

Reference: 30

Accuracy: Unknown

Notes: (1) Cross section values plotted in curve 1 refer to the dissociation of the H_2^+ ion via the $1s\sigma_g$ state of H_2^+. The energy of the slow particles is approximately zero. (2) Cross section values plotted in curve 2 refer to the dissociation of the H_2^+ ion via the $2p\sigma_u$ and also all excited electronic states of the H_2^+ ion. The energy of the slow particles is approximately 7 eV.

Chebyshev Fitting Parameters for Cross Sections

(via $1s\sigma_g$ state) E_{min} = 5.0E+03 eV/amu, E_{max} = 5.0E+04 eV/amu
(via $2p\sigma_u$ state) E_{min} = 5.0E+03 eV/amu, E_{max} = 5.0E+04 eV/amu

	A0	A1	A2	A3	A4	A5	A6
(via $1s\sigma_g$ state)	-87.2731	-.218951	-.522211	.0736627	.0971637	.00270336	-.0307172
(via $2p\sigma_u$ state)	-84.2975	-.438944	-.978064	.0211660	.0476037	.00159449	-.0175091

The fit represents cross section (via $1s\sigma_g$ state) with an rms deviation of 0.8%. The maximum deviation is 1.5% at 2.5E+04 eV/amu.

The fit represents cross section (via $2p\sigma_u$ state) with an rms deviation of 0.8%. The maximum deviation is 1.1% at 4.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$$H + H_2 \rightarrow H^- + (H + H^+)$$



Dissociative Double Ionization Cross Sections for

$$H^+ + H_2 \rightarrow H^+ + H^+ + H^+ + 2e$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.0E+05	4.39E+08	3.96E-19
1.5E+05	5.38E+08	1.79E-19
2.0E+05	6.21E+08	1.02E-19
3.0E+05	7.61E+08	4.90E-20
7.0E+05	1.16E+09	1.17E-20
1.0E+06	1.39E+09	7.16E-21
1.5E+06	1.70E+09	4.30E-21
2.0E+06	1.96E+09	3.26E-21
3.0E+06	2.40E+09	2.49E-21

Reference: 365

Accuracy: Within factor of 4

Note: The data are reported for the H_2 internuclear axis oriented at 90° and 30° relative to the ion beam direction and no apparent difference was found between the measured cross section values.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.0E+05 eV/amu, E_{max} = 3.0E+06 eV/amu

A0	A1	A2	A3	A4	A5	A6
-90.4247	-2.58979	.296205	.0536430	.00747606	.00133207	.00152342

The fit represents the above cross section with an rms deviation of 0.6%. The maximum deviation is 0.9% at 2.0E+05 eV/amu.

See appendix for Chebyshev fit details.

$$H^+ + H_2 \rightarrow H^+ + H^+ + H^+ + 2e$$



Cross Sections for the Dissociative Double Ionization of H_2 By Fast He^+

$$He^+ + H_2 \rightarrow He^+ + H^+ + H^+ + 2e$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.5E+03	6.95E+07	5.00E-18
3.0E+03	7.61E+07	4.22E-18
4.0E+03	8.79E+07	3.12E-18
7.0E+03	1.16E+08	1.62E-18
1.0E+04	1.39E+08	1.02E-18
1.5E+04	1.70E+08	5.89E-19
2.0E+04	1.96E+08	3.91E-19
4.0E+04	2.78E+08	1.40E-19
7.0E+04	3.68E+08	5.95E-20
9.0E+04	4.17E+08	4.02E-20

Reference: 482

Accuracy: The absolute cross section values are estimated to within a factor of 4.

Note: The data are reported for the H_2 internuclear axis oriented at 90° and 30° relative to the ion beam direction, and no apparent difference was found between the cross section values.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 2.5E+03 eV/amu, E_{max} = 9.0E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-84.2206	-2.43620	-.134853	.0237772	-.00240081	.000622206	-.00103783

The fit represents the above cross section with an rms deviation of 0.1%. The maximum deviation is 0.2% at 1.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$He^+ + H_2 \rightarrow He^+ + H^+ + H^+ + 2e$



Cross Sections for the Dissociative Ionization of H_2 by Fast He^{2+}

$$He^{2+} + H_2 \rightarrow He^{2+} + (H^+ + H) + e$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
3.8E+03	8.56E+07	1.10E-19
4.0E+03	8.79E+07	1.38E-19
5.0E+03	9.82E+07	2.74E-19
6.0E+03	1.08E+08	4.31E-19
7.0E+03	1.16E+08	5.63E-19
8.0E+03	1.24E+08	6.27E-19
9.0E+03	1.32E+08	6.48E-19
1.0E+04	1.39E+08	6.57E-19
1.5E+04	1.70E+08	6.65E-19
2.0E+04	1.96E+08	1.04E-18
2.5E+04	2.20E+08	1.81E-18

Reference: 95

Accuracy: 20%

Note: These cross sections values refer to the ionization with dissociation of H_2 via the $1s\sigma_g$ state of the H_2^+ ion.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 3.8E+03 eV/amu, E_{max} = 2.5E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-84.1813	1.12091	-.170391	.319582	.0335842	-.0467173	-.0198799

The fit represents the above cross section with an rms deviation of 1.0%. The maximum deviation is 1.3% at 5.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$$He^{2+} + H_2 \rightarrow He^{2+} + (H^+ + H) + e$$



Cross Sections for the Dissociative Ionization of H_2 by Fast He^{2+}

$$He^{2+} + H_2 \rightarrow He^{2+} + (H^+ + H) + e$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.5E+03	6.95E+07	1.46E-19
3.0E+03	7.61E+07	2.05E-19
4.0E+03	8.79E+07	4.28E-19
5.0E+03	9.82E+07	7.94E-19
6.0E+03	1.08E+08	1.05E-18
7.0E+03	1.16E+08	1.23E-18
8.0E+03	1.24E+08	1.35E-18
9.0E+03	1.32E+08	1.42E-18
1.0E+04	1.39E+08	1.46E-18
1.5E+04	1.70E+08	1.53E-18
2.0E+04	1.96E+08	2.23E-18
2.5E+04	2.20E+08	3.13E-18
3.0E+04	2.41E+08	4.22E-18

Reference: 95

Accuracy: 30%

Notes: (1) The cross section values are for the ionization with dissociation of H_2 via the $2p\sigma_u$ state of H_2^+. See reference 95. (2) The measured energy of the slow ion is approximately 7 eV.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 2.5E+03 eV/amu, E_{max} = 3.0E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-82.9850	1.52994	-.280153	.246249	.0922114	-.0977812	-.00881272

The fit represents the above cross section with an rms deviation of 2.1%. The maximum deviation is 2.9% at 1.5E+04 eV/amu.

See appendix for Chebyshev fit details.

$$He^{2+} + H_2 \rightarrow He^{2+} + (H^+ + H) + e$$



Cross Sections for Dissociative Electron Capture from H_2 By Fast He^{2+}

$$He^{2+} + H_2 \rightarrow He^{+} + (H^{+} + H)$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.3E+03	5.01E+07	2.77E-18
1.5E+03	5.38E+07	3.01E-18
2.0E+03	6.21E+07	3.47E-18
3.0E+03	7.61E+07	4.37E-18
4.0E+03	8.79E+07	5.25E-18
5.0E+03	9.82E+07	6.17E-18
6.0E+03	1.08E+08	6.96E-18
7.0E+03	1.16E+08	7.79E-18
8.0E+03	1.24E+08	8.51E-18
9.0E+03	1.32E+08	9.31E-18
1.0E+04	1.39E+08	1.02E-17
1.5E+04	1.70E+08	1.27E-17
2.0E+04	1.96E+08	1.29E-17
2.5E+04	2.20E+08	1.29E-17

Reference: 95

Accuracy: 20%

Note: The cross section values are for capture with dissociation from the $1s\sigma_g$ state of the H_2^+ ion.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.3E+03 eV/amu, E_{max} = 2.5E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-79.1499	.840316	-.0633082	-.0781071	-.0271766	.00205805	.00727217

The fit represents the above cross section with an rms deviation of 1.0%. The maximum deviation is 1.5% at 2.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$He^{2+} + H_2 \rightarrow He^{+} + (H^{+} + H)$



Cross Sections for Dissociative Electron Capture with Ionization in Collisions of Fast He^{2+} with H_2

$$He^{2+} + H_2 \rightarrow He^{+} + (H^{+} + H^{+}) + e$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.2E+03	4.81E+07	1.10E-16
1.5E+03	5.38E+07	1.01E-16
2.0E+03	6.21E+07	8.84E-17
2.5E+03	6.95E+07	8.01E-17
3.0E+03	7.61E+07	7.47E-17
3.5E+03	8.22E+07	6.84E-17
4.0E+03	8.79E+07	6.01E-17
5.0E+03	9.82E+07	5.04E-17
6.0E+03	1.08E+08	4.90E-17
7.0E+03	1.16E+08	5.11E-17
8.0E+03	1.24E+08	5.53E-17
9.0E+03	1.32E+08	6.10E-17
1.0E+04	1.39E+08	6.54E-17
1.5E+04	1.70E+08	7.88E-17
2.0E+04	1.96E+08	8.57E-17
2.5E+04	2.20E+08	8.70E-17

Reference: 95

Accuracy: 30%

Note: The measured energy of the slow H^{+} ions is approximately 9 eV.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.2E+03 eV/amu, E_{max} = 2.5E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-74.1562	-.104398	.277805	.00632922	-.107311	-.0247870	.0511068

The fit represents the above cross section with an rms deviation of 2.9%. The maximum deviation is 3.9% at 2.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$$He^{2+} + H_2 \rightarrow He^{+} + (H^{+} + H^{+}) + e$$



Cross Sections for Double (Dissociative) Ionization of H_2 By Fast He^{2+}

$$He^{2+} + H_2 \rightarrow He^{2+} + (H^+ + H^+) + 2e$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.5E+03	6.95E+07	1.44E-19
3.0E+03	7.61E+07	2.15E-19
4.0E+03	8.79E+07	4.24E-19
5.0E+03	9.82E+07	7.90E-19
6.0E+03	1.08E+08	1.07E-18
7.0E+03	1.16E+08	1.25E-18
8.0E+03	1.24E+08	1.35E-18
9.0E+03	1.32E+08	1.39E-18
1.0E+04	1.39E+08	1.44E-18
1.5E+04	1.70E+08	1.52E-18
2.0E+04	1.96E+08	2.24E-18
3.0E+04	2.41E+08	4.23E-18

Reference: 95

Accuracy: 30%

Note: The measured energy of the slow H^+ ions is approximately 9 eV.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 2.5E+03 eV/amu, E_{max} = 3.0E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-82.9515	1.54494	-.254833	.255864	.0749613	-.113726	-.0359609

The fit represents the above cross section with an rms deviation of 2.0%. The maximum deviation is 4.4% at 4.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$$He^{2+} + H_2 \rightarrow He^{2+} + (H^+ + H^+) + 2e$$



Cross Sections for (Dissociative) Double Electron Capture from H_2 By Fast He^{2+}

$$He^{2+} + H_2 \rightarrow He + (H^+ + H^+)$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.1E+03	4.61E+07	5.15E-17
1.5E+03	5.38E+07	3.00E-17
2.0E+03	6.21E+07	1.70E-17
2.5E+03	6.95E+07	1.45E-17
3.0E+03	7.61E+07	1.42E-17
4.0E+03	8.79E+07	1.68E-17
5.0E+03	9.82E+07	2.14E-17
6.0E+03	1.08E+08	2.55E-17
7.0E+03	1.16E+08	2.90E-17
8.0E+03	1.24E+08	3.23E-17
9.0E+03	1.32E+08	3.49E-17
1.0E+04	1.39E+08	3.75E-17
1.5E+04	1.70E+08	4.42E-17
2.0E+04	1.96E+08	4.59E-17
2.5E+04	2.20E+08	4.48E-17

Reference: 95

Accuracy: 15%

Notes: None

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.1E+03 eV/amu, E_{max} = 2.5E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-75.9688	.211347	.437455	-.355845	.0231353	.0708411	-.0517123

The fit represents the above cross section with an rms deviation of 1.2%. The maximum deviation is 2.9% at 2.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$He^{2+} + H_2 \rightarrow He + (H^+ + H^+)$



Cross Sections for the Conversion of Fast H_2^+ Ions Into Fast Protons in Passage Through H_2

$$H_2^+ + H_2 \rightarrow H^+_{Proj.}$$

Low Energy		High Energy	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
1.6E+00	3.31E-17	1.5E+03	2.09E-16
2.0E+00	6.46E-17	2.0E+03	2.26E-16
2.5E+00	8.09E-17	4.0E+03	2.42E-16
3.0E+00	8.66E-17	7.0E+03	2.31E-16
3.5E+00	8.67E-17	1.0E+04	2.13E-16
4.0E+00	8.58E-17	1.5E+04	1.88E-16
4.5E+00	8.33E-17	1.8E+04	1.80E-16
5.0E+00	7.96E-17	2.0E+04	1.86E-16
		3.0E+04	2.23E-16
		4.0E+04	2.30E-16
		5.0E+04	2.26E-16
		8.0E+04	1.99E-16
		1.0E+05	1.80E-16
		1.5E+05	1.46E-16
		2.0E+05	1.21E-16
		4.0E+05	7.43E-17
		7.0E+05	4.22E-17
		1.0E+06	2.70E-17
		1.5E+06	1.46E-17
		2.0E+06	1.11E-17
		4.0E+06	6.21E-18
		7.0E+06	4.11E-18
		1.0E+07	3.28E-18

<u>References:</u> 86, 208, 210, 214, 218, 219, 527, 528, 531, 557, 558, 559, 571, 590

<u>Accuracy:</u> 25%

<u>Note:</u> No attempt has been made to interpolate between the low-energy data and the data taken at higher energies.

Chebyshev Fitting Parameters for Cross Sections

Low Energy E_{min} = 1.6E+00 eV/amu, E_{max} = 5.0E+00 eV/amu
High Energy E_{min} = 1.5E+03 eV/amu, E_{max} = 1.0E+07 eV/amu

	A0	A1	A2	A3	A4	A5	A6
Low Energy	-74.4493	.351878	-.249279	.0781924	-.0295527	.00853617	-.00490330
High Energy	-74.9261	-2.19443	-.855834	.0421307	.216227	.0921147	-.0893079

The fit represents the Low Energy cross section with an rms deviation of 0.2%. The maximum deviation is 0.3% at 3.5E+00 eV/amu.

The fit represents the High Energy cross section with an rms deviation of 6.3%. The maximum deviation is 12.4% at 1.8E+04 eV/amu.

See appendix for Chebyshev fit details.

$$H_2^+ + H_2 \rightarrow H^+_{Proj.}$$



Cross Sections for the Conversion of Fast H_2^+ Ions Into Fast Protons in Passage Through H Atoms

$$H_2^+ + H \rightarrow H^+_{Proj.}$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.5E+03	5.38E+07	1.60E-16
2.0E+03	6.21E+07	1.83E-16
4.0E+03	8.79E+07	2.16E-16
7.0E+03	1.16E+08	2.20E-16
1.0E+04	1.39E+08	2.15E-16
1.5E+04	1.70E+08	2.06E-16
2.0E+04	1.96E+08	1.96E-16
4.0E+04	2.78E+08	1.69E-16
5.0E+04	3.11E+08	1.59E-16

Reference: 599

Accuracy: 25%

Notes: None

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.5E+03 eV/amu, E_{max} = 5.0E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-72.4228	-.0319629	-.156039	.0287214	-.00752456	.000228203	.000526627

The fit represents the above cross section with an rms deviation of 0.1%. The maximum deviation is 0.2% at 2.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$$H_2^+ + H \rightarrow H^+_{Proj.}$$



Cross Section for the Formation of Fast H Atoms by Passage of H_2^+ Through H_2

$$H_2^+ + H_2 \rightarrow H_{Proj.}$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.0E+03	6.21E+07	4.98E-16
4.0E+03	8.79E+07	7.09E-16
7.0E+03	1.16E+08	8.12E-16
1.0E+04	1.39E+08	8.22E-16
1.5E+04	1.70E+08	8.05E-16
2.0E+04	1.96E+08	7.44E-16
4.0E+04	2.78E+08	4.70E-16
7.0E+04	3.68E+08	2.21E-16
1.0E+05	4.39E+08	1.21E-16

References: 208, 210, 214, 218, 219, 271

Accuracy: 20%

Note: The data refer to the sum of the cross sections for $H_2^+ + H_2 \rightarrow (H^+ + H)_{proj.}$ and $H_2^+ + H_2 \rightarrow (H + H)_{proj.}$.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 2.0E+03 eV/amu, E_{max} = 1.0E+05 eV/amu

A0	A1	A2	A3	A4	A5	A6
-70.6702	-.632612	-.606521	-.0915143	-.0121710	.0168179	.0104797

The fit represents the above cross section with an rms deviation of 0.3%. The maximum deviation is 0.6% at 1.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$$H_2^+ + H_2 \rightarrow H_{Proj.}$$



Partial Cross Sections for Dissociation of Fast H_2^+ Ions in Passage Through H_2

$$H_2^+ + H_2 \rightarrow (H^+ + H) + H_2$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.0E+01	4.39E+06	6.14E-17
2.0E+01	6.21E+06	6.58E-17
4.0E+01	8.79E+06	6.96E-17
7.0E+01	1.16E+07	7.64E-17
1.0E+02	1.39E+07	8.16E-17
1.5E+02	1.70E+07	9.19E-17
2.0E+02	1.96E+07	1.01E-16
4.0E+02	2.78E+07	1.24E-16
7.0E+02	3.68E+07	1.47E-16
1.0E+03	4.39E+07	1.60E-16
1.5E+03	5.38E+07	1.65E-16
2.0E+03	6.21E+07	1.67E-16
4.0E+03	8.79E+07	1.68E-16
7.0E+03	1.16E+08	1.67E-16
1.0E+04	1.39E+08	1.65E-16
1.5E+04	1.70E+08	1.55E-16
2.0E+04	1.96E+08	1.47E-16
4.0E+04	2.78E+08	1.13E-16
7.0E+04	3.68E+08	8.34E-17
1.0E+05	4.39E+08	6.73E-17
1.5E+05	5.38E+08	5.12E-17
2.0E+05	6.21E+08	4.22E-17
4.0E+05	8.78E+08	2.58E-17
7.0E+05	1.16E+09	1.74E-17
1.0E+06	1.39E+09	1.34E-17

<u>References:</u> 271, 527, 528, 529, 560, 589, 591

<u>Accuracy:</u> $E < 5x10^4$ eV/amu, 40%
$E > 5x10^4$ eV/amu, 20%

<u>Notes:</u> (1) At the lower impact energies, the results of the various investigators depend on the initial vibrational-state distributions of the H_2^+ ion. (2) In the eV range, H_2^+ projectiles were used with D_2 target and D_2^+ projectiles with H_2 target (see Ref. 589).

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.0E+01 eV/amu, E_{max} = 1.0E+06 eV/amu

A0	A1	A2	A3	A4	A5	A6
-74.5723	-.555602	-.906658	-.238001	.110022	.0251065	.00343160

The fit represents the above cross section with an rms deviation of 2.6%. The maximum deviation is 5.8% at 4.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$$H_2^+ + H_2 \rightarrow (H^+ + H) + H_2$$



Cross Sections for the Formation of Fast Protons from H_2^+ in He

H_2^+ + He -> Total H^+

$\sigma_{H^+}(He)$

Energy (eV/amu)	Cross Section (cm^2)
1.0E+03	3.69E-17
1.5E+03	5.75E-17
2.0E+03	7.43E-17
4.0E+03	1.17E-16
7.0E+03	1.53E-16
1.0E+04	1.72E-16
1.5E+04	1.86E-16
2.0E+04	1.94E-16
3.0E+04	2.01E-16
3.5E+04	2.01E-16
4.0E+04	2.00E-16
7.0E+04	1.84E-16
1.0E+05	1.57E-16
1.5E+05	1.22E-16
2.0E+05	9.81E-17
4.0E+05	5.21E-17
6.0E+05	3.35E-17
6.5E+05	3.06E-17

<u>References:</u> 86, 527, 528, 531, 562, 570

<u>Accuracy:</u> 20%

<u>Notes:</u> (1) More recent cross-section data from Ref. 562 (between $2x10^3$ to $8x10^3$ eV/amu) exceed the earlier data by approximately 40% at $8x10^3$ eV/amu to approximately 360% at $2x10^3$ eV/amu. (2) $\sigma_{H^+}(He)$ refers to the sum: $\sigma(H^+ + H)_{Proj.} + 2\sigma(H^+ + H^+)_{Proj.}$

Chebyshev Fitting Parameters for Cross Sections

$\sigma_{H^+}(He)$ E_{min} = 1.0E+03 eV/amu, E_{max} = 6.5E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
$\sigma_{H^+}(He)$	-74.0323	-.0812968	-.900960	-.0399950	-.0205556	.0309465	.00869410

The fit represents the σ_{H^+} cross section with an rms deviation of 1.0%. The maximum deviation is 1.4% at 2.0E+05 eV/amu.

See appendix for Chebyshev fit details.

$H_2^+ + He \rightarrow H^+$



Cross Sections for Dissociation of Fast H_2^+ Ions in Passage Through He

$$H_2^+ + He \rightarrow (H^+ + H) + He$$

Low Energy		High Energy	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
9.7E+00	6.55E-17	2.0E+03	2.11E-16
1.1E+01	6.55E-17	4.0E+03	1.93E-16
1.2E+01	6.55E-17	7.0E+03	1.69E-16
1.3E+01	6.56E-17	1.0E+04	1.46E-16
1.3E+01	6.56E-17	1.5E+04	1.22E-16
1.5E+01	6.56E-17	2.0E+04	1.07E-16
1.5E+01	6.56E-17	4.0E+04	7.35E-17
1.6E+01	6.56E-17	7.0E+04	5.39E-17
1.7E+01	6.61E-17	1.0E+05	4.46E-17
1.8E+01	6.57E-17	1.5E+05	3.35E-17
1.9E+01	6.55E-17	2.0E+05	2.68E-17
2.0E+01	6.59E-17	4.0E+05	1.51E-17
2.1E+01	6.59E-17	7.0E+05	8.82E-18
2.2E+01	6.58E-17	1.0E+06	6.37E-18
2.3E+01	6.57E-17		
2.4E+01	6.55E-17		

References: 527, 528, 560, 562

Accuracy: $E > 5x10^4$ eV/amu, 20%
$2x10^3 < E < 8x10^3$ eV/amu, 15% (see Ref. 562)
$E < 10^2$ eV/amu, Unknown

Notes: (1) At lower impact energies the results of the various investigators depend on the initial vibrational state distribution of the H_2^+ ion. (2) The data at 10 to 25 eV/amu were taken with D_2^+ instead of H_2^+ as the projectile. (3) No attempt has been made to interpolate between the low and high energy data.

Chebyshev Fitting Parameters for Cross Sections

Low Energy E_{min} = 9.7E+00 eV/amu, E_{max} = 2.4E+01 eV/amu
High Energy E_{min} = 2.0E+03 eV/amu, E_{max} = 1.0E+06 eV/amu

	A0	A1	A2	A3	A4	A5	A6
Low Energy	-74.5250	.00171327	-.000929677	-.00133213	-.000620656	-.000378036	-.000458557
High Energy	-75.0072	-1.76360	-.334933	.000919287	-.0200220	.0135139	.0139802

The fit represents the Low Energy cross section with an rms deviation of 0.2%. The maximum deviation is 0.4% at 1.9E+01 eV/amu.

The fit represents the High Energy cross section with an rms deviation of 0.8%. The maximum deviation is 1.3% at 2.0E+05 eV/amu.

See appendix for Chebyshev fit details.

$H_2^+ + He \rightarrow (H^+ + H) + He$



Cross Sections for Formation of Fast H^+ for Fast H_2 Projectiles Passing Through H_2

$H_2 + H_2 \rightarrow H^+$ (Fast,Total)

Low E		High E	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
2.5E+03	2.04E-17	1.5E+05	6.13E-17
4.0E+03	2.62E-17	2.0E+05	4.61E-17
7.0E+03	3.61E-17	2.5E+05	3.70E-17
1.0E+04	4.52E-17	3.0E+05	3.13E-17
1.5E+04	5.74E-17	4.0E+05	2.56E-17
2.0E+04	6.71E-17	5.0E+05	2.25E-17
2.4E+04	6.89E-17	6.0E+05	2.12E-17
3.0E+04	6.25E-17		
4.0E+04	4.34E-17		
5.0E+04	2.80E-17		

References: 31, 535

Accuracy: Unknown

Notes: (1) The low energy data are from Ref. 31, while the high energy data are taken from Ref. 535. (2) This data involves the sum of the cross sections for the products: $\sigma(H^+ + H)_{Proj.} + 2\sigma(H^+ + H^+)_{Proj.} + \sigma(H^+ + H^-)_{Proj.}$. (3) No attempt has been made to join the high and low energy data sets.

Chebyshev Fitting Parameters for Cross Sections

Low E E_{min} = 2.5E+03 eV/amu, E_{max} = 5.0E+04 eV/amu
High E E_{min} = 1.5E+05 eV/amu, E_{max} = 6.0E+05 eV/amu

	A0	A1	A2	A3	A4	A5	A6
Low E	-75.7161	.371301	-.373363	-.209805	-.0677263	-.00389719	.0259767
High E	-75.8616	-.542987	.0642733	.0118753	-.00148111	.000221649	.00636358

The fit represents the Low E cross section with an rms deviation of 0.7%. The maximum deviation is 1.2% at 1.5E+04 eV/amu.

The fit represents the High E cross section with an rms deviation of 0.0%. The maximum deviation is 0.0% at 4.0E+05 eV/amu.

See appendix for Chebyshev fit details.

$H_2 + H_2 \rightarrow H^+$ (Fast, Total)



Cross Sections for the Formation of Fast H from Fast H_2 Projectiles Passing Through H_2

$H_2 + H_2 \rightarrow H$ (Fast,Total)

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
2.5E+03	6.95E+07	2.62E-16
3.0E+03	7.61E+07	2.89E-16
4.0E+03	8.79E+07	3.08E-16
5.0E+03	9.82E+07	3.08E-16
6.0E+03	1.08E+08	2.95E-16
7.0E+03	1.16E+08	2.85E-16
8.0E+03	1.24E+08	2.93E-16
9.0E+03	1.32E+08	3.13E-16
1.0E+04	1.39E+08	3.32E-16
1.3E+04	1.58E+08	3.46E-16
1.5E+04	1.70E+08	3.43E-16
2.0E+04	1.96E+08	3.16E-16
3.0E+04	2.41E+08	2.49E-16
4.0E+04	2.78E+08	2.00E-16
5.0E+04	3.11E+08	1.65E-16

References: 31, 536

Accuracy: 40%

Note: This data involves the sum of the cross sections for the products: $\sigma(H^+ + H)_{Proj.} + 2\sigma(H + H)_{Proj.}$. Variations in ion source conditions produce changes up to 10% in the measured values (see Ref. 536).

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 2.5E+03 eV/amu, E_{max} = 5.0E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-71.7329	-.200109	-.223241	-.0773361	-.0140887	.0563185	-.00955263

The fit represents the above cross section with an rms deviation of 2.5%. The maximum deviation is 4.4% at 7.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$H_2 + H_2 \rightarrow H$ (Fast, Total)



Total Destruction Cross Sections for Fast H_2 Projectile in H_2

$H_2 + H_2$ -> Total Destruction of Fast H_2 Projectile

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.3E+03	5.01E+07	1.59E-16
1.5E+03	5.38E+07	1.64E-16
2.0E+03	6.21E+07	1.76E-16
4.0E+03	8.79E+07	2.16E-16
7.0E+03	1.16E+08	2.64E-16
1.0E+04	1.39E+08	3.09E-16
1.5E+04	1.70E+08	3.45E-16
2.0E+04	1.96E+08	3.52E-16
4.0E+04	2.78E+08	3.17E-16
5.0E+04	3.11E+08	2.92E-16

Reference: 536

Accuracy: 20%

Notes: (1) This cross section is the sum of cross sections for all reactions that destroy the fast H_2 molecule in passage through H_2, i.e. those producing fast H_2^+, (H+H), ($H+H^+$), and (H^++H^+) products. (2) Large variations in ion source operating conditions were found to produce changes up to 10% in the measured cross section (see Ref. 536).

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.3E+03 eV/amu, E_{max} = 5.0E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-71.8671	.380700	-.137630	-.0886297	-.00459765	.0134628	.00197498

The fit represents the above cross section with an rms deviation of 0.6%. The maximum deviation is 1.2% at 7.0E+03 eV/amu.

See appendix for Chebyshev fit details.

$H_2 + H_2$ –> Total Destruction of Fast H_2 Projectiles



Total Destruction Cross Sections for H_3^+ Projectile Ions in H_2

$H_3^+ + H_2 \rightarrow$ Total Destruction of H_3^+

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.6E+03	5.56E+07	3.39E-16
2.0E+03	6.21E+07	3.78E-16
4.0E+03	8.79E+07	5.07E-16
7.0E+03	1.16E+08	6.10E-16
1.0E+04	1.39E+08	6.65E-16
1.5E+04	1.70E+08	7.06E-16
2.0E+04	1.96E+08	6.81E-16
3.0E+04	2.41E+08	5.93E-16
3.5E+04	2.60E+08	5.58E-16

Reference: 536

Accuracy: 20%

Notes: (1) This cross section is the sum of cross sections for all reactions that destroy the H_3^+ molecular ion in passage through H_2. (2) Large variations in ion source operating conditions were found to produce changes up to 10% in the measured cross sections (see Ref. 536).

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.6E+03 eV/amu, E_{max} = 3.5E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-70.3701	.284601	-.184305	-.0425670	-.00921848	.00698594	.00725332

The fit represents the above cross section with an rms deviation of 0.3%. The maximum deviation is 0.5% at 1.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$H_3^+ + H_2$ –> Total Destruction of H_3^+ Projectiles



Total Destruction Cross Sections for H_2^+ Projectile Ions in H_2

$H_2^+ + H_2 \rightarrow$ Total Destruction of H_2^+

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.5E+03	5.38E+07	8.12E-16
2.0E+03	6.21E+07	8.18E-16
4.0E+03	8.79E+07	8.30E-16
7.0E+03	1.16E+08	8.15E-16
1.0E+04	1.39E+08	8.18E-16
1.5E+04	1.70E+08	8.46E-16
2.0E+04	1.96E+08	7.75E-16
4.0E+04	2.78E+08	4.69E-16
5.0E+04	3.11E+08	3.81E-16

References: 536, 589

Accuracy: 20%

Notes: (1) This cross section is the sum of cross sections for all reactions that destroy the H_2^+ molecular ion in passage through H_2. (2) Large variations in ion source operating conditions were found to produce changes up to 10% in the measure cross sections (see Ref. 536).

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.3E+03 eV/amu, E_{max} = 5.0E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-69.7995	-.288081	-.216489	-.102343	-.0344599	.0155290	.0223268

The fit represents the above cross section with an rms deviation of 0.9%. The maximum deviation is 1.3% at 1.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$H_2^+ + H_2 \rightarrow$ Total Destruction of H_2^+ Projectiles



Cross Sections for Production of Fast H^+ From H_3^+ Projectiles in H_2

$H_3^+ + H_2 \rightarrow H^+$ (Total,Projectile)

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
4.0E+01	8.84E+06	6.00E-18
7.0E+01	1.16E+07	6.91E-18
1.0E+02	1.39E+07	7.61E-18
2.0E+02	1.96E+07	8.89E-18
4.0E+02	2.78E+07	1.19E-17
7.0E+02	3.68E+07	1.78E-17
1.0E+03	4.39E+07	2.49E-17
1.5E+03	5.38E+07	3.99E-17
2.0E+03	6.21E+07	5.41E-17
4.0E+03	8.79E+07	9.80E-17
7.0E+03	1.16E+08	1.42E-16
1.0E+04	1.39E+08	1.77E-16
2.0E+04	1.96E+08	2.14E-16
4.0E+04	2.78E+08	2.33E-16
7.0E+04	3.68E+08	2.39E-16
1.0E+05	4.39E+08	2.28E-16
2.0E+05	6.21E+08	1.74E-16
4.0E+05	8.78E+08	1.08E-16
6.0E+05	1.08E+09	7.15E-17

References: 86, 214, 535, 574

Accuracy: 30%

Note: Large variations in measured dissociation cross sections have been ascribed to the influence of H_3^+ ions formed in ion sources with varying degrees of vibrational excitation.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 4.0E+01 eV/amu, E_{max} = 6.0E+05 eV/amu

A0	A1	A2	A3	A4	A5	A6
-75.3369	1.74367	-.759749	-.559135	.0918355	.0438106	-.0940811

The fit represents the above cross section with an rms deviation of 3.4%. The maximum deviation is 5.3% at 7.0E+02 eV/amu.

See appendix for Chebyshev fit details.

$H_3^+ + H_2 \rightarrow H^+$ (Total, Projectile)



Cross Sections for Production of Fast H_2^+ From H_3^+ Projectiles in H_2

$H_3^+ + H_2 \rightarrow H_2^+$ (Total,Projectile)

Energy (eV/amu)	Cross Section (cm^2)
1.1E+02	1.34E-17
2.0E+02	1.99E-17
4.0E+02	2.93E-17
7.0E+02	4.17E-17
1.0E+03	5.09E-17
1.5E+03	6.53E-17
2.0E+03	7.96E-17
4.0E+03	1.18E-16
7.0E+03	1.29E-16
1.0E+04	1.29E-16
2.0E+04	1.18E-16
4.0E+04	9.62E-17
7.0E+04	7.75E-17
1.0E+05	6.62E-17
2.0E+05	4.69E-17
4.0E+05	2.86E-17
6.0E+05	2.03E-17

References: 86, 214, 535, 574

Accuracy: 30%

Note: Large variations in measured dissociation cross sections have been ascribed to the influence of H_3^+ ions formed in ion sources with varying degrees of vibrational excitation.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.1E+02 eV/amu, E_{max} = 6.0E+05 eV/amu

A0	A1	A2	A3	A4	A5	A6
-75.4231	.295854	-.985779	-.0762360	.0980699	-.0248092	-.0512818

The fit represents the cross section with an rms deviation of 2.4%. The maximum deviation is 4.1% at 4.0E+05 eV/amu.

See appendix for Chebyshev fit details.

$H_3^+ + H_2 \rightarrow H^{2+}$ (Total, Projectile)



Cross Sections for Production of Fast H from H_3^+ Projectiles in H_2

$H_3^+ + H_2 \rightarrow H$ (Total,Projectile)

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.5E+03	5.42E+07	3.20E-16
2.0E+03	6.21E+07	4.26E-16
4.0E+03	8.79E+07	6.94E-16
7.0E+03	1.16E+08	8.90E-16
1.0E+04	1.39E+08	9.69E-16
1.3E+04	1.58E+08	9.96E-16
2.0E+04	1.96E+08	9.02E-16
4.0E+04	2.78E+08	5.77E-16
7.0E+04	3.68E+08	3.56E-16
1.0E+05	4.39E+08	2.62E-16
2.0E+05	6.21E+08	1.45E-16
4.0E+05	8.78E+08	7.79E-17
6.0E+05	1.08E+09	5.48E-17

References: 214, 535

Accuracy: 30%

Note: Large variations in measured dissociation cross sections have been ascribed to the influence of H_3^+ ions formed in ion sources with varying degrees of vibrational excitation.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.5E+03 eV/amu, E_{max} = 6.0E+05 eV/amu

A0	A1	A2	A3	A4	A5	A6
-71.5391	-1.05347	-.826342	.203507	.0536140	-.0425785	-.0185315

The fit represents the above cross section with an rms deviation of 1.8%. The maximum deviation is 2.9% at 4.0E+05 eV/amu.

See appendix for Chebyshev fit details.

$H_3^+ + H_2 \rightarrow H$ (Total, Projectile)



Cross Sections for Production of Fast H_2 from H_3^+ Projectiles in H_2

$H_3^+ + H_2 \rightarrow H_2$ (Total,Projectile)

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
3.0E+01	7.61E+06	5.03E-17
4.0E+01	8.79E+06	5.18E-17
7.0E+01	1.16E+07	5.74E-17
1.0E+02	1.39E+07	6.29E-17
1.5E+02	1.70E+07	7.07E-17
2.0E+02	1.96E+07	7.82E-17
4.0E+02	2.78E+07	1.10E-16
7.0E+02	3.68E+07	1.48E-16
1.0E+03	4.39E+07	1.81E-16
1.5E+03	5.38E+07	2.25E-16
2.0E+03	6.21E+07	2.63E-16
4.0E+03	8.79E+07	3.64E-16
7.0E+03	1.16E+08	4.26E-16
8.0E+03	1.24E+08	4.31E-16
1.0E+04	1.39E+08	4.30E-16
1.5E+04	1.70E+08	3.83E-16
2.0E+04	1.96E+08	3.22E-16
4.0E+04	2.78E+08	1.50E-16
7.0E+04	3.68E+08	6.49E-17
1.0E+05	4.39E+08	3.65E-17
1.5E+05	5.38E+08	1.98E-17
2.0E+05	6.21E+08	1.34E-17
4.0E+05	8.78E+08	6.90E-18
6.0E+05	1.08E+09	6.09E-18

References: 214, 535, 585

Accuracy: 30%

Note: Large variations in measured dissociation cross sections have been ascribed to the influence of H_3^+ ions formed in ion sources with varying degrees of vibrational excitation.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 3.0E+01 eV/amu, E_{max} = 6.0E+05 eV/amu

A0	A1	A2	A3	A4	A5	A6
-74.8168	-.899995	-1.57067	-.379862	.384429	.264557	.0263143

The fit represents the above cross section with an rms deviation of 4.1%. The maximum deviation is 6.5% at 1.0E+05 eV/amu.

See appendix for Chebyshev fit details.

$H_3^+ + H_2 \rightarrow H_2$ (Total, Projectile)



Cross Sections for the Formation of Fast H^+ From the Dissociation of H_3^+ Projectiles in He

H_3^+ + He -> H^+ (Total)

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.0E+03	4.39E+07	2.19E-17
1.5E+03	5.38E+07	3.20E-17
2.0E+03	6.21E+07	4.18E-17
4.0E+03	8.79E+07	7.20E-17
7.0E+03	1.16E+08	1.02E-16
1.0E+04	1.39E+08	1.20E-16
1.5E+04	1.70E+08	1.36E-16
1.7E+04	1.81E+08	1.39E-16

Reference: 86

Accuracy: 20%

Notes: None

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.0E+03 eV/amu, E_{max} = 1.7E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-74.5849	.941781	-.146192	-.0143319	.000599232	-.00304737	.00231477

The fit represents the above cross section with an rms deviation of 0.1%. The maximum deviation is 0.2% at 1.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$H_3^+ + He \rightarrow H^+$ (Total)



Cross Sections for the Formation of Fast H_2^+ From the Dissociation of H_3^+ Projectiles in He

$$H_3^+ + He \rightarrow H_2^+$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.0E+03	4.39E+07	1.67E-17
1.5E+03	5.38E+07	2.50E-17
2.0E+03	6.21E+07	3.25E-17
4.0E+03	8.79E+07	5.48E-17
7.0E+03	1.16E+08	7.57E-17
1.0E+04	1.39E+08	9.00E-17
1.5E+04	1.70E+08	1.06E-16
1.7E+04	1.81E+08	1.10E-16

Reference: 86

Accuracy: 20%

Notes: None

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.0E+03 eV/amu, E_{max} = 1.7E+04 eV/amu

A0	A1	A2	A3	A4	A5	A6
-75.1127	.939201	-.132028	.00680760	.000497736	-.00339960	-.000620124

The fit represents the above cross section with an rms deviation of 0.0%. The maximum deviation is 0.0% at 1.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$H_3^+ + He \rightarrow H_2^+$



Total Cross Sections for the Dissociation of HeH^+ in H_2

$HeH^+ + H_2$ -> Total HeH^+ Dissociation Products

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.0E+04	1.39E+08	6.85E-16
1.5E+04	1.70E+08	6.33E-16
2.0E+04	1.96E+08	5.59E-16
4.0E+04	2.78E+08	3.65E-16
7.0E+04	3.68E+08	2.14E-16
1.0E+05	4.39E+08	1.41E-16
1.5E+05	5.38E+08	1.05E-16
2.0E+05	6.21E+08	9.07E-17

Reference: 610

Accuracy: 20%

Notes: (1) This reaction involves the total dissociation cross section obtained from the sum of the partial cross sections for the formation of fast projectile fragments, i.e. (He+H), (He^++H), (He+H^+) and (He^++H^+). (2) See Ref. 610 for partial cross sections in the 1-2 x 10^5 eV/amu energy range.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.0E+04 eV/amu, E_{max} = 2.0E+05 eV/amu

A0	A1	A2	A3	A4	A5	A6
-71.6345	-1.10463	-.128721	.0876009	.0373958	.00719887	-.0189657

The fit represents the above cross section with an rms deviation of 0.9%. The maximum deviation is 1.5% at 1.0E+05 eV/amu.

See appendix for Chebyshev fit details.

$HeH^+ + H_2 \rightarrow$ Total HeH^+ Dissociation



Total Cross Sections for the Dissociation of HeH^+ in He

HeH^+ + He -> Total HeH^+ Dissociation Products

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.0E+04	1.39E+08	2.09E-16
1.5E+04	1.70E+08	2.98E-16
2.0E+04	1.96E+08	3.14E-16
3.0E+04	2.41E+08	3.00E-16
4.0E+04	2.78E+08	2.73E-16
5.0E+04	3.11E+08	2.44E-16
6.0E+04	3.40E+08	2.15E-16
7.0E+04	3.68E+08	1.93E-16
8.0E+04	3.93E+08	1.73E-16
9.0E+04	4.17E+08	1.58E-16
1.0E+05	4.39E+08	1.44E-16
1.5E+05	5.38E+08	1.03E-16
2.0E+05	6.21E+08	8.54E-17

Reference: 610

Accuracy: 20%

Notes: (1) This reaction involves the total dissociation cross section obtained from the sum of the partial cross sections for the formation of fast projectile fragments, i.e. (He+H), (He^++H), (He+H^+) and (He^++H^+). (2) See Ref. 610 for partial cross sections in the 1-2 x 10^5 eV/amu energy range.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 1.0E+04 eV/amu, E_{max} = 2.0E+05 eV/amu

A0	A1	A2	A3	A4	A5	A6
-72.4377	-.561126	-.322720	.0959965	-.00323223	.017197	-.00704447

The fit represents the above cross section with an rms deviation of 0.3%. The maximum deviation is 0.6% at 6.0E+04 eV/amu.

See appendix for Chebyshev fit details.

$HeH^+ + He \rightarrow$ Total HeH^+ Dissociation



H. Cross Sections for Interchange Reactions

	Page
$H^+ + D_2 \rightarrow D^+ + HD$	H-2
$\rightarrow HD^+ + D$	H-2
$HD^+ + D_2 \rightarrow HD_2^+ + D$	H-4
$\rightarrow D_3^+ + H$	H-4
$HD^+(\upsilon=0) + He \rightarrow HeH^+ + D$	H-6
$HD^+(\upsilon=1) + He \rightarrow HeH^+ + D$	H-6
$HD^+(\upsilon=2) + He \rightarrow HeH^+ + D$	H-6
$HD^+(\upsilon=3) + He \rightarrow HeH^+ + D$	H-6
$HD^+(\upsilon=4) + He \rightarrow HeH^+ + D$	H-6
$H_2^+ + He \rightarrow HeH^+ + H$	H-8
$H_2^+ + H_2 \rightarrow H_3^+ + H$	H-10
$D_2^+ + D_2 \rightarrow D_3^+ + D$	H-12
$D_2^+(\upsilon=0) + H_2 \rightarrow HD_2^+ + H$	H-14
$D_2^+(\upsilon=1) + H_2 \rightarrow HD_2^+ + H$	H-14
$D_2^+(\upsilon=2) + H_2 \rightarrow HD_2^+ + H$	H-14
$D_2^+(\upsilon=3) + H_2 \rightarrow HD_2^+ + H$	H-14
$H_2^+(\upsilon=0) + D_2 \rightarrow HD_2^+ + D$	H-16
$H_2^+(\upsilon=1) + D_2 \rightarrow HD_2^+ + D$	H-16
$H_2^+(\upsilon=2) + D_2 \rightarrow HD_2^+ + D$	H-16
$H_2^+(\upsilon=3) + D_2 \rightarrow HD_2^+ + D$	H-16
$H_2^+(\upsilon=4) + D_2 \rightarrow HD_2^+ + D$	H-16

Cross Reactions for Interchange Reactions Between Protons and D_2

$$H^+ + D_2 \rightarrow D^+ + HD$$
$$\rightarrow HD^+ + D$$

$D^+ + HD$		$HD^+ + D$	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
4.5E-01	1.18E-15	2.2E+00	4.67E-18
7.0E-01	7.38E-16	4.0E+00	2.22E-17
1.0E+00	4.69E-16	4.5E+00	2.26E-17
1.5E+00	2.59E-16	7.0E+00	1.29E-17
2.0E+00	1.59E-16	1.0E+01	4.03E-18
4.0E+00	4.15E-17	1.5E+01	1.28E-18
7.0E+00	1.47E-17	2.0E+01	6.95E-19
1.0E+01	1.10E-17	4.0E+01	2.72E-19
1.5E+01	9.80E-18	7.0E+01	1.87E-19
2.0E+01	8.78E-18	1.0E+02	1.67E-19
4.0E+01	5.18E-18		
7.0E+01	2.90E-18		
1.0E+02	2.01E-18		

<u>References:</u> 602, 603, 604, 605, 606, 620

<u>Accuracy:</u> Unknown

<u>Notes:</u> (1) For proton energies about 5 eV the production of D^+ may proceed also via $H^+ + D_2 \rightarrow D^+ + D + H$. (2) For total reaction cross sections see: Maier (Ref. 602); Ochs and Teloy (Ref. 604); Schlier et al. (Ref. 605); and Holliday et al. (Ref. 603). (3) For relative yields of D^+ and HD^+ see: Krenos and Wolfgang (Ref. 620). (4) Rate coefficient for thermal energy is: $k = 3.7 \times 10^{-10}$ cm^3/s from Fehsenfeld et al. (Ref. 606).

<u>Chebyshev Fitting Parameters for Cross Sections</u>

$D^+ + HD$ E_{min} = 4.5E-01 eV/amu, E_{max} = 1.0E+02 eV/amu
$HD^+ + D$ E_{min} = 2.2E+00 eV/amu, E_{max} = 1.0E+02 eV/amu

	A0	A1	A2	A3	A4	A5	A6
$D^+ + HD$	-75.6468	-3.22911	.412113	.0393398	-.261749	.00129512	.120885
$HD^+ + D$	-82.0488	-2.46136	-.299817	.833081	-.341103	-.0343948	.0959803

The fit represents the $D^+ + HD$ cross section with an rms deviation of 4.9%. The maximum deviation is 9.3% at 7.0E+00 eV/amu.

The fit represents the $HD^+ + D$ cross section with an rms deviation of 3.5%. The maximum deviation is 7.0% at 1.0E+01 eV/amu.

See appendix for Chebyshev fit details.

$H^+ + D_2 \rightarrow D^+ + HD$
$H^+ + D_2 \rightarrow HD^+ + D$



Cross Sections for Interchange Reactions Between HD^+ and D_2

$$HD^+ + D_2 \rightarrow HD_2^+ + D$$
$$\rightarrow D_3^+ + H$$

$HD_2^+ + D$		$D_3^+ + H$	
Energy (eV/amu)	Cross Section (cm^2)	Energy (eV/amu)	Cross Section (cm^2)
6.0E-03	1.17E-14	6.0E-03	6.52E-15
7.0E-03	1.10E-14	7.0E-03	5.99E-15
1.0E-02	9.45E-15	1.0E-02	5.03E-15
1.5E-02	7.92E-15	1.5E-02	4.15E-15
2.0E-02	6.93E-15	2.0E-02	3.58E-15
4.0E-02	5.08E-15	4.0E-02	2.48E-15
7.0E-02	4.11E-15	7.0E-02	1.80E-15
1.0E-01	3.58E-15	1.0E-01	1.47E-15
1.5E-01	3.07E-15	1.5E-01	1.14E-15
2.0E-01	2.70E-15	2.0E-01	9.55E-16
4.0E-01	1.85E-15	4.0E-01	5.78E-16
7.0E-01	1.11E-15	7.0E-01	3.54E-16
1.0E+00	7.05E-16	1.0E+00	2.30E-16
1.5E+00	3.49E-16	1.5E+00	1.20E-16
2.0E+00	1.82E-16	2.0E+00	6.60E-17
4.0E+00	1.47E-17	4.0E+00	6.33E-18
4.3E+00	9.62E-18	4.3E+00	4.59E-18

<u>References:</u> 607, 608

<u>Accuracy:</u> Unknown

<u>Notes:</u> None

Chebyshev Fitting Parameters for Cross Sections

$HD_2^+ + D$ E_{min} = 6.0E-03 eV/amu, E_{max} = 4.3E+00 eV/amu
$D_3^+ + H$ E_{min} = 6.0E-03 eV/amu, E_{max} = 4.3E+00 eV/amu

	A0	A1	A2	A3	A4	A5	A6
$HD_2^+ + D$	-68.6683	-2.92522	-1.06891	-.570210	-.198731	-.0387800	-.0184986
$D_3^+ + H$	-70.3595	-3.11828	-.916067	-.449998	-.194126	-.0628098	-.00822327

The fit represents the $HD_2^+ + D$ cross section with an rms deviation of 1.3%. The maximum deviation is 2.9% at 4.3E+00 eV/amu.

The fit represents the $D_3^+ + H$ cross section with an rms deviation of 0.5%. The maximum deviation is 0.9% at 4.0E+00 eV/amu.

See appendix for Chebyshev fit details.

$HD^+ + D_2 \rightarrow HD_2^+ + D$
$HD^+ + D_2 \rightarrow D_3^+ + H$



Cross Sections for Interchange Reactions Between HD^+ and He

$$HD^+(\upsilon) + He \rightarrow HeH^+ + D$$

$\upsilon = 0$		$\upsilon = 1$		$\upsilon = 2$		$\upsilon = 3$		$\upsilon = 4$	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
1.8E-01	7.76E-18	1.9E-01	2.53E-17	2.0E-01	6.15E-17	2.0E-01	1.25E-16	1.8E-01	2.25E-16
2.0E-01	7.71E-18	2.0E-01	2.44E-01	3.0E-01	3.82E-17	2.5E-01	1.03E-16	2.0E-01	2.06E-16
3.0E-01	7.36E-18	3.0E-01	1.84E-17	4.0E-01	2.48E-17	3.0E-01	8.06E-17	3.0E-01	1.26E-16
4.0E-01	6.79E-18	4.0E-01	1.46E-17	5.0E-01	1.71E-17	4.0E-01	4.51E-17	4.0E-01	8.05E-17
5.0E-01	6.27E-18	5.0E-01	1.22E-17	6.0E-01	1.27E-17	5.0E-01	2.54E-17	5.0E-01	5.29E-17
6.0E-01	5.79E-18	6.0E-01	1.04E-17	7.0E-01	1.04E-17	6.0E-01	1.78E-17	6.0E-01	3.54E-17
7.0E-01	5.37E-18	7.0E-01	9.22E-18	8.0E-01	8.98E-18	7.0E-01	1.42E-17	7.0E-01	2.44E-17
8.0E-01	5.00E-18	8.0E-01	8.26E-18	9.0E-01	8.08E-18	8.0E-01	1.23E-17	8.0E-01	1.69E-17
9.0E-01	4.68E-18	9.0E-01	7.52E-18	1.0E+00	7.45E-18	9.0E-01	1.13E-17	9.0E-01	1.22E-17
1.0E+00	4.37E-18	1.0E+00	6.94E-18	1.5E+00	6.51E-18	1.0E+00	1.06E-17	1.0E+00	8.94E-18
1.5E+00	3.31E-18	1.5E+00	5.29E-18			1.5E+00	1.01E-17	1.5E+00	2.36E-18
1.7E+00	3.08E-18								

<u>References:</u> 609, 611

<u>Accuracy:</u> Unknown

<u>Notes:</u> (1) Cross sections are presented as a function of initial vibrational quantum number υ. (2) The HD^+ ion beam was prepared in a given vibrational state by selective photoionization of HD (Turner et al. Ref. 611). (3) The thermal-energy (~0.1 eV) rate coefficient k for the HD^+ + He reaction leading to all products in Smith et al. (Ref. 609) is: $k = (1.25 \pm 0.1)\times10^{-10}$ cm^3/s.

Chebyshev Fitting Parameters for Cross Sections

$\upsilon = 0$	E_{min} =	1.8E-01	eV/amu,	E_{max} =	1.7E+00	eV/amu
$\upsilon = 1$	E_{min} =	1.9E-01	eV/amu,	E_{max} =	1.5E+00	eV/amu
$\upsilon = 2$	E_{min} =	2.0E-01	eV/amu,	E_{max} =	1.5E+00	eV/amu
$\upsilon = 3$	E_{min} =	2.0E-01	eV/amu,	E_{max} =	1.5E+00	eV/amu
$\upsilon = 4$	E_{min} =	1.8E-01	eV/amu,	E_{max} =	1.5E+00	eV/amu

	A0	A1	A2	A3	A4	A5	A6
$\upsilon = 0$	-79.5223	-.474196	-.106022	.00910711	.00412170	.00308534	.00343162
$\upsilon = 1$	-78.0058	-.802720	.000681956	.0202372	.00181675	.000208519	.00196596
$\upsilon = 2$	-77.1958	-1.20851	.141092	.101616	-.00786021	-.0159891	.0143204
$\upsilon = 3$	-76.1495	-1.42278	.227207	.182149	-.0607895	-.0178064	.0326454
$\upsilon = 4$	-75.8801	-2.27104	-.369722	-.00864579	.00647117	.00107548	-.00589532

The fit represents the $\upsilon = 0$ cross section with an rms deviation of 0.1%. The maximum deviation is 0.1% at 4.0E-01 eV/amu.

The fit represents the $\upsilon = 1$ cross section with an rms deviation of 0.2%. The maximum deviation is 0.4% at 6.0E-01 eV/amu.

The fit represents the $\upsilon = 2$ cross section with an rms deviation of 0.2%. The maximum deviation is 0.5% at 6.0E-01 eV/amu.

The fit represents the $\upsilon = 3$ cross section with an rms deviation of 0.8%. The maximum deviation is 1.4% at 5.0E-01 eV/amu.

The fit represents the $\upsilon = 4$ cross section with an rms deviation of 0.2%. The maximum deviation is 0.6% at 8.0E-01 eV/amu.

See appendix for Chebyshev fit details.

$HD^+(\nu) + He \rightarrow HeH^+ + D, HeD^+ + H$



Cross Sections for Interchange Reactions Between H_2^+ and He

$$H_2^+ + He \rightarrow HeH^+ + H$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
5.6E-01	1.04E+06	5.13E-16
6.0E-01	1.08E+06	4.81E-16
7.0E-01	1.16E+06	4.17E-16
8.0E-01	1.24E+06	3.67E-16
9.0E-01	1.32E+06	3.24E-16
1.0E+00	1.39E+06	2.86E-16
1.5E+00	1.70E+06	1.66E-16
2.0E+00	1.96E+06	1.04E-16
3.0E+00	2.41E+06	4.91E-17
4.0E+00	2.78E+06	3.07E-17
5.0E+00	3.11E+06	2.23E-17
6.0E+00	3.40E+06	1.77E-17
7.0E+00	3.68E+06	1.49E-17
8.0E+00	3.93E+06	1.29E-17
9.0E+00	4.17E+06	1.16E-17
1.0E+01	4.39E+06	1.05E-17

References: 609, 611, 612, 613, 614, 615, 616, 617, 618, 619

Accuracy: Unknown

Notes: (1) For relative total cross sections see: Neynaber and Magnuson (Ref. 618). (2) For absolute cross sections as a function of vibrational states of H_2^+ see: Chupka and Russell (Ref. 616); Chupka et al. (Ref. 617); Turner et al. (Ref. 611); and Govers and Guyon (Ref. 619). (3) For absolute rate constants for production of HeH^+ as a function of the energy see: Von Koch and Friedman (Ref. 615); and Smith et al. (Ref. 609). (4) Rate coefficient k, at thermal energy for HeH^+ production is: $k = [1.4 \pm .15] \times 10^{-10}$ $cm^3/mol \cdot s$. (5) For data on $D_2^+ + He \rightarrow HeD^+$ see: Smith et al. (Ref. 609). (6) Rate coefficient k at thermal energy for HeD^+ production is: $k = [1.15 \pm 0.1] \times 10^{-10}$ $cm^3/mol.s$.

Chebyshev Fitting Parameters for Cross Sections

E_{min} = 5.6E-01 eV/amu, E_{max} = 1.0E+01 eV/amu

A0	A1	A2	A3	A4	A5	A6
-74.2766	-2.05811	-.0177676	.136014	.00555409	-.0251696	-.000631817

The fit represents the above cross section with an rms deviation of 0.6%. The maximum deviation is 1.4% at 3.0E+00 eV/amu.

See appendix for Chebyshev fit details.

$$H_2^+ + He \rightarrow HeH^+ + H$$



Cross Sections for Interchange Reactions Between H_2^+ and H_2

$$H_2^+ + H_2 \rightarrow H_3^+ + H$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.0E-02	1.39E+05	1.16E-14
2.0E-02	1.96E+05	1.03E-14
4.0E-02	2.78E+05	8.63E-15
7.0E-02	3.68E+05	7.20E-15
1.0E-01	4.39E+05	6.23E-15
2.0E-01	6.21E+05	4.62E-15
4.0E-01	8.79E+05	3.27E-15
7.0E-01	1.16E+06	2.43E-15
1.0E+00	1.39E+06	1.87E-15
1.5E+00	1.70E+06	1.25E-15
2.0E+00	1.96E+06	8.51E-16
4.0E+00	2.78E+06	2.09E-16
7.0E+00	3.68E+06	2.22E-17

<u>References:</u> 607, 612, 613, 621, 622, 623, 624, 625, 626, 627, 628, 629, 631, 632, 633, 636, 637

<u>Accuracy:</u> ± 45%

<u>Notes:</u> (1) For relative total cross sections see: Clow and Futrell (Ref. 625); and Vance and Bailey (Ref. 628). (2) For absolute rate constants as a function of the energy see: Clow and Futrell (Ref. 625); Bowers et al. (Ref. 626); and Reuben and Friedman (Ref. 627). (3) For energy distributions of H_3^+ see: Lees and Rol (Ref. 638); and Gentry et al. (Ref. 621). (4) For cross sections as a function of v see: Chupka et al. (Ref. 629); and Koyano and Tanaka (Ref. 633). (5) For k as a function of υ see: Theard and Hunterss (Ref. 613). (6) Rate coefficient k at thermal energy for production of H_3^+ is: $k = [2.0 \pm .15]\times10^{-9}$ cm^3/s, from Refs. 613, 621, 623, 625, 626, 627, 631, 637.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

E_{min} = 1.0E-02 eV/amu, E_{max} = 7.0E+00 eV/amu

A0	A1	A2	A3	A4	A5	A6
-67.8902	-2.58783	-1.02353	-.461932	-.229219	-.0763343	-.0170793

The fit represents the above cross section with an rms deviation of 1.2%. The maximum deviation is 2.7% at 2.0E+00 eV/amu.

See appendix for Chebyshev fit details.

$$H_2^+ + H_2 \rightarrow H_3^+ + H$$



Cross Sections for Interchange reactions Between D_2^+ and D_2

$$D_2^+ + D_2 \rightarrow D_3^+ + D$$

Energy (eV/amu)	Velocity (cm/s)	Cross Section (cm^2)
1.6E-01	5.56E+05	2.60E-15
2.0E-01	6.21E+05	2.43E-15
3.0E-01	7.61E+05	2.00E-15
4.0E-01	8.79E+05	1.69E-15
5.0E-01	9.82E+05	1.45E-15
6.0E-01	1.08E+06	1.29E-15
7.0E-01	1.16E+06	1.16E-15
8.0E-01	1.24E+06	1.06E-15
9.0E-01	1.32E+06	9.77E-16
1.0E+00	1.39E+06	9.07E-16
1.5E+00	1.70E+06	6.90E-16
2.0E+00	1.96E+06	5.55E-16
3.0E+00	2.41E+06	4.01E-16
4.0E+00	2.78E+06	3.15E-16
4.2E+00	2.85E+06	3.04E-16

<u>References:</u> 623, 625, 626, 627, 631, 647

<u>Accuracy:</u> Unknown

<u>Notes:</u> (1) For relative cross sections see: Clow and Futrell (Ref. 625). (2) For absolute rate constants as a function of energy see: Drewitz (Ref. 631); Harrison et al. (Ref. 623); Reuben and Friedman (Ref. 627); Clow and Futrell (Ref. 625); and Bowers et al. (Ref. 626). (3) For kinetic energy distributions of D_3^+ products see: Doverspike and Champion (Ref. 647). (4) Rate coefficient for production of D_3^+ is: $k = 1.55 \times 10^{-9}$ $cm^3/mol\ s$ from Refs. 623, 625, 626, 627, 631. The states of reactants are unknown.

<u>Chebyshev Fitting Parameters for Cross Sections</u>

E_{min} = 1.6E-01 eV/amu, E_{max} = 4.2E+00 eV/amu

A0	A1	A2	A3	A4	A5	A6
-69.1270	-1.09264	-.0827553	.0133959	-.0140139	.00530178	.00343451

The fit represents the above cross section with an rms deviation of 0.2%. The maximum deviation is .4% at 1.0E+00 eV/amu.

See appendix for Chebyshev fit details.

$$D_2^+ + D_2 \rightarrow D_3^+ + D$$



Cross Sections for Interchange Reactions Between D_2^+ and H_2

$$D_2^+(\upsilon) + H_2 \rightarrow HD_2^+ + H$$

$\upsilon = 0$		$\upsilon = 1$		$\upsilon = 2$		$\upsilon = 3$	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
5.8E-02	1.88E-15	5.7E-02	2.03E-15	5.8E-02	2.01E-15	5.8E-02	1.93E-15
6.0E-02	1.84E-15	6.0E-02	1.92E-15	6.0E-02	1.96E-15	6.0E-02	1.90E-15
7.0E-02	1.52E-15	7.0E-02	1.61E-15	7.0E-02	1.76E-15	7.0E-02	1.78E-15
8.0E-02	1.20E-15	8.0E-02	1.29E-15	8.0E-02	1.57E-15	8.0E-02	1.62E-15
9.0E-02	9.67E-16	9.0E-02	1.10E-15	9.0E-02	1.42E-15	9.0E-02	1.49E-15
1.0E-01	8.31E-16	1.0E-01	9.66E-16	1.0E-01	1.28E-15	1.0E-01	1.36E-15
1.5E-01	5.70E-16	1.5E-01	6.93E-16	1.5E-01	8.29E-16	1.5E-01	8.81E-16
2.0E-01	4.29E-16	2.0E-01	5.25E-16	2.0E-01	5.66E-16	2.0E-01	5.70E-16
3.0E-01	2.47E-16	2.5E-01	3.78E-16	2.5E-01	3.79E-16	2.5E-01	3.75E-16
3.5E-01	1.81E-16	3.0E-01	2.79E-16	3.0E-01	2.77E-16	3.0E-01	2.46E-16
4.0E-01	1.52E-16	3.5E-01	2.00E-16	3.5E-01	2.01E-16	3.5E-01	1.65E-16
4.6E-01	1.15E-16	4.0E-01	1.44E-16	4.0E-01	1.47E-16	4.0E-01	1.12E-16
		4.5E-01	1.04E-16	4.5E-01	1.04E-16	4.5E-01	7.55E-17
		4.6E-01	9.61E-17	4.6E-01	9.58E-17	4.6E-01	6.93E-17

References: 625, 626, 627, 634, 638, 647, 652

Accuracy: 25%

Notes: (1) For relative cross sections see: Clow and Futrell (Ref. 625). (2) For rate constants, energy dependence see: Bowers et al. (Ref. 626); and Clow and Futrell (Ref. 625). (3) For kinetic energy distributions of HD_2^+ products see: Lees and Rol (Ref. 638); Doverspike and Champion (Ref. 647), and Krenos et al. (Ref. 652). (4) Rate coeficient for reaction $D_2^+ + H_2 \rightarrow$ products is: $k = (3.0 \pm 0.6)\times10^{-9}$ cm^3/mols from Clow and Futrell (Ref. 625). (5) Data are presented as a function of υ, as reported in Anderson et al. (Ref. 634).

Chebyshev Fitting Parameters for Cross Sections

$\upsilon = 0$	E_{min} =	5.8E-02	eV/amu,	E_{max} =	4.6E-01	eV/amu
$\upsilon = 1$	E_{min} =	5.7E-02	eV/amu,	E_{max} =	4.6E-01	eV/amu
$\upsilon = 2$	E_{min} =	5.8E-02	eV/amu,	E_{max} =	4.6E-01	eV/amu
$\upsilon = 3$	E_{min} =	5.8E-02	eV/amu,	E_{max} =	4.6E-01	eV/amu

	A0	A1	A2	A3	A4	A5	A6
$\upsilon = 0$	-70.4879	-1.34605	-.0400746	-.0884724	.00327276	.0410225	-.0238082
$\upsilon = 1$	-70.3174	-1.41371	-.174436	-.131909	-.00594907	.0250115	-.0190350
$\upsilon = 2$	-70.1696	-1.46820	-.256068	-.0411502	-.0107313	-.00889023	-.0110482
$\upsilon = 3$	-70.3104	-1.61386	-.377596	-.0473991	-.00736333	.000736757	-.00411866

The fit represents the $\upsilon = 0$ cross section with an rms deviation of 1.3%. The maximum deviation is 3.3% at 3.5E-01 eV/amu.

The fit represents the $\upsilon = 1$ cross section with an rms deviation of 0.7%. The maximum deviation is 1.3% at 6.0E-02 eV/amu.

The fit represents the $\upsilon = 2$ cross section with an rms deviation of 0.7%. The maximum deviation is 1.4% at 2.5E-01 eV/amu.

The fit represents the $\upsilon = 3$ cross section with an rms deviation of 0.3%. The maximum deviation is 0.4% at 8.0E-02 eV/amu.

See appendix for Chebyshev fit details.

$$D^{2+}(\nu) + H_2 \rightarrow HD_2^{+} + H$$



Cross Sections for Interchange Reactions Between H_2^+ and D_2

$$H_2^+(\upsilon) + D_2 \rightarrow HD_2^+ + H$$

$\upsilon = 0$		$\upsilon = 1$		$\upsilon = 2$		$\upsilon = 3$		$\upsilon = 4$	
Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)	Energy (eV/amu)	σ (cm^2)
1.7E-01	2.92E-15	1.7E-01	2.53E-15	1.7E-01	2.46E-15	1.8E-01	2.45E-15	1.8E-01	2.47E-15
2.0E-01	2.91E-15	2.0E-01	2.43E-15	2.0E-01	2.41E-15	2.0E-01	2.39E-15	2.0E-01	2.40E-15
3.0E-01	2.64E-15	3.0E-01	2.15E-15	3.0E-01	2.12E-15	3.0E-01	2.13E-15	3.0E-01	2.14E-15
4.0E-01	2.26E-15	4.0E-01	1.91E-15	4.0E-01	1.91E-15	4.0E-01	1.91E-15	4.0E-01	1.95E-15
5.0E-01	1.92E-15	5.0E-01	1.72E-15	5.0E-01	1.71E-15	5.0E-01	1.71E-15	5.0E-01	1.80E-15
6.0E-01	1.62E-15	5.0E-01	1.68E-15	6.0E-01	1.54E-15	6.0E-01	1.58E-15	6.0E-01	1.67E-15
7.0E-01	1.36E-15	6.0E-01	1.49E-15	7.0E-01	1.42E-15	7.0E-01	1.46E-15	7.0E-01	1.56E-15
8.0E-01	1.14E-15	7.0E-01	1.32E-15	8.0E-01	1.30E-15	8.0E-01	1.37E-15	8.0E-01	1.47E-15
9.0E-01	9.72E-16	8.0E-01	1.19E-15	9.0E-01	1.21E-15	9.0E-01	1.29E-15	9.0E-01	1.39E-15
1.0E+00	8.51E-16	9.0E-01	1.08E-15	1.0E+00	1.13E-15	1.0E+00	1.23E-15	1.0E+00	1.32E-15
1.5E+00	5.48E-16	1.0E+00	9.81E-16	1.5E+00	8.39E-16	1.5E+00	9.64E-16	1.5E+00	1.06E-15
2.0E+00	4.24E-16	1.5E+00	6.61E-16	2.0E+00	6.65E-16	2.0E+00	7.74E-16	2.0E+00	8.60E-16
3.0E+00	3.02E-16	2.0E+00	5.13E-16	3.0E+00	4.65E-16	3.0E+00	5.21E-16	3.0E+00	5.53E-16
4.0E+00	2.39E-16	3.0E+00	3.71E-16	4.0E+00	3.58E-16	4.0E+00	3.73E-16	4.0E+00	3.55E-16
4.5E+00	2.19E-16	4.0E+00	3.07E-16	5.0E+00	2.90E-16	5.0E+00	2.85E-16	4.5E+00	2.84E-16
		5.0E+00	2.74E-16	5.4E+00	2.70E-16	5.2E+00	2.80E-16		

References: 625, 626, 627, 634, 638, 652

Accuracy: 25%

Notes: (1) For energy dependence of rate constants see: Bowers et al. (Ref. 626); Clow and Futrell (Ref. 625); and Reuben and Friedman (Ref. 627). (2) For kinetic energy distributions of HD_2^+ products see: Lees and Rol (Ref. 638); and Krenos et al. (Ref. 652). (3) For relative cross sections see: Clow and Futrell (Ref. 625). (4) Rate coefficient for reaction: $H_2^+ + D_2 \rightarrow$ Products is: $k = (3.2 \pm 0.6)\times10^{-9}$ cm^3/mol.s from Clow and Futrell (Ref. 625). (5) Data are presented as a function of υ, as reported in Anderson et al. (Ref. 634).

Chebyshev Fitting Parameters for Cross Sections

$\upsilon = 0$	E_{min} =	1.7E-01	eV/amu,	E_{max} =	4.5E+00	eV/amu
$\upsilon = 1$	E_{min} =	1.7E-01	eV/amu,	E_{max} =	5.0E+00	eV/amu
$\upsilon = 2$	E_{min} =	1.7E-01	eV/amu,	E_{max} =	5.4E+00	eV/amu
$\upsilon = 3$	E_{min} =	1.8E-01	eV/amu,	E_{max} =	5.2E+00	eV/amu
$\upsilon = 4$	E_{min} =	1.8E-01	eV/amu,	E_{max} =	4.5E+00	eV/amu

	A0	A1	A2	A3	A4	A5	A6
$\upsilon = 0$	-69.2456	-1.40946	-.128223	.139770	-.0170479	-.0330555	.00606671
$\upsilon = 1$	-69.2502	-1.19199	-.116093	.0911706	.0198857	-.0118162	-.000510451
$\upsilon = 2$	-69.1336	-1.12492	-.176104	.0165999	.000760447	.00273290	-.000454869
$\upsilon = 3$	-69.0426	-1.06802	-.213967	-.0368351	-.00493123	.0161013	.00775590
$\upsilon = 4$	-68.8659	-.992859	-.258499	-.0853432	-.0286104	-.00372291	.00520865

The fit represents the $\upsilon = 0$ cross section with an rms deviation of 1.3%. The maximum deviation is 2.0% at 1.0E+00 eV/amu.

The fit represents the $\upsilon = 1$ cross section with an rms deviation of 0.6%. The maximum deviation is 0.8% at 7.0E-01 eV/amu.

The fit represents the $\upsilon = 2$ cross section with an rms deviation of 0.3%. The maximum deviation is 0.7% at 3.0E-01 eV/amu.

The fit represents the $\upsilon = 3$ cross section with an rms deviation of 0.5%. The maximum deviation is 1.3% at 5.0E+00 eV/amu.

The fit represents the $\upsilon = 4$ cross section with an rms deviation of 0.2%. The maximum deviation is 0.3% at 3.0E+00 eV/amu.

See appendix for Chebyshev fit details.

$$H_2^+(\nu) + D_2 \rightarrow HD_2^+ + H$$



REFERENCES

REFERENCES

1. J. E. Bayfield, Phys. Rev. 185, 105 (1969).
2. V. A. Belyaev, B. G. Brezhnev, and E. M. Erastov, Sov. Phys.-JETP 25, 777 (1967).
3. W. L. Fite, R. T. Brackmann, and W. R. Snow, Phys. Rev. 112, 1161 (1958).
4. W. L. Fite, R. F. Stebbings, D. G. Hummer, and R. T. Brackmann, Phys. Rev. 119, 663 (1960).
5. W. L. Fite, A.C.H. Smith, and R. F. Stebbings, Proc. Roy. Soc. London, Sect. A268, 527 (1962).
6. H. B. Gilbody and G. Ryding, Proc. Roy. Soc. London, 291, 438 (1966).
7. P. Hvelplund and A. Andersen, Phys. Scr. 26, 375 (1982).
8. G. W. McClure, Phys. Rev. 148, 47 (1966).
9. J. H. Newman, J. D. Cogan, D. L. Ziegler, D. E. Nitz, R. D. Rundel, K. A. Smith, and R. F. Stebbings, Phys. Rev. A 25, 2976 (1982).
10. A. B. Wittkower, G. Ryding, and H. B. Gilbody, Proc. Phys. Soc. 89, 541 (1966).
11. J. E. Bayfield, Phys. Rev. Lett. 20, 1223 (1968).
12. Y. P. Chong and W. L. Fite, Phys. Rev. A 16, 933 (1977).
13. J. Hill, J. Geddes, and H. B. Gilbody, J. Phys. B 12, L341 (1979).
14. T. J. Morgan, J. Geddes, and H. B. Gilbody, J. Phys. B 6, 2118 (1973).
15. T. J. Morgan, J. Stone, and R. Mayo, Phys. Rev. A 22, 1460 (1980).
16. G. Ryding, A. B. Wittkower, and H. B. Gilbody, Proc. Phys. Soc. 89, 547 (1966).
17. T. Kondow, R. J. Girnius, Y. P. Chong, and W. L. Fite, Phys. Rev. A 10, 1167 (1974).
18. T. J. Morgan, J. Geddes, and H. B. Gilbody, J. Phys. B 6, 2118 (1973).
19. R. F. Stebbings, R. A. Young, C. L. Oxley, and H. Ehrhardt, Phys. Rev. 138, A1312 (1965).
20. E. W. Thomas, ed., Table 5-3-4, P. 99 in Excitation in Heavy Particle Collisions, Wiley-Interscience, New York, 1972 (Table contains corrections applied to Ref.19).
21. J. E. Bayfield and G. A. Khayrallah, Phys. Rev. A 12, 869 (1975).
22. R. E. Olson, A. Salop, R. A. Phaneuf, and F. W. Meyer, Phys. Rev. A 16, 1867 (1977).
23. W. L. Nutt, R. W. McCullough, K. Brady, M. B. Shah, and H. B. Gilbody, J. Phys. B 11, 1457 (1978).
24. M. B. Shah and H. B. Gilbody, J. Phys. B 11, 121 (1978).
25. M. B. Shah and H. B. Gilbody, J. Phys. B 7, 630 (1974).
26. J. F. Williams, Phys. Rev. 153, 116 (1967).
27. P. M. Stier and C. F. Barnett, Phys. Rev. 103, 896 (1956).

28. B. Van Zyl, T. Q. Le, and R. C. Amme, J. Chem. Phys. 74, 314 (1981).

29. F. Roussel, P. Pradel, and G. Spiess, Phys. Rev. A 16, 1854 (1977).

30. V. V. Afrosimov, G. A. Leiko, Yu. A. Mamaev, M. N. Panov, and N. V. Fedorenko, Sov. Phys.-JETP 35, 1070 (1972).

31. G. W. McClure, Phys. Rev. 134, A1226 (1964).

32. T. Jorgensen, Jr., C. E. Kuyatt, W. W. Lang, D. C. Lorents, and C. A. Sautter, Phys. Rev. 140, A1481 (1965).

33. J. Hill, J. Geddes, and H. B. Gilbody, J. Phys. B 12, 3341 (1979).

34. G. W. McClure, Phys. Rev. 166, 22 (1968).

35. von U. Schryber, Helv. Phys. Acta A 40, 1023 (1967).

36. V. V. Afrosimov, G. A. Leiko, Yu. A. Mamaev, and M. N. Panov, Sov. Phys.-JETP 29, 648 (1969).

37. C. F. Barnett and H. K. Reynolds, Phys. Rev. 109, 355 (1958).

38. K. H. Berkner, R. V. Pyle, and J. W. Stearns, Nucl. Fusion 10, 145 (1970).

39. W. H. Cramer, J. Chem. Phys. 35, 836 (1961).

40. F. J. de Heer, J. Schutten, and H. Moustafa, Physica 32, 1766 (1966).

41. M. G. Holliday, J. T. Muckerman, and L. Friedman, J. Chem. Phys. 54, 1058 (1971).

42. Von O. Hollricher, Z. Physik 187, 41 (1965).

43. D. W. Koopman, Phys. Rev. 154, 79 (1967).

44. G. W. McClure, Phys. Rev. 132, 1636 (1963).

45. F. Schwirzke, Z. Physik 157, 510 (1960).

46. B. Van Zyl and M. W. Gealy, Phys. Rev. A 35, 3741 (1987).

47. L. H. Toburen, M. Y. Nakai, and R. A. Langley, Phys. Rev. 171, 114 (1968).

48. J. F. Williams, Phys. Rev. 179, 240 (1969).

49. L. M. Welsh, K. H. Berkner, S. N. Kaplan, and R. V. Pyle, Phys. Rev. 158, 85 (1967).

50. V. F. Kozlov and S. A. Bondar, Sov. Phys.-JETP 23, 195 (1966).

51. J. F. Williams, Phys. Rev. 157, 97 (1967).

52. J. F. Williams, Phys. Rev. 150, 7 (1966).

53. L. H. Toburen and M. Y. Nakai, Phys. Rev. 177, 191 (1969).

54. Ya. M. Fogel, R. V. Mitin, V. F. Kozlov, and N. D. Romashko, Sov. Phys.-JETP 35, 390 (1959).

55. R. Hippler, W. Harbich, M. Faust, H. O. Lutz, and L. J. Dube, J. Phys. B 19, 1507 (1986).

56. M. B. Shah and H. B. Gilbody, J. Phys. B 9, 1933 (1976).

57. M. C. Van Hemert, E. F. Van Dishoeck, J. A. Van der Hart, and F. Koike, Phys. Rev. A 31, 2227 (1985).

58. T. G. Winter and N. F. Lane, Phys. Rev. A 17, 66 (1978).

59. M. B. Shah and H. B. Gilbody, Atomic Data for Fusion 3, 27 (1977), and private communication.

60. G. J. Lockwood, G. H. Miller, and J. M. Hoffman, Bull. Am. Phys. Soc. 21, 1266 (1976), and private communication.

61. C. Harel and A. Salin, J. Phys. B 16, 55 (1983).

62. E. Gustafsson and E. Lindholm, Ark. Physik 18, 219 (1960).

63. J. Moseley, W. Aberth, and J. R. Peterson, Phys. Rev. Lett. 24, 435 (1970).

64. R. D. Rundel, K. L. Aitken, and M.F.A. Harrison, J. Phys. B 2, 954 (1969).

65. B. Peart, R. Grey, and K. T. Dolder, J. Phys. B 9, L369 (1976).

66. T. D. Gaily and M.F.A. Harrison, J. Phys. B 3, L25 (1970).

67. S. Szucs, M. Karemera, M. Terao, and F. Brouillard, J. Phys. B 17, 1613 (1984).

68. B. Peart, M. A. Bennett, and K. T. Dolder, J. Phys. B 18, L439 (1985).

69. K. T. Dolder and B. Peart, Rep. Prog. Phys. 48, 1283 (1985).

70. R. Shakeshaft, Phys. Rev. A 18, 1930 (1978).

71. M. Kimura and W. R. Thorson, Phys. Rev. A 24, 1780 (1981).

72. M. C. Chidichimo-Frank and R. D. Piacentini, J. Phys. B 7, 548 (1974).

73. B. Peart, K. Rinn, and K. T. Dolder, J. Phys. B 16, 1461 (1983).

74. K. Rinn, F. Melchert, and E. Salzborn, J. Phys. B 18, 3783 (1985).

75. B. Peart, R. Grey, and K. T. Dolder, J. Phys. B 10, 2675 (1977).

76. G. C. Angel, E. C. Sewell, K. F. Dunn, and H. B. Gilbody, J. Phys. B 11, L297 (1978).

77. G. C. Angel, K. F. Dunn, E. C. Sewell, and H. B. Gilbody, J. Phys. B 11, L49 (1978).

78. T. G. Winter, Phys.Rev. A 25, 697 (1982).

79. T. G. Winter, G. J. Hatton, and N. F. Lane, Phys. Rev. A 22, 930 (1980).

80. W. Fritsch and C. D. Lin, J. Phys. B 15, 1255 (1982).

81. J.B.A. Mitchell, K. F. Dunn, G. C. Angel, R. Browning, and H. B. Gilbody, J. Phys. B 10, 1897 (1977).

82. J.B.H. Stedeford and J. B. Hasted, Proc. Roy. Soc. London A227, 466 (1955).

83. M. E. Rudd, R. D. DuBois, L. H. Toburen, C. A. Ratcliffe, and T. V. Goffe, Phys. Rev. A 28, 3244 (1983).

84. Z. Z. Latypov and A. A. Shaporenko, Sov. Phys.-Tech. Phys. 21, 1277 (1976).

85. R. K. Colegrave and D.B.L. Stephens, J. Phys. B 1, 856 (1968).

86. J. F. Williams and D.N.F. Dunbar, Phys. Rev. 149, 62 (1966).

87. V. V. Afrosimov, Yu. A. Mamaev, M. N. Panov, and N. V. Fedorenko, Sov. Phys.-Tech. Phys. 14, 109 (1969).

88. K. H. Berkner, S. N. Kaplan, G. A. Paulikas, and R. V. Pyle, Phys. Rev. 140, A729 (1965).

89. J. Desesquelles, G. D. Cao, and M. Dufay, C. R. Acad. Sci. Paris 262B, 1329 (1966).

90. L. I. Pivovar, M. T. Novikov, and V. M. Tubaev, Sov. Phys.-JETP 15, 1035 (1962).

91. A. Itoh, M. Asari, and F. Fukuzawa, J. Phys. Soc. Jpn. 48, 943 (1980).

92. J. E. Bayfield and G. A. Khayrallah, Phys. Rev. A 11, 920 (1975).

93. R. A. Baragiola and I. B. Nemirovsky, Nucl. Instrum Methods 110, 511 (1973).

94. S. K. Allison, Phys. Rev. 109, 76 (1958).

95. V. V. Afrosimov, G. A. Leiko, and M. N. Panov, Sov. Phys.-Tech. Phys. 25, 313 (1980).

96. M. E. Rudd, T. V. Goffe, A. Itoh, and R. D. DuBois, Phys. Rev. A 32, 829 (1985).

97. L. I. Pivovar, V. M. Tubaev, and M. T. Novikov, Sov. Phys.-JETP 14, 20 (1962).

98. H. B. Gilbody, K. F. Dunn, R. Browning, and C. J. Latimer, J. Phys. B 4, 800 (1971).

99. A. B. Wittkower, G. Levy, and H. B. Gilbody, Proc. Phys. Soc. 91, 862 (1967).

100. M. B. Shah and H. B. Gilbody, J. Phys. B 9, 2685 (1976).

101. S. K. Allison, J. Cuevas, and P. G. Murphy, Phys. Rev. 102, 1041 (1956).

102. H. Tawara, J. Phys. Soc. Jpn. 31, 871 (1971).

103. C. F. Barnett and P. M. Stier, Phys. Rev. 109, 385 (1958).

104. F. J. de Heer, J. Schutten, and H. Moustafa, Physica 32, 1793 (1966).

105. H. B. Gilbody, J. B. Hasted, J. V. Ireland, A. R. Lee, E. W. Thomas, and A. S. Whiteman, Proc. Roy. Soc., London A274, 40 (1963).

106. R.L.C. Wu and D. G. Hopper, Chem. Phys. 57, 385 (1981).

107. T. F. Moran and R. J. Conrads, J. Chem. Phys. 58, 3793 (1973).

108. H. Helm, J. Phys. B 10, 3683 (1977).

109. R. D. Rundel, D. E. Nitz, K. A. Smith, M. W. Geis, and R. F. Stebbings, Phys. Rev. A 19, 33 (1979).

110. E. A. Hinds and R. Novick, J. Phys. B 11, 2201 (1978).

111. R. Hegerberg, T. Stefansson, and M. T. Elford, J. Phys. B 11, 133 (1978).

112. B. I. Kikiani, Z. E. Saliya, and I. B. Bagdasarova, Sov. Phys.-Tech. Phys. 20, 364 (1975).

113. A. Galli, A. Giardini-Guidoni, and G. G. Volpi, Nuovo Cimento 26, 845 (1962).

114. R. F. Potter, J. Chem. Phys. 22, 974 (1954).

115. V. A. Belyaev, B. G. Brezhnev, and E. M. Erastov, Sov. Phys.-JETP 27, 924 (1968).

116. R. J. Conrads, W. Pomerance, and T. F. Moran, J. Chem. Phys. 57, 2468 (1972).

117. V. S. Nikolaev, I. S. Dmitriev, L. N. Fateeva, and Ya. A. Teplova, Sov. Phys.-JETP 13, 695 (1961).

118. F. L. Eisele and S. W. Nagy, J. Chem. Phys. 65, 752 (1978).

119. W. N. Shelton and P. A. Stoycheff, Phys. Rev. A 3, 613 (1971).

120. P. Mahadevan and G. D. Magnuson, Phys. Rev. 171, 103 (1968).

121. M. B. Shah, J. Geddes, and H. B. Gilbody, J. Phys. B 13, 4049 (1980).

122. E. P. Andreev, V. A. Ankudinov, and S. V. Bobashev, p. 309 in V International Conference on the Physics of Electronic and Atomic Collisions, Leningrad, USSR, July 17-23, 1967, Publishing House Nauka, Leningrad 1967.

123. J. E. Bayfield, Phys. Rev. 182, 115 (1969).

124. J. H. Birely and R. J. McNeal, Phys. Rev. A 5, 692 (1972).

125. R. H. Hughes, E. D. Stokes, S. Choe, and T. J. King, Phys. Rev. A 4, 1453 (1971).

126. L. Colli, F. Cristofori, G. E. Frigerio, and P. G. Sona, Phys. Lett. 3, 62 (1962).

127. H. B. Gilbody, Adv. At. Mol. Phys. 22, 143 (1986).

128. R. H. Hughes, T. J. King, and S. Choe, Phys. Rev. A 5, 644 (1972).

129. R. H. Hughes, H. R. Dawson, B. M. Doughty, D. B. Kay, and C. A. Stigers, Phys. Rev. 146, 53 (1966).

130. R. H. Hughes, H. R. Dawson, and B. M. Doughty, Phys. Rev. 164, 166 (1967).

131. J. C. Ford and E. W. Thomas, Phys. Rev. A 5, 1701 (1972).

132. D. H. Loyd and H. R. Dawson, Phys. Rev. A 11, 140 (1975).

133. I. D. Williams, J. Geddes, and H. B. Gilbody, J. Phys. B 15, 1377 (1982).

134. R. H. Hughes, C. A. Stigers, B. M. Doughty, and E. D. Stokes, Phys. Rev. A 1, 1424 (1970).

135. R. J. Knize, S. R. Lundeen, and F. M. Pipkin, Phys. Rev. A 29, 1114 (1984).

136. H. R. Dawson and D. H. Loyd, Phys. Rev. A 15, 43 (1977).

137. W. R. Hess, Phys. Rev. A 9, 2036 (1974).

138. W. R. Hess, Phys. Lett. 41A, 66 (1972).

139. R. H. Hughes, S. Lin, and L. L. Hatfield, Phys. Rev. 130, 2318 (1963).

140. K. H. Berkner, R. V. Pyle, J. W. Stearns, and J. C. Warren, Phys. Rev. 166, 44 (1968).

141. G. R. Hertel and W. S. Koski, J. Chem. Phys. 40, 3452 (1964).

142. P. H. Hvelplund, J. Heinemeier, E. H. Pedersen, and F. R. Simpson, J. Phys. B 9, 491 (1976).

143. L. J. Puckett, G. O. Taylor, and D. W. Martin, Phys. Rev. 178, 271 (1969).

144. M. E. Rudd, T. V. Goffe, and A. Itoh, Phys. Rev. A 32, 2128 (1985).

145. M. B. Shah and H. B. Gilbody, J. Phys. B 7, 256 (1974).

146. Z. Z. Latypov and A. A. Shaporenko, Sov. Phys.-Tech. Phys. 19, 976 (1975).

147. Z. Z. Latypov, N. N. Fedorenko, I. P. Flaks, and A. A. Shaporenko, Sov. Phys.-JETP Lett. 11, 116 (1970).

148. V. V. Afrosimov, G. A. Leiko, Yu. A. Mamaev, and M. N. Panov, Sov. Phys.-JETP 40, 661 (1975).

149. J. S. Cohen and J. N. Bardsley, Phys. Rev. A 18, 1004 (1978).

150. V. S. Nikolaev, L. N. Fateeva, I. S. Dmitriev, and Ya. A. Teplova, Sov. Phys.-JETP 14, 67 (1962).

151. T. Kusakabe, H. Hanaki, N. Nagai, K. Kuroda, N. Maeda, and M. Sakisaka, Nucl. Instrum. Methods 198, 577 (1982).

152. H. Schrey and B. Huber, Z. Phys. A 213, 401 (1975).

153. A. F. Ferguson and B. L. Moiseiwitsch, Proc. Phys. Soc. 74, 437 (1959).

154. R. J. Conrads, T. W. Nichols, J. C. Ford, and E. W. Thomas, Phys. Rev. A 7, 1928 (1973).

155. H. R. Dawson and D. H. Loyd, Phys. Rev. A 9, 166 (1974).

156. J. C. Ford and E. W. Thomas, Phys. Rev. A 5, 1694 (1972).

157. F. T. Chan and J. Eichler, Phys. Rev. A 20, 1841 (1979).

158. J. L. Edwards and E. W. Thomas, Phys. Rev. A 2, 2346 (1970).

159. E. P. Andreev, V. A. Ankudinov, and S. V. Bobashev, p. 307 in V International Conference on the Physics of Electronic and Atomic Collisions, Leningrad, USSR, July 17-23, 1967, Publishing House Nauka, Leningrad, 1967.

160. J. Lenormand, J. Phys. (Paris) 37, 699 (1976).

161. J. E. Bayfield, Atomic Physics Vol. 4, p. 397, ed. G. Zu. Putlitz, E. W. Weber, and A. Winnacker, Plenum Press, New York, 1974.

162. V. A. Esaulov, J. Phys. B 13, 4039 (1980).

163. F. J. de Heer, J. Van Eck, and J. Kistemaker, Vol. 1, p. 73, in VI International Conference on Ionization Phenomena in Gases, Paris, July 8-13, 1963.

164. J. S. Risley, F. J. de Heer, and C. B. Kerkdijk, p. 93 in IX International Conference on the Physics of Electronic and Atomic Collisions, Seattle, July 24-30, 1975, University of Washington Press Seattle, WA, 1976.

165. J. S. Risley, F. J. de Heer, and C. B. Kerkdijk, J. Phys. B 11, 1759 (1978).

166. F. J. de Heer, Advances in Atomic and Molecular Physics Vol. 2, P. 327, ed. D. R. Bates and I. Estermann, Academic Press, New York, 1966.

167. E. P. Andreev, V. A. Ankudinov, S. V. Bobashev, and V. B. Matveev, Sov. Phys-JETP 25, 232 (1967).

168. C. W. Newby, J. Phys. B 18, 1781 (1985).

169. M. Kimura and W. R. Thorson, Phys. Rev. A 24, 3019 (1981).

170. S. Datta and S. C. Mukherjee, J. Phys. B 13, 539 (1980).

171. G. J. Hatton, N. F. Lane, and T. G. Winter, J. Phys. B 12, L571 (1979).

172. D. Ciric, D. Dijkkamp, E. Vlieg, and F. J. de Heer, J. Phys. B 18, L17 (1985).

173. D. Ciric, D. Dijkkamp, E. Vlieg, and F. J. de Heer, J. Phys. B 18, 4745 (1985).

174. B. H. Bransden, C. J. Noble, and J. Chandler, J. Phys. B 16, 4191 (1983).

175. T. G. Winter and G. J. Hatton, Phys. Rev. A 21, 793 (1980).

176. H. J. Ludde and D. M. Dreizler, J. Phys. B 15, 2713 (1982).

177. G. A. Khayrallah and J. E. Bayfield, Phys. Rev. A 11, 930 (1975).

178. Dz. S. Belkic and R. K. Janev, J. Phys. B 6, 1020 (1973).

179. S. C. Mukherjee, K. Roy, and N. C. Sil, J. Phys. B 6, 467 (1973).

180. K. E. Banyard and B. J. Szuster, J. Phys. B 10, L503 (1977).

181. Dz. S. Belkic and R. Gayet, J. Phys. B 10, 1923 (1977).

182. J. W. Gallagher, B. H. Bransden, and R. K. Janev, J. Phys. Chem. Ref. Data 12, 873 (1983).

183. E. P. Andreev, V. A. Ankudinov, and S. V. Bobashev, Sov. Phys.-JETP 23, 375 (1966).

184. V. Dose, Helv. Phys. Acta 39, 683 (1966).

185. D. Pretzer, B. Van Zyl, and R. Geballe, p. 618 in III International Conference on the Physics of Electronic and Atomic Collisions, London, July 22-26, 1963, North-Holland Publishing Co., Amsterdam, 1964.

186. D. Pretzer, B. Van Zyl, and R. Geballe, Phys. Rev. Lett. 10, 340 (1963).

187. B. Van Zyl, M. W. Gealy, and H. Neumann, Phys. Rev. A 35, 4551 (1987).

188. G. H. Dunn, R. Geballe, and D. Pretzer, Phys. Rev. 128, 2200 (1962).

189. D. H. Crandall and D. H. Jaecks, Phys. Rev. A 4, 2271 (1971).

190. D. Jaecks, B. Van Zyl, and R. Geballe, Phys. Rev. 137, A340 (1965).

191. R. L. Fitzwilson and E. W. Thomas, Phys. Rev. A 3, 1305 (1971).

192. B. Van Zyl, M. W. Gealy, and H. Neumann, Phys. Rev. A 33, 2333 (1986).

193. R. J. McNeal and J. H. Birely, p. 815 in VII International Conference on the Physics of Electronic and Atomic Collisions, Amsterdam, July 26-30, 1971, North-Holland Publishing Co., Amsterdam, 1971.

194. D. A. Dahlberg, D. K. Anderson, and I. E. Dayton, Phys. Rev. 170, 127 (1968).

195. B. Van Zyl, D. Jaecks, D. Pretzer, and R. Geballe, Phys. Rev. 158, 29 (1967).

196. R. E. Miers, A. S. Schlachter, and L. W. Anderson, Phys. Rev. 183, 213 (1969).

197. H. B. Gilbody, K. F. Dunn, R. Browning, and C. J. Latimer, p. 785 in VII International Conference on the Physics of Electronic and Atomic Collisions, Amsterdam, July 26-30, 1971, North-Holland Publishing Co., Amsterdam, 1971.

198. H. B. Gilbody, R. Browning, G. Levy, A. I. McIntosh, and K. F. Dunn, J. Phys. B 1, 863 (1968).

199. R. C. Isler and R. D. Nathan, Phys. Rev. A 6, 1036 (1972).

200. G. N. Polyakova, V. F. Erko, and A. V. Zats, Opt. Spectrosc. 29, 219 (1970).

201. P. J. MacVicar-Whelan and W. L. Borst, Phys. Rev. A 1, 314 (1970).

202. D. P. Sural, S. C. Mukherjee, and N. C. Sil, Phys. Rev. 164, 156 (1967).

203. R. Hippler, K -H. Schartner, and H. F. Beyer, J. Phys. B 11, L337 (1978).

204. L. W. Muller and and F. J. de Heer, Physica 48, 345 (1970).

205. F. J. de Heer, L. W. Muller, and R. Geballe, Physica 31, 1745 (1965).

206. C. E. Head and R. H. Hughes, Phys. Rev. 139, A1392 (1965).

207. F. J. de Heer and J. Van den Bos, Physica 31, 365 (1965).

208. D. R. Sweetman, Proc. Roy. Soc. London A256, 416 (1960).

209. H. B. Gilbody, Phys. Scr. 24, 712 (1981).

210. R. Schmid, Z. Phys. 161, 550 (1961).

211. D. W. Vance and T. L. Bailey, J. Chem. Phys. 44, 486 (1966).

212. H. L. Rothwell, B. Van Zyl, and R. C. Amme, J. Chem. Phys. 61, 3851 (1974).

213. C. J. Latimer, R. Browning, and H. B. Gilbody, J. Phys. B 2, 1055 (1969).

214. G. W. McClure, Phys. Rev. 130, 1852 (1963).

215. N. Noda, J. Phys. Soc. Jpn. 43, 1021 (1977).

216. H. C. Hayden and R. C. Amme, Phys. Rev. 172, 104 (1968).

217. Yu. S. Gordeev and M. N. Panov, Sov. Phys.-Tech. Phys. 9, 656 (1964).

218. A. Bouliou, M. Sarret, and P. Frere, C. R. Acad. Sci. Paris 296, 1217 (1983).

219. E. S. Solov'ev, R. N. Il'in, V. A. Oparin, and N. V. Fedorenko, Sov. Phys-JETP 26, 1097 (1968).

220. H. B. Gilbody and J. B. Hasted, Proc. Roy. Soc. London A238, 334 (1956).

221. F. M. Campbell, R. Browning, and C. J. Latimer, J. Phys. B 14, 3493 (1981).

222. T. F. Moran, K. J. McCann, M. R. Flannery, and D. L. Albritton, J. Chem. Phys. 65, 3172 (1976).

223. R. N. Stocker and H. Neumann, J. Chem. Phys. 61, 3852 (1974).

224. E. L. Duman and A. V. Evseev, Sov. Phys.-JETP 58, 1108 (1983).

225. C-Y. Lee and A. E. DePristo, J. Chem. Phys. 80, 1116 (1984).

226. T. F. Moran, M. R. Flannery, and D. L. Albritton, J. Chem. Phys. 62, 2869 (1975).

227. M. R. Flannery, J. V. Hornstein, and T. F. Moran, Chem. Phys. Lett. 32, 455 (1975).

228. Z. Karpas, V. Anicich, and W. T. Huntress, Jr., J. Chem. Phys. 70, 2877 (1979).

229. R. D. Smith, D. L. Smith, and J. H. Futrell, Int. J. Mass Spectrom Ion Phys. 19, 369 (1976).

230. S. L. Varghese, W. Waggoner, and C. L. Cocke, Phys. Rev. A 29, 2453 (1984).

231. R. D. DuBois and L. H. Toburen, Phys. Rev. A 31, 3603 (1985).

232. R. D. DuBois, Phys. Rev. A 32, 3319 (1985).

233. F. Aumayr and H. Winter, Phys. Rev. A 31, 67 (1985).

234. M. B. Shah, D. S. Elliott, and H. B. Gilbody, J. Phys. B 18, 4245 (1985).

235. W. Gruebler, P. A. Schmelzbach, V. Konig, and P. Marmier, Helv. Phys. Acta 43, 254 (1970).

236. R. N. Il'in, V. A. Oparin, E. S. Solov'ev, and N. V. Fedorenko, Sov. Phys.-JETP Lett. 2, 197 (1965).

237. R. N. Il'in, V. A. Oparin, E. S. Solov'ev, and N. V. Fedorenko, Sov. Phys.-Tech. Phys. 11, 921 (1967).

238. B. A. D'yachkov, Sov. At. Energy 27, 958 (1969).

239. O. Auciello, E. V. Alonso, and R. A. Baragiola, Phys. Rev. A 13, 985 (1976).

240. D. Dijkkamp, A. Boellaard, and F. J. de Heer, Nucl. Instrum. Methods Phys. Res. B 9, 377 (1985).

241. M. Sasao, A. Matsumoto, A. Nishizawa, K. N. Sato, S. Takagi, S. Amamiya, T. Masuda, Y. Tsurita, Y. Kanamori, Y. Haruyama, and F. Fukuzawa, p. 569 in XIII International Conference on the Physics of Electronic and Atomic Collisions, Berlin, July 27-Aug. 2, 1983.

242. R. W. McCullough, T. V. Goffe, M. B. Shah, M. Lennon, and H. B. Gilbody, J. Phys. B 15, 111 (1982).

243. K. Kadota, D. Dijkkamp, R. L. van der Woude, A. de Boer, P. G. Yan, and F. J. de Heer, J. Phys. B 15, 3275 (1982).

244. G. A. Murray, J. Stone, M. Mayo, and T. J. Morgan, Phys. Rev. A 25, 1805 (1982).

245. K. Kadota, D. Dijkkamp, R. L. van der Woude, P. G. Yan, and F. J. de Heer, Phys. Lett. A 88, 135 (1982).

246. M. Sasao, K. Sato, A. Matsumoto, A. Nishizawa, S. Takagi, S. Amemiya, T. Masuda, Y. Tsurita, F. Fukuzawa, Y. Haruyama, and Y. Kanamori, J. Phys. Soc. Jpn. 55, 102 (1986).

247. M. B. Shah, T. V. Goffe, and H. B. Gilbody, J. Phys. B 11, L233 (1978).

248. W. Seim, A. Muller, I. Wirkner-Bott, and E. Salzborn, J. Phys. B 14, 3475 (1981).

249. I. Wirkner-Bott, W. Seim, A. Muller, P. Kester, and E. Salzborn, J. Phys. B 14, 3987 (1981).

250. S. K. Allison, J. Cuevas, and M. Garcia-Munoz, Phys. Rev. 120, 1266 (1960).

251. I. S. Dmitriev, V. S. Nikolaev, Yu. A. Tashaev, and Ya. Teplova, Sov. Phys.-JETP 40, 1017 (1975).

252. K. E. Banyard and G. W. Shirtcliffe, Phys. Rev. A 30, 604 (1984).

253. A. L. Ford, J. F. Reading, and R. L. Becker, J. Phys. B 15, 3257 (1982).

254. P. H. Hvelplund, J. Phys. B 9, 1555 (1976).

255. F. Aumayr, G. Lakits, W. Husinsky, and H. Winter, J. Phys. B 18, 2493 (1985).

256. F. Aumayr, M. Fehringer, and H. Winter, J. Phys. B 17, 4201 (1984).

257. A. M. Ermolaev, J. Phys. B 17, 1069 (1984).

258. F. Aumayr, M. Fehringer, and H. Winter, J. Phys. B 17, 4185 (1984).

259. J. L. Barrett and J. J. Leventhal, Phys. Rev. A 23, 485 (1981).

260. R. Hippler, W. Harbich, H. Madeheim, K. Kleinpoppen, and H. O. Lutz, Phys. Rev. A 35, 3139 (1987).

261. J. T. Park, Adv. At. Mol. Phys. 19, 67 (1983).

262. P. G. Yan, R. L. van der Woude, D. Dijkkamp, and F. J. de Heer, Phys. Scr. T3, 120 (1983).

263. B. H. Bransden and A. M. Ermolaev, Phys. Lett. 84A, 316 (1981).

264. E. J. Shipsey, L. T. Redmon, J. C. Browne, and R. E. Olson, Phys. Rev. A 18, 1961 (1978).

265. R. Johnsen and M. A. Biondi, Phys. Rev. A 18, 996 (1978).

266. J. N. Bardsley, J. S. Cohen, and J. M. Wadehra, Phys. Rev. A 19, 2129 (1979).

267. F. Aumayr and H. Winter, J. Phys. B 18, L741 (1985).

268. C. J. Anderson, R. J. Girnius, A. M. Howald, and L. W. Anderson, Phys. Rev. A 22, 822 (1980).

269. H. Tawara, Atomic Data Nucl. Data Tables 22, 491 (1978).

270. Ya. M. Fogel, V. A. Ankudinov, D. V. Pilipenko, and N. V. Topolia, Sov. Phys.-JETP 7, 400 (1958).

271. D. M. Davidovic and R. K. Janev, Phys. Rev. 186, 89 (1969).

272. A. Dalgarno and M.R.C. McDowell, Proc. Phys. Soc. A69, 615 (1956).

273. W. Schwab, G. B. Baptista, E. Justiniano, R. Schuch, H. Vogt, and E. W. Weber, J. Phys. B 20, 2825 (1987).

274. B. Peart and R. A. Forrest, J. Phys. B 12, L23 (1979).

275. F. Brouillard, W. Claeys, G. Poulaert, G. Rahmat, and G. Van Wassenhove, J. Phys. B 12, 1253 (1979).

276. K. T. Dolder, "The measurement of inelastic ion-ion and electron-ion collisions" p. 373 in Physics of Ion-Ion and Electron-Ion Collisions, Plenum Press, New York, 1983.

277. B. Peart, K. Rinn, and K. T. Dolder, J. Phys. B 16, 2831 (1983).

278. M. F. Watts, K. F. Dunn, and H. B. Gilbody, J. Phys. B 19, L355 (1986).

279. K. Rinn, F. Melchert, K. Rink, and E. Salzborn, J. Phys. B 19, 3717 (1986).

280. D. Fussen and C. Kubach, J. Phys. B 19, L31 (1986).

281. D. G. Hummer, R. F. Stebbings, W. L. Fite, and L. M. Branscomb, Phys. Rev. 119, 668 (1960).

282. R. D. DuBois, Phys. Rev. A 33, 1595 (1986).

283. F. Melchert, K. Rink, K. Rinn, E. Salzborn, and N. Grun, J. Phys. B 20, L223 (1987).

284. R. J. Allan, J. Phys. B 19, L683 (1986).

285. R. Abrines and I. C. Percival, Proc. Phys. Soc., London 88, 861 (1966).

286. R. J. McNeal, D. C. Clark, and R. A. Klingberg, Phys. Rev. A 2, 131 (1970).

287. H. H. Fleischmann and R. A. Young, Phys. Rev. 178, 254 (1969).

288. R. E. Olson, J. Phys. B 11, L227 (1978).

289. F. Brouillard and D.J.W. Delfosse, p. 159 in V International Conference on the Physics of Electronic and Atomic Collisions, Leningrad, USSR, July 17-23, 1976, Publishing House Nauka, Leningrad, 1967.

290. M. Terao, M. Cherkani, S. Szucs, and F. Brouillard, p. 620 in XIV International Conference on the Physics of Electronic and Atomic Collisions, Palo Alto, July 24-30, 1985.

291. M. Terao, S. Szucs, M. Cherkani, F. Brouillard, and R. J. Allan, Europhys. Lett. 1, 123 (1986).

292. B. Peart, R. Grey, and K. T. Dolder, J. Phys. B 9, L373 (1976).

293. T. D. Gaily and M.F.A. Harrison, J. Phys. B 3, 1098 (1970).

294. B. Peart and M. A. Bennett, J. Phys. B 19, L321 (1986).

295. C. -L. Liao, C. -X. Liao, and C. Y. Ng, J. Chem. Phys. 81, 5672 (1984).

296. R. C. Dehmel, H. K. Chau, and H. H. Fleischmann, Atomic Data 5, 231 (1973).

297. A. B. Wittkower, G. Levy, and H. B. Gilbody, p. 205 in V International Conference on the Physics of Electronic and Atomic Collisions, Leningrad, USSR, July 17-23, 1976, Publishing House Nauka, Leningrad, 1967.

298. A. B. Wittkower, G. Levy, and H. B. Gilbody, Proc. Phys. Soc. 91, 306 (1967).

299. H. Tawara and A. Russek, Rev. Mod. Phys. 45, 178 (1973).

300. N. V. Fedorenko, Sov. Phys. Tech. Phys. 15, 1947 (1971).

301. P. H. Hvelplund and A. Andersen, Phys. Scr. 26, 370 (1982).

302. M. A. Prasad and K. Unnikrishnan, Phys. Rev. A 22, 514 (1980).

303. R. Smythe and J. W. Toevs, Phys. Rev. 139, A15 (1965).

304. N. Noda, J. Phys. Soc. Jpn. 41, 625 (1976).

305. D. V. Pilipenko, and Ya. M. Fogel, Sov. Phys.-JETP 15, 646 (1962).

306. B. Donnally and W. Sawyer, p. 488 in VI International Conference on the Physics of Electronic and Atomic Collisions, MIT, July 28-Aug 2, The MIT Press, Cambridge, Massachusetts, 1969.

307. K. H. Berkner, S. N. Kaplan, and R. V. Pyle, Phys. Rev. 134, A1461 (1964).

308. R. Curran and T. M. Donahue, Phys. Rev. 118, 1233 (1960).

309. H. H. Fleischmann, C. F. Barnett, and J. A. Ray, Phys. Rev. A 10, 569 (1974).

310. J. H. Montague, Phys. Rev. 81, 1026 (1951).

311. F. Roussel, P. Pradel, and G. Spiess, p. 462 in X International Conference on the Physics of Electronic and Atomic Collisions, Paris, July 21-27, 1977, North-Holland Publishing Co., 1977.

312. F. Roussel, P. Pradel, A. S. Schlachter, and G. Spiess, p. 27 in IX International Conference on the Physics Electronic and Atomic Collisions, Seattle, July 24-30, 1975, University of Washington Press, Seattle, WA, 1975.

313. K. A. Smith, M. D. Duncan, M. W. Geis, and R. D. Rundel, J. Geophys. Res. 81, 2231 (1976).

314. K. A. Smith, M. D. Duncan, M. W. Geis, and R. D. Rundel, p. 25 in IX International Conference on the Physics Electronic and Atomic Collisions, Seattle, WA, July 24-30, 1975, University of Washington Press, Seattle, WA, 1975.

315. E. S. Solov'ev, R. N. In'in, V. A. Oparin, and N. V. Fedorenko, Sov. Phys.-JETP 15, 459 (1962).

316. S. K. Allison, Rev. Mod. Phys. 30, 1137 (1958).

317. A. M. Kudryavtsev and A. F. Sorokin, Sov. Phys.-Tech. Phys. 22, 1368 (1977).

318. J. S. Risley, Comments At. Mol. Phys. 12, 215 (1983).

319. N. V. Fedorenko, p. 54 in Lectures Summer School on Physics Ionized Gases, Herceq-Novi, Yugoslavia, 1970.

320. E. H. Pedersen and L. Larsen, J. Phys. B 12, 4099 (1979).

321. G. I. Dimov and V. G. Dudnikov, Sov. Phys.-Tech. Phys. 11, 919 (1967).

322. Y. Nakai, A. Kikuchi, T. Shirai, and M. Sataka, Japan Atomic Energy Research Institute Report JAERI-M 83-143, September (1983).

323. R. J. McNeal, D. C. Clark, and R. A. Klingberg, Phys. Rev. A 2, 131 (1970).

324. R. N. Il'in, V. V. Afrosimov, and N. V. Fedorenko, Sov. Phys.-JETP 9, 29 (1959).

325. F. Schwirzke, Z. Phys. 157, 540 (1960).

326. G. Spiess, A. Valance, and P. Pradel, Phys. Rev. A 6, 746 (1972).

327. H. B. Gilbody, R. Browning, R. M. Reynolds, and G. I. Riddell, p. 1080 in VII International Conference on the Physics of Electronic and Atomic Collisions, Amsterdam, July 26-30, 1971, North-Holland Publishing CO., Amsterdam, 1971.

328. R. H. Hughes and P. O. Berge, Phys. Rev. A 11, 906 (1975).

329. H. B. Gilbody and J. L. Corr, J. Phys. B 7, 1953 (1974).

330. H. B. Gilbody, R. Browning, R. M. Reynolds, and G. I. Riddell, J. Phys. B 4, 94 (1971).

331. V. Dose and R. Gunz, J. Phys. B 5, 636 (1972).

332. A. Cornet, W. Claeys, V. Lorent, J. Jureta, and D. Fussen, J. Phys. B 17, 2643 (1984).

333. J.D.A. McKee, J. Geddes, and H. B. Gilbody, J. Phys. B 12, 1701 (1979).

334. R. H. Hughes and S. S. Choe, Phys. Rev. A 6, 1413 (1972).

335. M. E. Rudd, Y.-K. Kim, D. H. Madison and J. W. Gallagher, Rev. Mod. Phys. 57, 965 (1985).

336. M. B. Shah, D. S. Elliott, and H. B. Gilbody, J. Phys. B 20, 2481 (1987).

337. M. B. Shah and H. B. Gilbody, J. Phys. B 14, 2361 (1981).

338. M. B. Shah and H. B. Gilbody, p. 791 in XII International Conference on the Physics of Electronic and Atomic Collisions, Gatlinburg, TN, July 15-21, 1981.

339. J. T. Park, p. 109 in XII International Conference on the Physics of Electronic and Atomic Collisions, Gatlinburg, TN, July 15-21, 1981. (Invited Papers)

340. J. T. Park, J. E. Aldag, J. M. George, and J. L. Peacher, p. 753 in IX International Conference on the Physics of Electronic and Atomic Collisions, Seattle, July 24-30, 1975, University of Washington Press, Seattle, WA, 1975.

341. J. T. Park, J. E. Aldag, J. M. George, J. L. Peacher, and J. H. McGuire, Phys. Rev. A 15, 508 (1977).

342. H. B. Gilbody, IEEE Trans. Nucl. Sci. NS-30, 1201 (1983).

343. H. B. Gilbody and J. V. Ireland, Proc. R. Soc. London A277, 137 (1963).

344. M. E. Rudd, Phys. Rev. A 28, 3244 (1983).

345. M. E. Rudd, Phys. Rev. A 20, 787 (1979).

346. M. B. Shah and H. B. Gilbody, J. Phys. B 15, 3441 (1982).

347. M. E. Rudd and T. Jorgensen, Jr., Phys. Rev. 131, 666 (1963).

348. V. V. Afrosimov, R. N. Il'in, and N. V. Fedorenko, Sov. Phys.-JETP 7, 968 (1958).

349. L. H. Toburen and W. E. Wilson, Phys. Rev. A 5, 247 (1972).

350. J. W. Hooper, E. W. McDaniel, D. W. Martin, and D. S. Harmer, Phys. Rev. 121, 1123 (1961).

351. C. E. Kuyatt and T. Jorgensen, Jr., Phys. Rev. 130, 1444 (1963).

352. Ia. M. Fogel, L. I. Krupnik, and B. G. Safronov, Sov. Phys.-JETP 1, 415 (1955).

353. L. I. Pivovar and Yu. Z. Levchenko, Sov. Phys.-JETP 25, 27 (1967).

354. H. B. Gilbody and A. R. Lee, Proc. R. Soc. London A 274, 365 (1963).

355. H. B. Gilbody and J. B. Hasted, Proc. R. Soc. London A 246, 382 (1957).

356. V. V. Afrosimov, G. A. Leiko, Yu. A. Mamaev, M. N. Panov, and N. V. Fedorenko, p. 114 in VI International Conference on the Physics of Electronic and Atomic Collisions, MIT, July 28-Aug 2, 1969, The MIT Press, Cambrdige, Massachusetts, 1969.

357. M. E. Rudd, T. V. Goffee, R. D. DuBois, L. H. Toburen, and C. A. Ratcliffe, p. 413 in XIII International Conference on the Physics of Electronic and Atomic Collisions, Berlin, (1983).

358. J. P. Keene, Philos. Mag. 40, 369 (1949).

359. E. W. McDaniel, J. W. Hooper, D. W. Martin, and D. S. Harmer, Vol. 1, p. 60 in V International Conference on Ionization Phenomena in Gases, Munich, August 28-September 1, 1961, North-Holland Publishing Co., Amsterdam, 1962.

360. A. K. Edwards, R. M. Wood and R. L. Ezell, Phys. Rev. A 34, 4411 (1986).

361. B. G. Lindsay, F. B. Yousif, F. R. Simpson, and C. J. Latimer, J. Phys. B 20, 2759 (1987).

362. R. Browning and H. B. Gilbody, J. Phys. B 1, 1149 (1968).

363. Yu. Z. Levchenko, L. I. Pivovar, and G. D. Tolstolutskaya, Sov. Phys.-JETP 37, 1004 (1973).

364. H. B. Gilbody, J. B. Hasted, A. R. Lee, J. V. Ireland, E. W. Thomas, and A. S. Whiteman, Vol. 1, p. 91 in VI International Conference on Ionization Phenomena in Gases, Paris, July 8-13, 1963.

365. A. K. Edwards, R. M. Wood, and R. L. Ezell, Phys. Rev. A 31, 99 (1985).

366. N. V. Fedorenko, V. V. Afrosimov, R. N. Il'in, and E. S. Solov'ev, Vol. 1, p. 47 in IV International Conference on Ionization Phenomena in Gases, Uppsala, 1959.

367. S. Wexler, J. Chem. Phys. 41, 1714 (1964); 44, 2221 (1966).

368. J. W. Hooper, D. S. Harmer, D. W. Martin, and E. W. McDaniel, Phys. Rev. 125, 2000 (1962).

369. M. E. Rudd, C. A. Sautter, and C. L. Bailey, Phys. Rev. 151 20 (1966).

370. M. Becker and A. Scharmann, Z. Naturforsch. A 24, 854 (1969).

371. J. T. Park and F. D. Schowengerdt, Phys. Rev. 185, 152 (1969).

372. L. J. Puckett and D. W. Martin, Phys. Rev. A 1, 1432 (1970).

373. N. Stolterfoht, Z. Phys. 248, 81 (1971).

374. Z. Z. Latypov and A. A. Shaporenko, Sov. Phys-JETP Lett. 18, 258 (1973).

375. P. Hvelplund, H. K. Haugen, and H. Knudsen, Phys. Rev. A 22, 1930 (1980).

376. R. D. DuBois and S. T. Manson, Phys. Rev. A 35, 2007 (1987).

377. D. K. Gibson and I. D. Reid, J. Phys. B 19, 3265 (1986).

378. R. J. McNeal and J. H. Birely, Rev. Geophys. Space Phys. 11, 633 (1973).

379. M. E. Rudd and D. H. Madison, Phys. Rev. A 14, 128 (1976).

380. J. H. Miller, L. H. Toburen, and S. T. Manson, Phys. Rev. A 27, 1337 (1983).

381. H. K. Haugen, L. H. Andersen, P. Hvelplund, and H. Knudsen, Phys. Rev. A 26, 1962 (1982).

382. L. H. Andersen, IEEE Trans. Nucl. Sci. NS-30, 973 (1983).

383. M. B. Shah and H. B. Gilbody, J. Phys. B 18, 899 (1985).

384. H. Knudsen, L. H. Andersen, P. Hvelplund, G. Astner, H. Cederquist, H. Danared, L. Liljeby, and K.-G. Rensfelt, J. Phys. B 17, 3545 (1984).

385. D. W. Martin, L. J. Puckett, and G. O. Taylor, p. 207 in V International Conference on the Physics of Electronic and Atomic Collisions, Leningrad, USSR, July 17-23, 1967, Publishing House Nauka, Leningrad, 1967.

386. L. H. Andersen, P. Hvelplund, H. Knudsen, S. P. Moller, K. Elsener, K.-G. Rensfelt, and E. Uggerhoj, Phys. Rev. Lett. 57, 2147 (1986).

387. L. H. Andersen, P. Hvelplund, H. Knudsen, S. P. Moller, A. H. Sorensen, K. Elsener, K.-G. Rensfelt, and E. Uggerhoj, Phys. Rev. A 36, 3612 (1987).

388. R. D. DuBois, L. H. Toburen, and M. E. Rudd, Phys. Rev. A 29, 70 (1984).

389. W. Schon, S. Krudener, F. Melchert, K. Rinn, M. Wagner, and E. Salzborn, J. Phys. B 20, L759 (1987).

390. S. W. Nagy, W. J. Savola, Jr. and E. Pollack, Phys. Rev. 177, 71 (1969).

391. A. S. Dickinson and D.J.W. Hardie, J. Phys. B 12, 4147 (1979).

392. A. Jognaux, F. Brouillard, and S. Szucs, J. Phys. B 11, L669 (1978).

393. B. Peart and K. Dolder, J. Phys. B 12, 4155 (1979).

394. R. E. Olson, J. R. Peterson, and J. T. Mosely, J. Chem. Phys. 53, 3391 (1970).

395. M. W. Gealy and B. Van Zyl, Phys. Rev. A 36, 3091 (1987).

396. M. W. Gealy and B. Van Zyl, Phys. Rev. A 36, 3100 (1987).

397. W. H. Cramer and J. H. Simons, J. Chem. Phys. 26, 1272 (1957).

398. E. H. Pedersen and L. Larsen, J. Phys. B 12, 4085 (1979).

399. E. H. Pedersen and P. Hvelplund, J. Phys. B 7, 132 (1974).

400. P. Hvelplund and E. H. Pedersen, Phys. Rev. A 9, 2434 (1974).

401. S. K. Allison, Phys. Rev. 110, 670 (1958).

402. F. Bastien, E. Depuydt, and A. Sinturei, C. R. Acad. Sci. Paris B 269 459 (1969).

403. D. W. Martin, L. J. Puckett, and G. O. Taylor, p. 280 in IV International Conference on the Physics of Electronic and Atomic Collisions, Quebec, Canada, August 2-6, 1965, Science Bookcrafter, Inc., New York, 1965.

404. E. H. Pedersen and P. Hvelplund, p. 522 in VII International Conference on the Physics of Electronic and Atomic Collisions, Amsterdam, July 26-30, 1971, North-Holland Publishing Co., Amsterdam, 1971.

405. A. Itoh, K. Ohnishi, and F. Fukuzawa, J. Phys. Soc. Jpn. 49, 1513 (1980).

406. K. Tobita and H. Takeuchi, J. Phys. Soc. Jpn. 55, 4231 (1986).

407. Ya. M. Fogel, V. A. Ankudinov, D. V. Pilipenko, Sov. Phys.-JETP 11, 18 (1960).

408. H. B. Gilbody, K. F. Dunn, R. Browning, and C. J. Latimer, J. Phys. B 3, 1105 (1970).

409. K. Okuno, p. 26 in Nagoya University Report IPPJ-AM-9, December, (1978).

410. E. H. Pedersen, J. Heinemeier, L. Larsen, and J. V. Mikkelsen, J. Phys. B 13, 1167 (1980).

411. R. E. Miers and L. W. Anderson, Phys. Rev. A 1, 819 (1970).

412. E. E. Benton, E. E. Ferguson, F. A. Matsen, and W. W. Robertson, Phys. Rev. 128, 206 (1962).

413. E. H. Pedersen, Phys. Rev. A 15, 53 (1977).

414. E. H. Pedersen, Phys. Rev. A 10, 110 (1974).

415. R. W. McCullough, T. V. Goffe, and H. B. Gilbody, J. Phys. B 11, 2333 (1978).

416. E. H. Pedersen, Phys. Rev. Lett. 42, 440 (1979).

417. X. Urbain, A. Giusti-Suzor, D. Fussen, and C. Kubach, J. Phys. B 19, L273 (1986).

418. G. Poulaert, F. Brouillard, W. Claeys, J. W. McGowan, and G. Van Wassenhove, J. Phys. B 11, L671 (1978).

419. G. Poulaert, F. Brouillard, W. Claeys, P. DeFrance, and J. W. McGowan, p. 876 in XI International Conference on the Physics of Electronic and Atomic Collisions, Kyoto, August 29-September 4, 1979.

420. M. J. Shaw, R. C. Bolden, R. S. Hemsworth, and N. D. Twiddy, Chem. Phys. Lett. 8, 148 (1971).

421. J. S. Howard, J. P. Riola, R. D. Rundel, and R. F. Stebbings, J. Phys. B 6, L109 (1973).

422. J. Fort, J. J. Laucagne, A. Pesnelle, and G. Watel, Phys. Rev. A 14, 658 (1976).

423. J. Fort, J. J. Laucagne, A. Pesnelle, and G. Watel, Chem. Phys. Lett. 37, 60 (1976).

424. J. Fort, J. J. Laucagne, A. Pesnelle, and G. Watel, p. 302 in X International Conference on the Physics of Electronic and Atomic Collisions, Paris, July 21-27, 1977, North-Holland Publishing Co., Amsterdam, 1977.

425. G. D. Magnuson and R. H. Neynaber, J. Chem. Phys. 60, 3385 (1974).

426. C. Manus, A. Pesnelle, and G. Watel, p. 525 in X International Conference on the Physics of Electronic and Atomic Collisions, Paris, July 21-27, 1977, North-Holland Publishing Co., Amsterdam, 1977.

427. H. Morgner and A. Niehaus, J. Phys. B 12, 1805 (1979).

428. R. H. Neynaber, p. 287 in XI International Conference on the Physics of Electronic and Atomic Collisions, Kyoto, August 29-September 4, 1979.

429. R. H. Neynaber and G. D. Magnuson, J. Chem. Phys. 62, 4953 (1975).

430. R. H. Neynaber and S. Y. Tang, J. Chem. Phys. 69, 4851 (1978).

431. J. Fort, J. J. Laucagne, A. Pesnelle, and G. Watel, Phys. Rev. A 18, 2063 (1978).

432. A. P. Hickman and H. Morgner, J. Chem. Phys. 67, 5484 (1977).

433. C. Manus, Physica 82C, 165 (1976).

434. R. C. Bolden, R. S. Hemsworth, M. J. Shaw, and N. D. Twiddy, J. Phys. B 3, 61 (1970).

435. J. S. Cohen and R. K. Preston, p. 310 in X International Conference on the Physics of Electronic and Atomic Collisions, Paris, July 21-27, 1977, North-Holland Publishing Co., Amsterdam, 1977.

436. F. B. Dunning and A.C.H. Smith, J. Phys. B 3, L60 (1970).

437. J. Fort, T. Bolzinger, D. Corno, T. Ebding, and A. Pesnelle, Phys. Rev. A 18, 2075 (1978).

438. J. Fort, T. Bolzinger, D. Corno, and T. Ebding, p. 314 in X International Conference on the Physics of Electronic and Atomic Collisions, Paris, July 21-27, 1977, North-Holland Publishing Co., Amsterdam, 1977.

439. H. Hotop and A. Niehaus, Z. Phys. 215, 395 (1968).

440. P. A. Jerram and A.C.H. Smith, J. Phys. B 18, 1747 (1985).

441. C. R. Jones and W. W. Robertson, J. Chem. Phys. 49, 4241 (1968).

442. D. W. Martin and P. E. Siska, J. Chem. Phys. 82, 2630 (1985).

443. A. Munzer and A. Niehaus, J. Electron Spectrosc. Relat. Phenom. 23, 367 (1981).

444. R. H. Neynaber and G. D. Magnuson, J. Chem. Phys. 61, 749 (1974).

445. J. R. Penton and E. E. Muschlitz, Jr., J. Chem. Phys. 49, 5083 (1968).

446. A. Pesnelle and S. Runge, p. 559 in XIII International Conference on the Physics of Electronic and Atomic Collisions, Berlin, July 27-August 2, 1983.

447. P. Pradel and J. J. Laucagne, J. Phys. (Paris) 44, 1263 (1983).

448. J. P. Riola, J. S. Howard, R. D. Rundel, and R. F. Stebbings, J. Phys. B 7, 376 (1974).

449. T. Sinda and A. Pesnelle, p. 1107 in VII International Conference on the Physics of Electronic and Atomic Collisions, Amsterdam, July 26-30, 1971, North-Holland Publishing Co., Amsterdam, 1971.

450. L. T. Specht, K. D. Foster, and E. E. Muschlitz, Jr., Chem. Phys. 63, 1582 (1975).

451. G. E. Veatch and H. J. Oskam, Phys. Rev. A 8, 389 (1973).

452. W. P. West, T. B. Cook, F. B. Dunning, R. D. Rundel, and R. F. Stebbings, J. Chem. Phys. 63, 1237 (1975).

453. A. L. Schmeltekopf and F. C. Fehsenfeld, J. Chem. Phys. 53, 3173 (1970).

454. W. Lindinger, A. L. Schmeltekopf, and F. C. Fehsenfeld, J. Chem. Phys. 61, 2890 (1974).

455. W. P. Sholette and E. E. Muschlitz, Jr., J. Chem. Phys. 36, 3368 (1962).

456. R. H. Neynaber, G. D. Magnuson, and J. K. Layton, J. Chem. Phys. 57, 5128 (1972).

457. F. Melchert, K. Rink, K. Rinn, and E. Salzborn, J. Phys. B 20, L797 (1987).

458. H. B. Gilbody, p. 223 in XII International Conference on the Physics of Electronic and Atomic Collisions, Gatlinburg, TN, July 15-21, 1981. (Invited Paper)

459. R. Browning, C. J. Latimer, and H. B. Gilbody, J. Phys. B 3, 667 (1970).

460. N. G. Utterback, Phys. Rev. Lett. 12, 295 (1964).

461. V. A. Gusev, G. N. Polyakova, V. F. Erko, Ya. M. Fogel, and A. V. Zats, p. 809 in VI International Conference on the Physics of Electronic and Atomic Collisions, MIT, July 28-August 2, 1969, The MIT Press, Cambridge, Massachusetts, 1969.

462. E. S. Solov'ev, R. N. Il'in, V. A. Oparin, and N. V. Fedorenko, p. 692 in III International Conference on the Physics of Electronic and Atomic Collisions, London, July 22-26, 1963, North-Holland Publishing Co., Amsterdam, 1964.

463. E. S. Solov'ev, R. N. Il'in, V. A. Oparin, and N. V. Fedorenko, Sov. Phys.-JETP 18, 342 (1964).

464. G. N. Polyakova, V. A. Gusev, V. F. Erko, Ya. M. Fogel, and A. V. Zats, Sov. Phys.-JETP 31, 637 (1970).

465. S. Krasner, Phys. Rev. 99, 520 (1955).

466. A. B. Wittkower, G. Levy, and H. B. Gilbody, Proc. Phys. Soc. 91, 862 (1967).

467. H. Tawara, J. Phys. Soc. Jpn. 31, 236 (1971).

468. W. L. Williams and F. M. Goldberg, p. 1087 in VII International Conference on the Physics of Electronic and Atomic Collisions, Amsterdam, July 26-30, 1971, North-Holland Publishing Co., Amsterdam, 1971.

469. A. B. Wittkower, G. Levy, and H. B. Gilbody, p. 202 in V International Conference on the Physics of Electronic and Atomic Collisions, Leningrad, USSR, July 17-23, 1967, Publishing House Nauka, Leningrad, 1967.

470. Y. Kanamori, Y. Haruyama, T. Kido, and F. Fukuzawa, Phys. Lett. A 101, 391 (1984).

471. I. S. Dmitriev, N. F. Vorob'ev, Zh. M. Konovalova, V. S. Nikolaev, V. N. Novozhilova, Ya. A. Teplova, and Yu. A. Fainberg, Sov. Phys.-JETP 57, 1157 (1983).

472. V. S. Nikolaev, V. S. Senashenko, V. A. Sidorovich, and V. Yu. Shafer, Sov. Phys.-Tech. Phys. 23, 789 (1978).

473. N. G. Utterback, p. 661 in III International Conference on the Physics of Electronic and Atomic Collisions, London, July 22-26, 1963, North-Holland Publishing Co., Amsterdam, 1964.

474. H. C. Hayden and N. G. Utterback, Phys. Rev. 135, A1575 (1964).

475. H. Fetz and H. Bussmann, p. 222 in V International Conference on the Physics of Electronic and Atomic Collisions, Leningrad, USSR, July 17-23, 1967, Publishing House Nauka, Leningrad, 1967.

476. M. B. Shah, T. V. Goffe, and H. B. Gilbody, J. Phys. B 10, L723 (1977).

477. M. B. Shah and H. B. Gilbody, J. Phys. B 8, 372 (1975).

478. D. W. Martin, R. A. Langley, D. S. Harmer, J. W. Hooper, and E. W. McDaniel, Phys. Rev. 136, A385 (1964).

479. R. A. Langley, D. W. Martin, D. S. Harmer, J. W. Hooper, and E. W. McDaniel, Phys. Rev. 136, A379 (1964).

480. L. I. Pivovar, Yu. Z. Levchenko, and A. N. Grigor'ev, Sov. Phys.-JETP 27, 699 (1968).

481. R. Browning, C. J. Latimer, and H. B. Gilbody, J. Phys. B 2, 534 (1969).

482. A. K. Edwards, R. M. Wood, and R. L. Ezell, Phys. Rev. A 31, 3972 (1985).

483. P. R. Jones, E. P. Ziemba, H. A. Moses, and E. Everhart, Phys. Rev. 113, 182 (1959).

484. N. V. Fedorenko, V. V. afrosimov, and D. M. Kaminker, Sov. Phys-Tech. Phys. 1, 1861 (1956).

485. N. V. Fedorenko and V. V. Afrosimov, Sov. Phys.-Tech. Phys. 1, 1872 (1956).

486. A. R. Lee and H. B. Gilbody, p. 123 in VI International Conference on Ionization Phenomena in Gases, Paris, July 8-13, 1963.

487. H. Fetz, M. Saar, and H. Scheidt, p. 1143 in VII International Conference on the Physics of Electronic and Atomic Collisions, Amsterdam, July 26-30, 1971, North-Holland Publishing Co., Amsterdam, 1971.

488. H. Knudsen, L. H. Andersen, H. K. Haugen, and P. Hvelplund, Phys. Scr. 26, 132 (1982).

489. I. S. Dmitriev, V. S. Nikolaev, L. M. Fateeva, and Ya. A. Teplova, Sov. Phys.-JETP 15, 11 (1962).

490. A. R. Lee and H. B. Gilbody, p. 672 in III International Conference on the Physics of Electronic and Atomic Collisions, London, July 22-26, 1963, North-Holland Publishing Co., Amsterdam, 1964.

491. K. F. Man, W. Steckelmacher, and M. W. Lucas, J. Phys. B 19, 4171 (1986).

492. N. V. de Castro Faria, F. L. Freire, Jr., and A. G. de Pinho, Phys. Rev. A 37, 280 (1988).

493. E. Snitzer, Phys. Rev. 89, 1237 (1953).

494. R. M. Wood, A. K. Edwards, and R. L. Ezell, Phys. Rev. A 34, 4415 (1986).

495. S. H. Be, T. Tonuma, H. Kumagai, H. Shibata, M. Kase, T. Kambara, I. Kohno, and H. Tawara, J. Phys. B 19, 1771 (1986).

496. A. K. Edwards, R. M. Wood, and R. L. Ezell, Phys. Rev. A 32, 1346 (1985).

497. D. W. Martin, R. A. Langley, J. W. Hooper, D. S. Harmer, and E. W. McDaniel, p. 679 in III International Conference on the Physics of Electronic and Atomic Collisions, London, July 22-26, 1963, North-Holland Publishing CO., Amsterdam, 1964.

498. B. I. Kikiani, G. M. Mirianashvili, Z. E. Saliya, and I. G. Bagdasarova, Sov. Phys.-Tech. Phys. 21, 634 (1976).

499. N. V. Fedorenko, Sov. Phys.-Usp. 2, 526 (1959).

500. R. H. Hughes, H. M. Petefish, and H. Kisner, Phys. Rev. A 5, 2103 (1972).

501. R. H. Neynaber and S. Y. Tang, Chem. Phys. 69, 213 (1982).

502. R. H. Neynaber and S. Y. Tang, p. 613 in XII International Conference on the Physics of Electronic and Atomic Collisions, Gatlinburg, TN, July 15-21, 1981.

503. D. Rapp and W. E. Francis, J. Chem. Phys. 37, 2631 (1962).

504. P. Gangopadhyay and J. N. Bardsley, Chem. Phys. Lett. 96, 379 (1983).

505. D. P. Hodgkinson and J. S. Briggs, J. Phys. B 9, 255 (1976).

506. R. K. Janev and L. P. Presnyakov, J. Phys. B 13, 4233 (1980).

507. D.S.F. Crothers and J. F. McCann, J. Phys. B 16, 3229 (1983).

508. R. E. Olson and A. Salop, Phys. Rev. A 16, 531 (1977).

509. D.J.W. Hardie and R. E. Olson, J. Phys. B 16, 1983 (1983).

510. M. B. Shah and H. B. Gilbody, J. Phys. B 14, 2361 (1981).

511. J. E. Golden and J. H. McGuire, J. Phys. B 9, L11 (1976).

512. C. R. Garibotti and J. E. Miraglia, Phys. Rev. A 25, 1440 (1982).

513. A. C. Whittier, Can J. Phys. 32, 275 (1954).

514. E. E. Muschlitz, Jr., T. L. Bailey, and J. H. Simons, J. Chem. Phys. 24, 1202 (1956).

515. J. B. Hasted and R. A. Smith, Proc. Roy. Soc. London Sect. A 235, 349 (1956).

516. E. E. Muschlitz, Jr., T. L. Bailey, and J. H. Simons, J. Chem. Phys. 26, 711 (1957).

517. P. H. Rose, R. J. Connor, and R. P. Bastide, Bull. Am. Phys. Soc. II 3, 40 (1958).

518. J. F. Williams, Phys. Rev. 154, 9 (1957).

519. F. R. Simpson and H. B. Gilbody, J. Phys. B 5, 1959 (1972).

520. J. S. Risley and R. Geballe, Phys. Rev. A 9, 2485 (1974).

521. J. Heinemeier, P. Hvelplund, and F. R. Simpson, J. Phys. B 9, 2669 (1976).

522. J. S. Risley, p. 619 in XI International Conference on the Physics of Electronic and Atomic Collisions, Kyoto, August 29-September 4, 1979, North-Holland Publishing Co., Amsterdam, 1980. (Invited Paper)

523. W. J. Lichtenberg, K. Bethge, and H. Schmidt-Bocking, J. Phys. B 13, 343 (1980).

524. J. Geddes, J. Hill, M. B. Shah, T. V. Goffe, and H. B. Gilbody, J. Phys. B 13, 319 (1980).

525. H. B. Gilbody, Proc. US-Mexico Joint Seminar on the Atomic Physics of Negative Ions, April 1-4, 1981 (Mexico), Notas de Fisica 5, 146 (1982).

526. Y. Nakai, T. Shirai, T. Tabata, and R. Ito, At. Data Nucl. Data Tables 37, 69 (1987).

527. L. I. Pivovar, V. M. Tubaev, and M. T. Novikov, Sov. Phys.-JETP 13, 23 (1961).

528. J. Guidini, C. R. Acad. Sci. Paris 253, 829 (1961).

529. N. N. Tunitskii, E. S. Zhurkin, and M. V. Tikhomirov, Sov. Phys.-JETP Lett. 12, 210 (1970).

530. M. Vogler and B. Meierjohann, p. 711 in IX International Conference on the Physics of Electronic and Atomic Collisions, Seattle, July 24-30, 1975, University of Washington Press, Seattle, WA, 1975.

531. R. N. Il'in, B. I. Kikiani, V. A. Oparin, E. S. Solov'ev, and N. V. Fedorenko, Sov. Phys.-JETP 19, 817 (1964).

532. A. Itoh, M. E. Rudd, and T. V. Goffe, p.406 in XIV International Conference on the Physics of Electronic and Atomic Collisions, Palo Alto, July 24-30, 1985.

533. W. G. Graham, C. J. Latimer, R. Browning, and H. B. Gilbody, J. Phys. B 7, L405 (1974).

534. P. S. Rudolph and C. E. Melton, J. Chem. Phys. 45, 2227 (1966).

535. K. H. Berkner, T. J. Morgan, R. V. Pyle, and J. W. Stearns, Phys. Rev. A 8, 2870 (1973).

536. I. D. Williams, J. Geddes, and H. B. Gilbody, J. Phys. B 17, 811 (1984).

537. J. H. McGuire, E. Salzborn, and A. Muller, Phys. Rev. A 35, 3265 (1987).

538. R. D. DuBois, Phys. Rev. A 36, 2585 (1987).

539. H. Shibata, S. H. Be, T. Tonuma, H. Kumagai, M. Kase, T. Kombara, I. Kohno, and H. Tawara, p. 407 in XIV International Conference on the Physics of Electronic and Atomic Collisions, Palo Alto, July 24-30, 1985.

540. Z. Z. Latypov, I. P. Flaks, and A. A. Shaporenko, Sov. Phys.-JETP 30, 1076 (1970).

541. M. S. Huq, L. D. Doverspike, and R. L. Champion, p. 401 in XIII International Conference on the Physics of Electronic and Atomic Collisions, Berlin, July 27-August 2, 1983.

542. M. S. Huq, L. D. Doverspike and R. L. Champion, Phys. Rev. A 27, 2831 (1983).

543. T. L. Bailey, C. J. May, and E. E. Muschlitz, Jr., J. Chem. Phys. 26, 1446 (1957).

544. J. S. Risley, Phys. Rev. A 10, 731 (1974).

545. R. L. Champion, L. D. Doverspike, and S. K. Lam, Phys. Rev. A 13, 617 (1976).

546. J. P. Gauyacq and V. A. Esaulov, p. 655 in XI International Conference on the Physics of Electronic and Atomic Collisions, Kyoto, August 29-September 4, 1979, North-Holland publishing Co., Amsterdam, 1980. (Invited papers)

547. C. J. Anderson, R. J. Girnius, A. M. Howald, and L. W. Anderson, Phys. Rev. A 22, 822 (1980).

548. M. S. Huq, L. D. Doverspike, R. L. Champion, and V. A. Esaulov, J. Phys. B 15, 951 (1982).

549. J. S. Risley and R. E. Olson, US-Mexico Joint Seminar on the Atomic Physics of Negative Ions, April 1-4, 1981 (Mexico). Notas de Fisica 5, 108 (1982).

550. L. D. Doverspike and R. L. Champion, US-Mexico Joint Seminar on the Atomic Physics of Negative Ions, April 1-4, 1981 (Mexico). Notas de Fisica 5, 1 (1982).

551. N. Andersen, T. Ansersen, L. Jepsen, and J. Macek, J. Phys. B 17, 2281 (1984).

552. D. P. Almeida, N. V. de Castro Faria, F. L. Freire, Jr., E. C. Montenegro, and A. G. de Pinho, Phys. Rev. A 36, 16 (1987).

553. J. B. Hasted. Proc. Roy. Soc. London A 212, 235 (1952).

554. J. Geddes, J. Hill, M. B. Shah, T. V. Goffe, and H. B. Gilbody, p. 610 in XI International Conference on the Physics of Electronic and Atomic Collisions, Kyoto, August 29-September 4, 1979.

555. V. A. Esaulov, p. 608 in XI International Conference on the Physics of Electronic and Atomic Collisions, Kyoto, August 29-September 4, 1979.

556. Ya. M. Fogel, V. A. Ankudinov, and R. E. Slabospitskii, Sov. Phys.-JETP 5, 382 (1957).

557. C. F. Barnett and J. A. Ray, Atomic Collisions Processes, p. 743, ed. M.R.C. McDowell, North Holland Publ. Co., Amsterdam, 1964.

558. K. H. Berkner, S. N. Kaplan, R. V. Pyle, and J. W. Stearns, Phys. Rev. 146, 91 (1966).

559. S. E. Kupriyanov, N. N. Tunitskii, and A. A. Prov, Sov. Phys.-Tech. Phys. 8, 932 (1964).

560. E. S. Zhurkin, M. V. Tikhomirov, and N. N. Tunitskii, Sov. Phys.-Tech. Phys. 16, 495 (1971).

561. J. S. Risley and P. D. Tennyson, p. 614 in XI International Conference on the Physics of Electronic and Atomic Collisions, Kyoto, August 29-September 4, 1979, 1979.

562. Y. Suzuki, T. Kaneko, M. Tomita, and M. Sakisaka, J. Phys. Soc. Jpn. 55, 3037 (1986).

563. G. Tisone and L. M. Branscomb, Bull. Am. Phys. Soc. 9, 535 (1964).

564. G. H. Gillespie, Phys. Lett. 70A, 329 (1979).

565. T.J.M. Boyd, B. L. Moiseiwitsch, and A. L. Stewart, Proc. Phys. Soc. London A70, 110 (1957).

566. S. L. Willis, G. Peach, M.R.C. McDowell, and J. Banerji, J. Phys. B 18, 3939 (1985).

567. Ya. M. Fogel, Sov. Phys.-JETP 5, 499 (1957).

568. V. A. Esaulov, p. 175 in XIV International Conference on the Physics of Electronic and Atomic Collisions, Palo Alto, July 24-30, 1986.

569. B. Peart, R. Grey, and K. T. Dolder, J. Phys. B 9, 3047 (1976).

570. E. P. Kanter, P. J. Cooney, D. S. Gemmell, K. O. Groeneveld, W. J. Pietsch, A. J. Ratkowski, Z. Vager, and B. J. Zabransky, Phys. Rev. A 20, 834 (1979).

571. M. Vogler and B. Meierjohann, Z. Phys. A 283, 11 (1977).

572. D. Fussen and W. Claeys, J. Phys. B 17, L89 (1984).

573. A. M. Ermolaev, J. Phys. B 21, 81 (1988).

574. G. Lange, B. Huber, and K. Wiesemann, Z. Phys. A 281, 21 (1977).

575. P. M. Windham, P. J. Joseph, and J. A. Weinman, Phys. Rev. 109, 1193 (1958).

576. D. R. Sweetman, Proc. Roy. Soc. London 76, 998 (1960).

577. G. Ryding, A. B. Wittkower, and P. H. Rose, Phys. Rev. 174, 149 (1968).

578. D. J. Nicholas, C. W. Trowbridge, and W. D. Allen, Phys. Rev. 167, 38 (1968).

579. R. D. BuBois and S. T. Manson, Phys. Rev. Lett. 57, 1130 (1986).

580. E. H. Pedersen and P. Hvelplund, J. Phys. B 6, 2600 (1975).

581. J. Heinemeier, P. Hvelplund, and F. R. Simpson, J. Phys. B 8, 1880 (1975).

582. M. J. Coggiola, US-Mexico Joint Seminar on the Atomic Physics of Negative Ions, April 1-4, 1981 (Mexico). Notas de Fisica 5, 78 (1982).

583. F. Rahman and B. Hird, Atomic Data Nucl Data Tables 35, 123 (1986).

584. D. P. Almeida, N. V. de Castro Faria, F. L. Freire, Jr., E. C. Montenegro, and A. G. de Pinho, Nucl. Instrum. Methods Phys. Res. B 24-25, 228 (1987).

585. B. A. Huber, U. Schulz, and K. Wiesemann, Phys. Lett. 79A, 58 (1980).

586. J. T. Park, Advances in Atomic and Molecular Physics Vol. 19, p. 67, ed. D. R. Bates and B. Bederson, Academic Press, New York, 1983.

587. Yu. F. Bydin, Sov. Phys.-JETP 23, 23 (1966).

588. R. W. McCullough, M. B. Shah, M. Lennon, and H. B. Gilbody, J. Phys. B 15, 791 (1982).

589. C. W. Eaker and J. L. Muzyka, Chem. Phys. Lett. 119, 169 (1985).

590. T. F. Moran and J. R. Roberts, J. Chem. Phys. 49, 3411 (1968).

591. E. S. Zhurkin, V. A. Kaminskii, M. V. Tikhomirov, and N. N. Tunitskii, Z. Tech. Fiz. USSR 43, 405 (1973).

592. J. Hill, J. Geddes, and H. B. Gilbody, J. Phys. B 12, 2875 (1979).

593. T. J. Morgan, J. Geddes, and H. B. Gilbody, J. Phys. B 7, 142 (1974).

594. M. r. Flannery, Phys. Rev. 183, 241 (1969).

595. B. M. McLaughlin and K. L. Bell, J. Phys. B 16, 3797 (1983).

596. R. Shingal, B. H. Bransden, and D. R. Flower, J. Phys. B 20, L477 (1987).

597. I. Khurana, R. Srivastava, and A. N. Tripathi, Can. J. Phys. 64, 826 (1986).

598. B. M. McLaughlin and K. L. Bell, J. Phys. B 20, L217 (1987).

599. G. W. McClure, Phys. Rev. 153, 182 (1967).

600. R. H. Hughes and R. F. Kouri, J. Chem. Phys. 57, 4503 (1972).

601. B. Van Zyl, M. W. Gealy, and H. Neumann, Phys. Rev. A 28, 176 (1983).

602. W. B. Maier,III, J. Chem. Phys. 54, 2732 (1971).

603. M. G. Holliday, J. T. Muckerman, and L. Friedman, J. Chem. Phys. 54, 1058 (1971).

604. G. Ochs and E. Teloy, J. Chem. Phys. 61, 4930 (1974).

605. Ch. Schlier, U. Nowotny, and E. Teloy, Chem. Phys. 111, 401 (1987).

606. F. C. Fehsenfeld, D. L. Albritton, Y. A. Bush, P. G. Fournier, T. R. Govers, and J. Fournier, J. Chem. Phys. 61, 2150 (1974).

607. C. H. Douglass, D. J. McClure, and W. R. Gentry, J. Chem. Phys. 67, 4931 (1977).

608. C. H. Douglass, D. J. Mcclure, D. J. Trevor, and W. R. Gentry, p. 584 in IX International Conference on the Physics of Electronic and Atomic Collisions, Seattle, July 24-30, 1973, University of Washington Press, Seattle, WA, 1975.

609. R. D. Smith, D. L. Smith, and J. H. Futrell, Int. J. Mass Spectrom. Ion Phys. 19, 369 (1976).

610. J. W. Stearns, K. H. Berkner, R. V. Pyle, B. P. Briegleb, and M. L. Warren, Phys. Rev. A 4, 1960 (1971).

611. T. Turner, O. Dutuit, and Y. T. Lee, J. Chem. Phys. 81, 3475 (1984).

612. C. F. Giese and W. B. Maier, III, J. Chem. Phys. 39, 739 (1963).

613. L. P. Theard and W. T. Hunterss, Jr., J. Chem. Phys. 60, 2840 (1974).

614. J. A. Rutherford and D. A. Vroom, J. Chem. Phys. 58, 4076 (1973).

615. H. von Koch and L. Friedman, J. Chem. Phys. 38, 1115 (1963).

616. W. A. Chupka and M. E. Russell, J. Chem. Phys. 49, 5426 (1968).

617. W. A. Chupka, J. Berkowitz, and M. E. Russell, p. 71 in VI International Conference on the Physics of Electronic and Atomic Collisions, MIT, July 28-August 2, 1969, The MIT Press, Cambridge, Massachusetts, 1969.

618. R. H. Neynaber and G. D. Magnuson, J. Chem. Phys. 59, 825 (1973).

619. T. R. Govers and P. M. Guyon, Chem. Phys. 113, 425 (1987).

620. J. Krenos and R. Wolfgang, J. Chem. Phys. 52, 5961 (1970).

621. W. R. Gentry, D. J. McClure, and C. H. Douglass, Rev. Sci. Instrum. 46, 367 (1975).

622. M. Saporoschenko, J. Chem. Phys. 42, 2760 (1965).

623. A. G. Harrison, A. Ivko, and T. W. Shannon, Can. J. Chem. 44, 1351 (1966).

624. L. T. Specht, K. D. Foster, and E. E. Muschlitz, Jr. J. Chem. Phys. 63, 1582 (1975).

625. R. P. Clow and J. H. Futrell, Int. J. Mass Spectrom. Ion Phys. 8, 119 (1972).

626. M. T. Bowers, D. D. Elleman, and J. King, Jr., J. Chem. Phys. 50, 4787 (1969).

627. B. G. Reuben and L. Friedman, J. Chem. Phys. 37, 1636 (1962).

628. D. W. Vance and T. L. Bailey, J. Chem. Phys. 44, 486 (1966).

629. W. A. Chupka, M. E. Russell, and K. Refaey, J. Chem. Phys. 48, 1518 (1968).

630. T. Terao and R. A. Back, J. Phys. Chem. 73, 3884 (1969).

631. H. J. Drewitz, Int. J. Mass Spectrom. Ion Phys. 19, 313 (1976).

632. R. H. Neynaber and S. M. Trujillo, Phys. Rev. 167, 63 (1968).

633. I. Koyano and K. Tanaka, J. Chem. Phys. 72, 4858 (1980).

634. S. L. Anderson, F. A. Houle, D. Gerlich, and Y. T. Lee, J. Chem. Phys. 75, 2153 (1981).

635. W. R. Gentry and G. Ringer, J. Chem. Phys. 67, 5396 (1977).

636. T. F. Moran and J. R. Roberts, J. Chem. Phys. 49, 3411 (1968).

637. P. Warneck, J. Chem. Phys. 46, 502 (1967).

638. A. B. Lees and P. K. Rol, J. Chem. Phys. 61, 4444 (1974).

639. J. H. Birely and R. J. McNeal, Phys. Rev. A 5, 257 (1972).

640. J. Grosser and W. Kruger, p. 337 in XIII International Conference on the Physics of Electronic and Atomic Collisions, Berlin, July 27-August 2, 1983.

641. J. Grosser and W. Kruger, Z. Phys. A 318, 25 (1984).

642. R. H. Hughes and S.-S. Choe, Phys. Rev. A 5, 1758 (1972).

643. R. H. Hughes and S.-S. Choe, Phys. Rev. A 5, 656 (1972).

644. R. Krotkov and J. Stone, Phys. Rev. A 22, 473 (1980).

645. A. L. Orbeli, E. P. Andreev, V. A. Ankudinov, and V. M. Dukelskii, Sov. Phys.-JETP 30, 63 (1970).

646. I. Sauers and E. W. Thomas, Phys. Rev. A 10, 822 (1974).

647. L. D. Doverspike and R. L. Champion, J. Chem. Phys. 46, 4718 (1967).

648. V. A. Ankudinov, E. P. Andreev, and A. L. Orbeli, p. 312 in V International Conference on the Physics of Electronic and Atomic Collisions, Leningrad, USSR, July 17-23, 1967, Publishing House Nauka, Leningrad, 1967.

649. D. Hasselkamp, A. Scharmann, and K. H. Schartner, p. 284 in X International Conference on the Physics of Electronic and Atomic Collisions, Paris, July 21-27, 1977, North-Holland Publishing Co., Amsterdam, 1977.

650. C. Benoit and J. P. Gauyacq, J. Phys. B 9, L391 (1976).

651. V. Dose, R. Gunz, and V. Meyer, Helv. Phys. Acta 41, 269 (1968).

652. J. R. Krenos, K. K. Lehmann, J. C. Tully, P. M. Hierl, and G. P. Smith, Chem. Phys. 16, 109 (1976).

653. M. B. Shah, D. S. Elliott, P. McCallion, and H. B. Gilbody, J. Phys. B 21, 2455 (1988).

654. M. E. Rudd, J. Quant. Spectrosc. Radiat. Transfer 31, 387 (1984).

655. J. L. Edwards and E. W. Thomas, Phys. Rev. 165, 16 (1968).

656. K. A. Quickert and D. J. Le Roy, J. Chem. Phys. 53, 1325 (1970).

657. F. C. Fehsenfeld, A. L. Schmeltekopf, P. D. Goldan, H. I. Schiff, and E. E. Ferguson, J. Chem. Phys. 44, 4087 (1966).

658. P. H. Dawson and A. W. Tickner, J. Chem. Phys. 45, 4330 (1966).

659. A. A. Westenberg and N. de Haas, J. Chem. Phys. 47, 1393 (1967).

660. E. Graham, IV, D. R. James, W. C. Keever, I. R. Gatland, D. L. Albritton, and E. W. McDaniel, J. Chem. Phys. 59, 4648 (1973).

661. K. R. Ryan and I. G. Graham, J. Chem. Phys. 59, 4260 (1973).

662. F. C. Fehsenfeld, C. J. Howard, and E. E. Ferguson, J. Chem. Phys. 58, 5841 (1973).

663. N. G. Adams, D. K. Bohme, and E. E. Ferguson, J. Chem. Phys. 52, 5101 (1970).

664. D. W. Trainor, D. O. Ham, and F. Kaufman, J. Chem. Phys. 58, 4599 (1973).

665. T. J. Kvale, D. G. Seely, D. M. Blankenship, E. Redd, T. J. Gay, M. Kimura, E. Rille, J. L. Peacher, and J. T. Park, Phys. Rev. A 32, 1369 (1985).

666. D. L. Smith and J. H. Futrell, Chem. Phys. Lett. 40, 229 (1976).

667. M. Kneba, U. Wellhausen, and J. Wolfrum, Ber. Bunsenges. Phys. Chem. 83, 940 (1979).

668. R. Johnsen and M. A. Biondi, J. Chem. Phys. 61, 2112 (1974).

669. G. P. Glass and B. K. Chaturvedi, J. Chem. Phys. 77, 3478 (1982).

670. O. J. Orient, Chem. Phys. Lett. 52, 264 (1977).

671. A. L. Schmeltekopf and H. P. Broida, J. Chem. Phys. 39, 1261 (1963).

672. V. Aquilanti, A. Galli, A. Giardini-Guidoni, and G. G. Volpi, J. Chem. Phys. 43, 1969 (1965).

673. E. C. Beaty and P. L. Petterson, Phys. Rev. 137, A346 (1965).

674. E. B. Gordon, B. I. Ivanov, A. P. Perminov, V. E. Balalaev, A. N. Ponomarev, and V. V. Filatov, Chem. Phys. Lett. 58, 425 (1978).

675. D. J. Le Roy, B. A. Ridley, and K. A. Quickert, Disscuss. Faraday Soc. 44, 92 (1967).

676. A. V. Phelps, p. 690 in XV International Conference on the Physics of Electronic and Atomic Collisions, Brighton, UK, July 22-28, 1987.

677. W.G.F. Blair and H. B. Gilbody, J. Phys. B 6, 483 (1973).

678. J. Van Eck, F. J. de Heer, and J. Kistemaker, Physica 30, 1171 (1964).

679. J. M. Robinson and H. B. Gilbody, Proc. Phys. Soc. London 92, 589 (1967).

680. R. Hippler and K.-H. Schartner, J. Phys. B 7, 618 (1974).

681. J. T. Park, J. M. George, J. L. Peacher, and J. E. Aldag, Phys. Rev. A 18, 48 (1978).

682. J. T. Park and F. D. Schowengerdt, Phys. Rev. 185, 152 (1969).

683. D. Krause and E. A. Soltysik, Phys. Rev. 175, 142 (1968).

684. E. W. Thomas, ed. p. 134 in Excitation in Heavy Particle Collisions, Wiley-Interscience, New York, 1972.

685. D. Hasselkamp, R. Hippler, A. Scharmann, and K. H. Schartner, Z. Phys. 248, 254 (1971).

686. J. van den Bos, G. J. Winter, and F. J. de Heer, Physica 40, 357 (1968).

687. A. Scharmann and K.-H. Schartner, Z. Phys. 228, 254 (1969).

688. E. W. Thomas and G. D. Bent, Phys. Rev. 164, 143 (1967).

689. J. G. Dodd and R. H. Hughes, Phys. Rev. 135, A618 (1964).

690. A. Denis, M. Dufay, and M. Gaillard, C. R. Acad. Sci. Paris Ser. B 264, 440 (1967).

691. J. Hill, J. Geddes, and H. B. Gilbody, J. Phys. B 13, 951 (1980).

692. V. Dose and W. Hett, J. Phys. B 7, L454 (1974).

693. E. P. Andreev, V. A. Ankudinov, and S. V. Bobashev, Opt. Spectrosc. 16, 103 (1964).

694. R. D. Nathan and R. C. Isler, Phys. Rev. Lett. 26, 1091 (1971).

695. D. Dowek, D. Dhuicq, V. Sidis, and M. Barat, Phys. Rev. A 26, 746 (1982).

696. F. J. de Heer and J. van den Bos, Physica 31, 365 (1965).

697. G. Rahmat, G. Vassilev, and J. Baudon, J. Phys. (Paris) 41, 319 (1980).

698. D. Hasselkamp, R. Hippler, A. Scharmann, and K.-H. Schartner, Z. Phys. 257, 43 (1972).

699. V. Pol, W. Kauppila, and J. T. Park, Phys. Rev. A 8, 2990 (1973).

700. R. A. Young, R. F. Stebbings, and J. W. McGowan, Phys. Rev. 171, 85 (1968).

701. J.D.A. McKee, J. R. Sheridan, J. Geddes, and H. B. Gilbody, J. Phys. B 10, 1679 (1977).

702. V. A. Ankudinov, S. V. Bokashev, and E. P. Andreev, Sov. Phys.-JETP 25, 236 (1967).

703. W. G. F. Blair, R. W. McCullough, F. R. Simpson, and H. B. Gilbody, J. Phys. B 6, 1265 (1973).

704. D. Bordenave-Montesquieu, A. Boutonnet, C. Bergnes, and R. Dagnac, J. Phys. B 16, 1937 (1983).

705. R. Okasaka, Y. Konishi, Y. Sato, and K. Fukuda, J. Phys. B 20, 3771 (1987).

706. M. B. Shah and H. B. Gilbody, J. Phys. B 7, L122 (1974).

707. S. H. Dworetsky, R. Novick, and N. Tolk, p. 294 in VI International Conference on the Physics of Electronic and Atomic Collisions, MIT, July 28-August 2, 1969, The MIT Press, Cambridge, Massachusetts, 1969.

708. S. H. Dworetsky, R. Novick, W. W. Smith, and N. Tolk, Phys. Rev. Lett. 18, 939 (1967).

709. J. T. Park, J. E. Aldag, J. M. George, and J. L. Peacher, Phys. Rev. A 14, 608 (1976).

710. J. T. Park, J. E. Aldag, and J. M. George, Phys. Rev. Lett. 34, 1253 (1975).

711. D. R. Bates and G. Griffing, Proc. Phys. Soc. London 66, 64 (1953).

712. R. A. Young, R. F. Stebbings, and J. W. McGowan, Phys. Rev. 171, 85 (1968).

713. D. R. Schoonover and J. T. Park, Phys. Rev. A 3, 228 (1971).

714. A. I. Dashchenko, I. P. Zapesochnyi, A. I. Imre, V. S. Bukstich, F. F. Danch, and V. A. Kel'man, Sov. Phys.-JETP 40, 249 (1975).

715. J. Fayeton, J. C. Houver, M. Barat, and F. Masnou-Seeuws, J. Phys. B 9, 461 (1976).

716. J. E. Aldag, J. L. Peacher, P. J. Martin, V. Sutcliffe, J. George, E. Redd, J. T. Kvale, D. M. Blankenship, and J. T. Park, Phys. Rev. A 23, 1062 (1981).

717. D. R. Schoonover and J. T. Park, p. 839 in VII International Conference on the Physics of Electronic and Atomic Collisions, Amsterdam, July 26-30, 1971, North-Holland Publishing Co., Amsterdam, 1971.

718. M. Kimura, T. Egawa, and T. Iwai, p. 554 in XI International Conference on the Physics of Electronic and Atomic Collisions, Kyoto, August 29-September 4, 1979, 1979.

719. V. Kempter, F. Veith, and L. Zehnle, J. Phys. B 8, 1041 (1975).

720. R. W. McCullough, F. R. Simpson, and H. B. Gilbody, J. Phys. B 6, L322 (1973).

721. R. E. Olson, E. J. Shipsey, and J. C. Browne, J. Phys. B 8, 905 (1975).

722. J. P. Gauyacq, J. Phys. B 9, 3067 (1976).

723. V. Kempter, G. Riecke, F. Veith, and L. Zehnle, J. Phys. B 9, 3081 (1976).

724. A. Gleizes, A. Bordenave-Montesquieu, and P. Benoit-Cattin, J. Phys. B 14, 4545 (1981).

725. A. Bordenave-Montesquieu, P. Benoit-Cattin, M. Boudjema, and A. Gleizes, p. 996 in X International Conference on the Physics of Electronic and Atomic Collisions, Paris, July 21-27, 1977, North-Holland Publishing Co., Amsterdam, 1967.

726. F. D. Schowengerdt, S. R. Smart, and M. E. Rudd, Phys. Rev. A 7, 560 (1973).

727. R. W. McCullough, F. R. Simpson, and H. B. Gilbody, J. Phys. B 6, L322 (1973).

728. M. Kimura, T. Egawa, and T. Iwai, p. 554 in XI International Conference on the Physics of Electronic and Atomic Collisions, Kyoto, August 29-September 4, 1979, 1979.

729. D. V. Elakhovsky, Yu. V. Zaitsev, and A. D. Khakhaev, p. 346 in XIII International Conference on the Physics of Electronic and Atomic Collisions, Berlin, July 27-August 2, 1983.

730. R. W. McCullough, W.G.F. Blair, F. R. Simpson, and H. B. Gilbody, p. 847 in VIII International Conference on the Physics of Electronic and Atomic Collisions, Beograd, July 16-20, 1973.

731. A. M. Ermolaev, R. N. Hewitt, R. Shingal, and M.R.C. McDowell, J. Phys. B 20, 4507 (1987).

732. F. Aumayr, G. Lakits, and H. Winter, Z. Phys. D 6, 145 (1987).

733. K. Kadota, D. Dijkkamp, R. L. van der Woude, P. G. Yan, and F. J. de Heer, J. Phys. B 15, 3297 (1982).

734. G. D. Myers, J. L. Barrett, and J. J. Leventhal, Phys. Rev. A 20, 797 (1979).

735. A. M. Ermolaev and R. N. Hewitt, Nucl. Instrum. Methods Phys. Res. B 9, 487 (1985).

736. J. L. Barrett and J. J. Leventhal, Phys. Rev. A 23, 485 (1981).

737. M. B. Shah, D. S. Elliott, and H. B. Gilbody, p. 405 in XIV International Conference on the Physics of Electronic and Atomic Collisions, Palo Alto, July 24-30, 1985.

738. R. E. Olson, J. Phys. B 15, L163 (1982).

739. S. N. Tiwary and D. K. Rai, J. Chem. Phys. 68, 2427 (1978).

740. F. Aumayr and H. Winter, Institut fur Allgemeine Physik Report 1/87, March (1987).

APPENDIX 1

Calculation of Cross Sections and Rate Coefficients from Chebyshev Fitting Parameters

APPENDIX 1

Calculation of Cross Sections and Rate Coefficients from Chebyshev Fitting Parameters

For ease of numerical data retrieval, the method of least-squares fitting to the recommended cross sections and rate coefficients using the Chebyshev orthogonal polynomials has been used throughout this compilation. In general, it produced high accuracy fits to the recommended data.

A measure of the accuracy of the Chebyshev fit representation to the recommended cross sections has been calculated and appears at the bottom of each cross section page as a relative rms error with a maximum deviation. In a handful of cases, the recommended cross sections had associated Chebyshev fits with an error of greater than 100% rms error. These fits have not been included and only the recommended cross section data is presented.

The recommended cross-section and rate coefficient data have been fitted to the following analytical expression:

$$\ell n[\sigma(E)] = A_0/2 + \sum_{i=1}^{k} A_i T_i(X) \ ,$$

where k is the smallest number of coefficients that provides an accurate fit (usually k=6), and

$$X = [(\ell nE - \ell nE_{min}) - (\ell nE_{max} - \ell nE)] \ / \ (\ell nE_{max} - \ell nE_{min}),$$

and where $T_i(X)$ are the Chebyshev orthogonal polynomials. Fitting to the natural logarithms of the cross section and energy values increases the Chebyshev fit accuracy by reducing the ranges of values for the cross section and energy by several orders of magnitude.

The values of fitting parameters A_i, E_{min}, and E_{max} are given for each reaction after each cross-section or rate-coefficient table. In each case, the minimum number of parameters A_i are listed which are required to produce a satisfactory fit to the data. The cross section σ is given in units of cm^2, with collision energies E in units of eV/amu. The rate coefficient α is in units of cm^3/s, with Maxwellian temperatures T in units of eV. (We follow the convention used frequently in plasma physics of expressing temperature in eV energy units, recognizing that kT is understood, where k is Boltzmann's constant.) The analytic representation of the cross section or rate coefficient should be used only in the region between E_{min} and E_{max}. The expressions for the Chebyshev polynomials are reproduced below for convenience.

$T_1(X) = X$
$T_2(X) = 2\ X^2 - 1$
$T_3(X) = 4\ X^3 - 3\ X$
$T_4(X) = 8\ X^4 - 8\ X^2 + 1$
$T_5(X) = 16\ X^5 - 20\ X^3 + 5\ X$
$T_6(X) = 32\ X^6 - 48\ X^4 + 18\ X^2 - 1$
$T_7(X) = 64\ X^7 - 112\ X^5 + 56\ X^3 - 7\ X$
$T_8(X) = 128\ X^8 - 256\ X^6 + 160\ X^4 - 32\ X^2 + 1$

All the rate-coefficient and cross-section data in this volume may be reproduced by using either of the model Chebyshev fitting programs listed at the end of this Appendix. For each reaction and pair of reactants there are given Chebyshev fitting parameters to be used as input to the CHEBFOR Fortran or CHEB Basic code.

In the plots of recommended data the Chebyshev or alternative fits have been added as dashed lines to give a visual inspection of the accuracy of the fit. Also, the pages containing cross section tables include rms deviations of the fit to the recommended data over the energy range from E_{min} to E_{max}.

The sample programs listed below utilize recursion relations to rapidly generate the Chebyshev polynomials and require input of the parameters E_{min}, E_{max}, and the Chebyshev coefficients A0, A1, A2, ... to enable calculation of a rate coefficient or cross section at a given energy. The programs direct the user to input these parameters at the proper time during program execution. To simplify data input, the tabulations of coefficients contain the minimum number of significant digits required to produce an accurate representation of the polynomial fit.

The ALADDIN database system may also be used to generate the cross-section data directly from the fitting coefficients. See Appendix 2 for details.

Sample FORTRAN Program

```
      Program CHEBFOR
C     This Fortran program is derived from a program in 'Elementary Numerical Analysis: An
C     Algorithmic Approach' , S. D. Conte and C. de Boor, McGraw-Hill, Inc., P.254, 1972.
      DIMENSION D(10)
      REAL*4 ANS, ANS2
1000  TYPE 14
14    FORMAT('   IS THIS A FIT TO RATE COEFFICIENTS? ; ENTER "Y" FOR YES, "N" FOR NO')
      ACCEPT 199, ANS
199   FORMAT(A1)
1010  TYPE 15
15    FORMAT('   ENTER THE NUMBER OF COEFFICIENTS')
      ACCEPT 200, NTERMS
200   FORMAT(I2)
```

Sample FORTRAN Program (Cont'd.)

```
      IF(ANS .EQ. 'N') GO TO 16
      TYPE 1
1     FORMAT(' **** ENTER Emin (eV) *****')
      ACCEPT *, EMIN
      TYPE 9
9     FORMAT(' **** ENTER Emax (eV) *****')
      ACCEPT *, EMAX
      GO TO 25
19    FORMAT(' **** ENTER Emin (eV/amu) ****')
16    TYPE 19
      ACCEPT *, EMIN
      TYPE 29
29    FORMAT(' **** ENTER Emax (eV/amu) ****')
      ACCEPT *, EMAX
25    EMINL=ALOG(EMIN)
      EMAXL=ALOG(EMAX)
      TYPE 2, NTERMS
2     FORMAT('  ENTER ', I2, ' COEFFICIENTS ')
      ACCEPT *, (D(J), J=1,NTERMS)
100   IF (ANS .EQ. 'N')GO TO 24
      TYPE 3, EMIN, EMAX
3     FORMAT('  INPUT TEMPERATURE (eV) BETWEEN ',1PE10.1,' AND ',1PE10.1,' FOR THE RATE
   & COEFFICIENT CALCULATION  ')
      GO TO 26
24    TYPE 23, EMIN, EMAX
23    FORMAT('  INPUT ENERGY (eV/amu) BETWEEN ',1PE10.1,' AND ',1PE10.1,' FOR THE CROSS SECTION
   & CALCULATION  ')
C     TAKE ENERGY FOR CALCULATION AND USE THREE TERM RECURRENCE RELATION
26    K=NTERMS
      CHEB=D(K)
      ACCEPT *, X
      IF(X .LT. 0.) GO TO 101
      X=ALOG(X)
      K=K-1
      XNORM=(X-EMINL-(EMAXL-X))/(EMAXL-EMINL)
      TWOX=2.*XNORM
      PREV2=0.
10    PREV=CHEB
      IF(K .EQ. 1)  GO TO 20
      CHEB=D(K)+TWOX*PREV-PREV2
      PREV2=PREV
      K=K-1
      GO TO 10
20    CHEB=.5*D(1)+XNORM*PREV-PREV2
      IF(ANS .EQ. 'Y') TYPE 69, EXP(CHEB)
69    FORMAT('  RATE COEFFICIENT (CM3/S) = ',1PE12.4)
      IF(ANS .EQ. 'N') TYPE 45, EXP(CHEB)
45    FORMAT('  CROSS SECTION (CM2) = ',1PE12.4)
      TYPE 48
48    FORMAT('  DO YOU WANT ANOTHER CALCULATION FOR THIS FIT?, ENTER "Y" FOR YES; "N" FOR NO')
      ACCEPT 199, ANS2
      IF(ANS2 .EQ. 'Y') GO TO 100
      TYPE 49
49    FORMAT('  DO YOU WANT A DIFFERENT FIT CALCULATION?, ENTER "Y" FOR YES;  "N" FOR NO')
      ACCEPT 199, ANS2
      IF(ANS2 .EQ. 'Y') GO TO 1000
101   END
```

Sample BASIC Program

```
5  REM CHEB.BAS
10 REM CHEB PROGRAM IS A BASIC PROGRAM DERIVED FROM A FORTRAN
20 REM PROGRAM IN "ELEMENTARY NUMERICAL ANALYSIS: AN ALGORITHMIC
30 REM APPROACH", S. D. CONTE AND C. de BOOR, 30 REM McGRAW-HILL, INC., P
40 REM 254, 1972.
50    DIM D(10)
60    INPUT "IS THE A RATE FIT?; ENTER 'Y' FOR YES, 'N' FOR NO   ",ANS$
70    IF ANS$="Y" THEN INPUT "ENTER Emin (eV)  ",EMIN
80    IF ANS$="Y" THEN INPUT "ENTER Emax (eV)  ",EMAX
90    IF ANS$="N" THEN INPUT "ENTER Emin (eV/amu)  ",EMIN
100    IF ANS$="N" THEN INPUT "ENTER Emax (eV/amu)  ",EMAX
110    INPUT "HOW MANY COEFFICIENTS FOR THIS FIT?";NTERMS
120   EMINL=LOG(EMIN)
130   EMAXL=LOG(EMAX)
140   PRINT USING"ENTER # COEFFICIENTS";NTERMS
150   FOR J=1 TO NTERMS
160   INPUT D(J):NEXT J
170   IF ANS$="Y" THEN PRINT USING "INPUT ENERGY (eV/amu) BETWEEN ###.# AND #.#^^^^ FOR THE RATE
COEFFICIENT CALCULATION";EMIN;EMAX
180   IF ANS$="N" THEN PRINT USING "INPUT ENERGY (eV/amu) BETWEEN ###.# AND #.#^^^^ FOR THE CROSS
SECTION CALCULATION";EMIN;EMAX
190 REM GET ENERGY FOR CALCULATION AND USE THREE TERM RECURRENCE
200 REM RELATION
210   INPUT X
220   K=NTERMS
230   CHEB=D(K)
240   IF X<0 THEN END
250   X=LOG(X)
260   K=K-1
270   IF K=0 THEN END
280   XNORM=(X-EMINL-(EMAXL-X))/(EMAXL-EMINL)
290   TWOX=2*XNORM
300   PREV2=0
310   PREV=CHEB
320   IF K=1 GOTO 370
330   CHEB=D(K)+TWOX*PREV-PREV2
340   PREV2=PREV
350   K=K-1
360   GOTO 310
370   CHEB=.5*D(1)+XNORM*PREV-PREV2
380   IF ANS$="Y" THEN PRINT USING"##.##^^^^ = RATE COEFFICIENT (CM3/S)";EXP(CHEB)
390   IF ANS$="N" THEN PRINT USING"##.##^^^^ = CROSS SECTION (CM2)";EXP(CHEB)
400   GOTO 170
410   END
```

APPENDIX 2

ALADDIN Database System and Data Files

APPENDIX 2

ALADDIN Database System and Data Files

ALADDIN[1,2] is an atomic physics database system which has been developed by R. A. Hulse of Princeton Plasma Physics Laboratory in conjunction with the Atomic and Molecular Data Unit of the International Atomic Energy Agency (IAEA) to provide a broadly-based standard medium for the exchange and management of atomic data. ALADDIN has been adopted by the IAEA as the standard international atomic physics data exchange format for magnetic confinement fusion applications.

ALADDIN consists of a data format definition together with supporting software for both interactive searches and access to the data by plasma modelling and other computer codes. The ALADDIN software has been written in strict FORTRAN-77 to ensure compatibility with the widest possible range of scientific computers. The data are stored in straightforward ASCII text files either in tabular x,y or parametrized format. Provision has been made within ALADDIN to incorporate the subroutines which are needed to extract the atomic data from each ALADDIN data entry. The Chebyshev polynomial fits which are presented for the data in this volume are one example of such a parametrization.

All recommended cross-section data from the compilations in the **Atomic Data for Fusion** series are available in ALADDIN format (including those from this volume). Table 2.1 contains a sample ALADDIN data entry from each chapter of this volume. Each entry contains a page-number label to expedite locating specific data within the file. Complete ALADDIN data files for this series and a copy of the ALADDIN Database program are available on PC-DOS-formatted diskettes from:

Controlled Fusion Atomic Data Center
Oak Ridge National Laboratory
P. O. Box 2008
MS-6372
Oak Ridge, TN 37831-6372, U.S.A.

A copy of the ALADDIN Manual[2] which contains instructions for its use and a dictionary of abbreviations and symbolic notations for collision processes is also available from the above address. The ALADDIN code and reference material are also available directly from the IAEA (see address below).

The ALADDIN data files and FORTRAN program may also be transmitted readily by electronic mail, and are available by contacting:

phaneuf@orn.MFENET or phaneuf@orph01.BITNET

With input from a number of international data centers, the Atomic and Molecular Data Unit of the IAEA is assembling a larger ALADDIN data base of atomic and molecular processes relevant to fusion applications, and is providing officially recommended data in ALADDIN format. Further information may be obtained by contacting:

Atomic and Molecular Data Unit
Nuclear Data Section
International Atomic Energy Agency
P. O. Box 100
A-1400 Vienna, Austria

or by electronic mail at:

RNDS@IAEA1.BITNET

---

1. R. A. Hulse, "The ALADDIN Atomic Physics Database System," pp. 63-72 in **Atomic Processes in Plasmas**, AIP Conference Proceedings 206, ed. Y. K. Kim and R. C. Elton, AIP, New York, 1990.

2. **ALADDIN Manual, A System for Storage, Exchange and Management of Atomic and Molecular Data for Fusion**, IAEA-NDS-AM-17, Atomic and Molecular Data Unit, Nuclear Data Section, International Atomic Energy Agency, Vienna, June 1989.

TABLE 2.1

Sample ALADDIN Data File

```
$ CX  H [+1]  H [+0]   H [+0]  H [+1]
& XS   EVAL  ACC=B  ORNL-CFADC  DOC=ORNL-6086  A-22  6/90  #CHEB
-72.6656        -5.49142        -3.42948       -1.98377      -0.878009
 -0.198932       0.0837431      0.121252        0.0827182  0.12  630000.

$ CXSS  He [+2]  H [+0] (G)   He [+1] (2s)  H [+1]
& XS   EVAL  ACC=C  ORNL-CFADC  DOC=ORNL-6086  B-89  6/90  #CHEB
-89.4268        -0.438776       -8.73454        0.130314       -0.0101210
  0.0343355      0.227699       -0.188121      -0.130396   23.  2500000.

$ EXC  H [+0] (G)  H [+0] (G)   H [+0] (2s)  H [+0]
& XS   EVAL  ACC=?  ORNL-CFADC  DOC=ORNL-6086  C-2  8/89  #CHEB
-79.1098        -0.784062       -0.651723       0.0462865      0.114234
 -0.00487654    -0.0114391       0.0            0.0     2000.   90000.

$ ION  H [+1]  H [+0]   H [+1]  H [+1]  e
& XS   EVAL  ACC=B  ORNL-CFADC  DOC=ORNL-6086  D-6  5/90  #CHEB
-75.505348206  -0.409784913  -0.996553540   0.355360299  -0.049000371
 -0.034998417   0.043782476   0.0           0.0    9400.    1500000.

$ STRIP  H [+0]  H [+0]   H [+1]  H [+0]  e
& XS   EVAL  ACC=B  ORNL-CFADC  DOC=ORNL-6086  E-2  5/90  #CHEB
-77.1793823242 -0.4514082074 -1.5175991058  0.5439749956 -0.0355154276
  0.0025895350 -0.0750899166  0.0           0.0     1200. 3500000.

$ ELDET  H [-1]  H [+0]   H [+0]  H [+0]  e
& XS   EVAL  ACC=C  ORNL-CFADC  DOC=ORNL-6086  F-2  5/90  #CHEB
-72.0698776245 -2.4792380333 -0.9363254905  0.0886530057  0.0175392255
  0.0545278080 -0.0645320490  0.0           0.0      110. 4000000.

$ DISION2  H [+0]  H(2) [+0]   H [+0]  H [+1]  H [+1]  2e
& XS   EVAL  ACC=?  ORNL-CFADC  DOC=ORNL-6086  G-2  6/90  #CHEB
-82.5844192505  1.5008718967 -0.5141620040 -0.0138262566  0.0274858437
 -0.0085383281 -0.0116637163  0.0           0.0     5000. 50000.

$ PX  H [+1]  D(2) [+0]   D [+1]  HD [+0]
& XS   EVAL  ACC=?  ORNL-CFADC  DOC=ORNL-6086  H-2  9/89  #CHEB
-75.6467819214 -3.2291088104  0.4121128917  0.0393397696 -0.2617489398
  0.0012951158  0.1208853275  0.0           0.0     0.45    100.
```

APPENDIX 3

Atomic Physics

Handy Conversion Units and Formulas

APPENDIX 3

Atomic Physics

Handy Conversion Units and Formulas

Conversion Units

1 electron volt = 1 eV = 8065.76 cm^{-1} = 23.069 kcal/mol = 1.6022×10^{-12} ergs = 1.6022×10^{-19} joules

1 atomic unit = 1 a.u. = 2 Rydbergs = 27.21 eV = 1 Hartree

1 atomic mass unit = 1 a.m.u. = 1.6605×10^{-24} gm

Atomic velocity = velocity of 13.6 eV electron = 2.18×10^{8} cm/sec = velocity of 24.97 keV proton

Atomic radius = radius of first Bohr orbit of H° = 5.29×10^{-9} cm

1 angstrom = 1 Å = 1×10^{-8} cm

Electron mass, $m_e = 9.110 \times 10^{-28}$ gm

Electron charge, $e = 4.8032 \times 10^{-10}$ statcoulomb, or 1.6011×10^{-19} coulombs

Proton mass = 1.673×10^{-24} gm

Atomic unit of time = atomic radius/atomic velocity = 2.42×10^{-17} sec

Boltzmann constant, $k = 1.381 \times 10^{-16}$ erg/°K

1 mm Hg = 133.3 pascals = 1 torr; 1 mBar = 100 pascals

Atomic cross section = $\pi \times (\text{atomic radius})^2 = \pi a_o^2 = 8.797 \times 10^{-17}$ cm^2

1 mol = 6.022169×10^{23} molecules = Avogadro's number

Loschmidt's number = number of molecules at 1 mm Hg at 0°C = 2.687×10^{16} cm^{-3}

Formulas

Velocity (non-relativistic) (cm/sec) = $1.38 \times 10^6 \, (E/m)^{1/2}$, where E = energy of the particle in eV, of mass m in a.m.u.

Energy levels in hydrogen-like atoms,

$$E_n(\text{eV}) = \frac{-13.6 \, Z^2}{n^2};$$

where Z is the atomic number and n is the total quantum number.

Wavelength of a photon,

$$\lambda(\text{Å}) = \frac{1.2398 \times 10^4}{E},$$

where E is the photon energy in eV.

Wavelength of a non-relativistic electron,

$$\lambda(\text{Å}) = \frac{17.34}{T^{1/2}},$$

where T is the temperature of the electrons in eV.

Relative velocity (cm/s) = $V_r = (V_1^2 + V_2^2 - 2V_1V_2\cos\theta)^{1/2}$, where V_1 = velocity of particle 1 in cm/sec, V_2 = velocity of particle 2, and θ = angle of approach.

Energy (center of mass) (eV) =

$$\frac{M_1M_2}{M_1+M_2}\left[\frac{E_1}{M_1}+\frac{E_2}{M_2}-2\left(\frac{E_1E_2}{M_1M_2}\right)^{1/2}\cos\theta\right]$$

where M_1, E_1 are the mass and energy of particle 1 and M_2, E_2 are the mass and energy of particle 2, respectively, and θ is the angle of approach.

Plasma Physics

Handy Formulas

*Gaussian units are used in the formulas below except for the following: Temperatures, T_e of electrons, and T_i of ions in eV. Mass of ions, m_i is expressed in units of proton mass, m_p by $\mu = m_i/m_p$; B in gauss; k is Boltzmann's constant; n_e and n_i are number densities of electrons and ions, respectively, in cm^{-3}; Z is the charge state of an ion; λ is wavelength in Å; c is the speed of light in vacuum (3×10^{10} cm/sec); and frequency is ω (rad/sec) = $2\pi \cdot f$(cycles/sec)

For Electrons:

Plasma frequency, ω_{pe} (rad/sec) $= 5.64 \times 10^4\ n_e^{1/2}$
Gyrofrequency, ω_{ce} (rad/sec) $= 1.76 \times 10^7\ B$
Gyroradius, r_e (cm) $= 2.38\ T_e^{1/2}\ B^{-1}$
Electron thermal velocity, v_{te} (cm/sec) $= 4.19 \times 10^7\ T_e^{1/2}$

Electron deBroglie length, λ (Å) $= \frac{17.34}{T_e^{1/2}}$

For Ions:

Plasma frequency, ω_{pi} (rad/sec) $= 1.32 \times 10^3\ Z\ \mu^{-1/2}\ n_i^{1/2}$
Gyrofrequency, ω_{ci} (rad/sec) $= 9.58 \times 10^3\ Z\ \mu^{-1}\ B$
Gyroradius, r_i (cm) $= 1.02 \times 10^2\ \mu^{1/2}\ Z^{-1/2}\ T_i^{1/2}\ B^{-1}$
Ion thermal velocity, v_{ti} (cm/sec) $= 9.79 \times 10^5\ \mu^{-1/2}\ T_i^{1/2}$

Drift velocity of particles in a constant electric field, E (volts/cm), perpendicular to a constant magnetic field, B (gauss), independent of mass m, or charge Z, of the particle is given by:

$$v_E \text{ (cm/sec)} = 1 \times 10^8 \frac{E}{B}$$

Phase velocity, v_ϕ (cm/sec) $= \omega/k$, where ω is in rad/sec and k is the wavenumber in cm^{-1}.

Group velocity, v_g (cm/sec) $= d\omega/dk$

Alfven velocity, v_A (cm/sec) $= 2.18 \times 10^{11}\ \mu^{-1/2}\ n_i^{-1/2}\ B$

Debye length, λ_D (cm) $= 7.43 \times 10^2\ T_e^{1/2}\ n_e^{-1/2}$

Maxwellian velocity distribution for a constant temperature plasma, for one-dimension:

$$F_{max}(v) \ (cm^{-3}) = n \cdot (m/2\pi kt)^{1/2} \exp(-mv^2/2kt) \ ,$$

where t is in °K, m is the particle's mass in gm, n is the number density (cm^{-3}), k is Boltzmann's constant (1.38×10^{-16} erg/°K), and v is the particle's velocity (cm/sec) for one-dimension.

Saha thermal equilibrium:

$$n_i/n_n \approx 2.4 \times 10^{15} \ (t^{3/2}/n_i) \ \exp(-U_i/kt) \ ,$$

where n_n is the number density of neutrals in cm^{-3}, U_i is the ionization energy of the gas in ergs, and t is the temperature of the gas in °K.

INTERNAL DISTRIBUTION

1. R. J. Colchin
2. R. N. Compton
3. E. C. Crume
4. W. K. Dagenhart
5. S. Datz
6. R. A. Dory
7-11. D. C. Gregory
12-16. J. W. Hale
17. H. H. Haselton
18-22. C. C. Havener
23. J. T. Hogan
24-28. H. T. Hunter
29-33. D. P. Hutchinson
34. R. C. Isler
35-39. M. I. Kirkpatrick
40. C. C. Klepper
41. H. F. Krause
42-46. C. H. Ma
47-51. F. W. Meyer
52-56. R. A. Phaneuf
57-61. R. K. Richards
62. J. A. Rome
63. P. M. Ryan
64. I. A. Sellin
65-69. J. S. Thompson
70. C. C. Tsai
71-75. K. L. Vander Sluis
76. J. H. Whealton
77. T. L. White
78. J. B. Wilgen
79-83. P. A. Zeijlmans
84. Central Research Library
85. Document Reference Section
86-87. Laboratory Records
88. Laboratory Records, RC
89. ORNL Patent Section
90. Fusion Energy Div. Library
91. Fusion Eng. Design Center

EXTERNAL DISTRIBUTION - U.S.

92. Office of Assistant Manager, Energy Research and Development, DOE-ORO
93. Dr. J. M. Ajello, Stop 183-601, Jet Propulsion Lab., Calif. Inst. of Technology, 4800 Oak Grove Dr., Pasadena, CA 91109
94. Dr. I. Alexeff, Dept. Elect. Eng., Ferris Hall, Univ. of Tennessee, Knoxville, TN 37996-2100
95. Dr. M. A. Ali, Chem. Dept., Howard Univ., Washington, DC 20059
96. Dr. P. Allison, Los Alamos National Lab., P.O. Box 1663, MS H-818, Los Alamos, NM 87545
97. Dr. R. C. Amme, Dept. of Physics, Univ. of Denver, Denver, CO 80208
98. Prof. L. W. Anderson, Dept. of Physics, Univ. of Wisconsin, Madison, WI 53706
99. Dr. E. H. Avrett, Center for Astrophysics, 60 Garden St., Cambridge, MA 02138
100. Dr. J. F. Baur, General Atomics, P.O. Box 85608, San Diego, CA 92138
101. Prof. B. Bayer, Ohio State Univ., 1011 Smith Lab., Physics Library, 174 West 18th Ave., Columbus, OH 43210
102. Dr. J. E. Bayfield, Dept. of Physics, Univ. of Pittsburgh, Pittsburgh, PA 15260
103. Prof. B. Bederson, Dept. of Physics, New York Univ., 4 Washington Place, New York, NY 10003
104. Prof. G. Bekefi, Dept. of Physics, Room 36-213, Massachusetts Inst. of Techn., Cambridge, MA 02139
105. Dr. R. D. Bengtson, Dept. of Physics, Univ. of Texas, Austin, TX 78712-1081
106. Prof. W. R. Bennett, 102 Dunham Laboratory, Yale Univ., New Haven, CT 06520
107. Dr. K. H. Berkner, Bldg. 50, Rm. 149, 1 Cyclotron Rd., Lawrence Berkeley Lab., Berkeley, CA 94720
108. Prof. R. S. Berry, Dept. of Chem., The Univ. of Chicago, 5735 South Ellis Ave., Chicago, IL 60637
109. Dr. M. Bitter, Plasma Physics Lab., Princeton Univ., P.O. Box 451, Princeton, NJ 08544
110. Dr. D. C. Boice, Space Science Div., Southwest Research Inst., P.O. Drawer 28510, 6220 Culebra Rd., San Antonio, TX 78228-0510
111. Dr. G. O. Brink, Dept. Physics & Astronomy, State Univ. of New York at Buffalo, Amherst, NY 14260
112. Dr. S. Bross, Univ. of Missouri-Rolla, Dept. of Physics, Rolla, MO 65401
113. Dr. R. Bruch, Dept. of Physics, Univ. of Nevada, Reno, Reno, NV 89557-0058
114. Dr. K. H. Burrell, General Atomics, P.O. Box 85608, San Diego, CA 92186-9704
115. Dr. J. Callaway, Dept. of Phys. and Astronomy, Louisiana State Univ., Baton Rouge, LA 70803-4001
116. Dr. J. D. Callen, Nucl. Eng. Dept., ERB, 1500 Johnson Dr., Univ. of Wisconsin, Madison, WI 53706-1687
117. Dr. D. Casperson, Group P-1, MS E-526, Los Alamos National Lab., Los Alamos, NM 87545

118. Dr. S. Chakrabarti, Space Sciences Lab., Univ. California, Berkeley, Berkeley, CA 94720
119. Dr. R. L. Champion, Dept. of Physics, College of William & Mary, Williamsburg, VA 23185
120. Dr. F. Chang, Astronaut Office/Code CB, Lyndon B. Johnson Space Center, Houston, TX 77058
121. Dr. M. H. Chen, LLNL, V Div., L-296, P.O. Box 808, Livermore, CA 94550
122. Dr. C. K. Choi, School of Nuclear Eng., Purdue Univ., West Lafayette, IN 47907
123. Dr. Y. P. Chong, LLNL, P.O. Box 808, MS L-390, Livermore, CA 94550
124. Prof. Shih-I Chu, Dept. of Chem., Univ. of Kansas, Lawrence, KS 66045-0046
125. Dr. D. A. Church, Physics Dept., Texas A- Univ., College Station, TX 77843
126. Dr. A. Chutjian, Jet Propulsion Lab., Mail Stop 183-601, California Inst. of Techn., 4800 Oak Grove Dr., Pasadena, CA 91109
127. Dr. E. J. Clothiaux, Dept. of Physics, Auburn Univ., Auburn, AL 36849-3501
128. Dr. C. L. Cocke, Dept. of Physics, Cardwell Hall, Kansas State Univ., Manhattan, KS 66502
129. Dr. F. H. Coensgen, Radiation Lab., L-637, LLNL, P.O. Box 808, Livermore, CA 94550
130. Dr. M. J. Coggiola, Molecular Physics Lab., SRI International, 333 Ravenswood Ave., Menlo Park, CA 94025-3493
131. Dr. S. A. Cohen, Plasma Physics Lab., James Forrestal Campus, P.O. Box 451, Princeton Univ., Princeton, NJ 08544
132. Prof. G. J. Collins, Colorado State Univ., Dept. Electrical Eng., Fort Collins, CO 80523
133. Prof. R. W. Conn, School of Eng., Inst. of Plasma & Fusion Research, Univ. of California, Los Angeles, CA 90024
134. Dr. K. Connor, Plasma Dynamics Lab., ECSE Dept., Rensselaer Polytechnic Inst., Troy, NY 12180-3590
135. Dr. D. L. Cook, Organization 1260, Sandia Labs., P.O. Box 5800, Albuquerque, NM 87185
136. Dr. D. H. Crandall, ER-54, GTN, Dept. of Energy, Washington, DC 20545
137. Dr. E. Crawford, Spectra Technology, Inc., 2755 Northup Way, Bellevue, WA 98004-1495
138. Dr. A. Dalgarno, Harvard-Smithsonian Center for Astrophysics, 60 Garden St., MS 14, Cambridge, MA 02138
139. Dr. R. A. Dandl, Applied Microwave Plasma Concepts, 2075-N Corte Del Nogal, Carlsbad, CA 92009
140. Dr. R. Danford, Intl. Research & Evaluation, Attn: Input Processing Director, 21098 IRE Control Center, Eagan, MN 55121
141. Mr. B. Davis, Atom Sciences Inc., 114 Ridgeway Center, Oak Ridge, TN 37830
142. Dr. H. Davis, LANL, P.O. Box 1663, MS E-526, Los Alamos, NM 87545
143. Dr. J. Davis, Code 4720, Radiation Hydro. Branch, Naval Research Lab., Washington, DC 20375-5000
144. Prof. J. J. Dorning, Dept. of NEEP, Thornton Hall (Reactor Facility), Univ. of Virginia, Charlottesville, VA 22901
145. Dr. R. Doyas, Lawrence Livermore National Lab., P.O. Box 808, Livermore, CA 94550
146. Dr. R. Dubois, Pacific Northwest Labs., P.O. Box 999 (P8-47), Richland, WA 99352
147. Dr. R. J. Dukart, Div. 1262, Sandia National Labs., P.O. Box 5800, Albuquerque, NM 87185
149. Dr. R. W. Dunford, PHY-203, Argonne National Lab., 9700 S. Cass Ave., Argonne, IL 60439
150. Dr. G. H. Dunn, JILA, Campus Box 440, Univ. of Colorado, Boulder, CO 80309-0440
151. Dr. H. F. Dylla, Princeton Plasma Physics Lab., Princeton Univ., P.O. Box 451, Princeton, NJ 08543
152. EG&G Idaho, Inc., Attn: Inel Techn. Library-I.F., P.O. Box 1625, Idaho Falls, ID 83415
153. Dr. R. C. Elton, Code 4780, Naval Research Lab., Washington, DC 20375-5000
154. Prof. G. A. Emmert, Nuclear Eng. Dept., Univ. of Wisconsin, 153 Eng. Research Bldg., Madison, WI 53706
155. Dr. B. Feinberg, Lawrence Berkeley Lab., Univ. of California, MS 51-208, Berkeley, CA 94720
156. Dr. U. Feldman, Code 4174, Naval Research Lab., Washington, DC 20375
157. Dr. J. H. Fink, 4023 East Ave., Hayward, CA 94542
158. Dr. R. K. Fisher, L-510, General Atomics, P.O. Box 85608, San Diego, CA 92186-9784
159. Dr. M. R. Flannery, Georgia Inst. of Techn., School of Physics, Atlanta, GA 30332-0430
160. Dr. H. H. Fleischmann, Dept. of Applied Physics, Clark Hall, Cornell Univ., Ithaca, NY 14853-2501
161. Dr. R. Fortner, Lawrence Livermore National Lab., V Division, P.O. Box 808, L-296, Livermore, CA 94550
162. Dr. T. K. Fowler, Univ. of California, Berkeley, Dept. Nuclear Engineering, Etcheverry Hall, Rm 4153, Berkeley, CA 94720
163. Dr. R. S. Freund, AT&T Bell Labs., Rm. 1D-256, 600 Mountain Ave., Murray Hill, NJ 07974
164. Dr. J. W. Gallagher, Physics A-323, NIST, Gaithersburg, MD 20899
165. Dr. A. Garscadden, Aero Propulsion Lab., AFWAL-POOC-3, Bldg. 450, Wright-Patterson AFB, Dayton, OH 45433

166. Dr. T. J. Gay, Univ. of Missouri-Rolla, Dept. of Physics, 102 Physics Bldg., Rolla, MO 65401-0249
167. Dr. K. W. Gentle, Dept. of Physics, Univ. of Texas, Austin, TX 78712
168. Dr. G. H. Gillespie, G. H. Gillespie Assoc., P.O. Box 2961, Del Mar, CA 92014
169. Dr. H. J. Goldberg, HO-36, Westinghouse Hanford Co., 3200 George Washington Way, Richland, WA 99352
170. Dr. M. J. Gouge, U. S. Dept. of Energy, Oak Ridge Operations Office, Federal Bldg. Rm. 2024, Oak Ridge, TN 37830
171. Dr. A.E.S. Green, ICAAS SSRB, Univ. of Florida, Gainesville, FL 32611
172. Dr. H. R. Griem, Lab. for Plasma\Fusion, Univ. of Maryland, College Park, MD 20742-3511
173. Dr. L. Grodzins, Physics Dept., MIT, Rm 26-421, Cambridge, MA 02139
174. Dr. R. J. Groebner, MS 13-463, 13/463, General Atomics, P.O. Box 85608, San Diego, CA 92186-9784
175. Prof. R. A. Gross, Dept. of Applied Physics, Columbia Univ., New York, NY 10027
176. Dr. G. E. Guest, Applied Microwave Plasma Concepts, Inc., 2075-N Corte Del Nogal, Carlsbad, CA 92009
177. Dr. S. Hagmann, Dept. of Physics, Kansas State Univ., Manhattan, KS 66506
178. Dr. W. Halverson, Spire Corp., Patriots Park, Bedford, MA 01730
179. Dr. T. G. Heil, Dept. of Phys. & Astronomy, Univ. of Georgia, Athens, GA 30602
180. Dr. F. A. Herrero, Code 692, NASA Goddard, Greenbelt, MD 20771
181. Dr. R. A. Hess, Lab. for Plasma Research, Univ. of Maryland, College Park, MD 20742-3511
182. Dr. K. W. Hill, Plasma Physics Lab., Princeton Univ., P.O. Box 451, Princeton, NJ 08544
183. Dr. E. Hinnov, Plasma Physics Lab., Princeton Univ., Princeton, NJ 08544
184. Dr. J. R. Hiskes, L-630, LLNL, P.O. Box 5511, Livermore, CA 94550
185. Dr. E. B. Hooper, Jr., Microwave Tokamak Experiment, LLNL, P.O. Box 5511, L-537, Livermore, CA 94550
186. Dr. Ki-Ki Hosea, Dept. of Physics, Texas A&M Univ., College Station, TX 77843-4242
187. Dr. A. M. Howald, 02/518, General Atomics, P.O. Box 85608, San Diego, CA 92186-9704
188. Dr. W. F. Huebner, Southwest Research Inst., Instrum. & Space Research Div., 6220 Culebra Rd., San Antonio, TX 78228-0510
189. Dr. R. Hulse, Plasma Physics Lab., Princeton Univ., P.O. Box 451, Princeton, NJ 08543
190. IFS/FRC Library, c/o Saralynn Stewart, Fusion Research Center, The Univ. of Texas, Austin, TX 78712-1081
191. International Research & Evaluation, Attn: IRE-ITTD Input Processing Dir., 21098 IRE Control Ctr., Eagan, MN 55121
192. Dr. M. Ishimoto, The Johns Hopkins Univ., Applied Physics Lab., Johns Hopkins Rd., Laurel, MD 20723-6099
193. Dr. F. C. Jahoda, CTR Div., Los Alamos National Lab., P.O. Box 1663, MS F-639, Los Alamos, NM 87545
194. JILA Information Center, University of Colorado, Campus Box 440, Boulder, CO 80309-0440
195. Prof. R. E. Johnson, Dept. of Nuclear Eng. and Eng. Physics, Thornton Hall, Univ. of Virginia, Charlottesville, VA 22903-2442
196. Dr. R. R. Johnston, Science Applications, Inc., Suite B-31, 5150 El-Camino Real, Los Altos, CA 94022
197. Dr. E. G. Jones, Systems Research Labs., 2800 Indian Ripple Rd., Dayton, OH 45440-3696
198. Dr. K. W. Jones, DAS-Bldg. 815, Brookhaven National Lab., Upton, NY 11973
199. Dr. W. D. Jones, Dept. of Physics, Univ. of South Florida, Tampa, FL 33620
200. Dr. T. Kammash, Dept. of Nuclear Eng., Univ. of Michigan, Ann Arbor, MI 48104
201. Dr. H. P. Kelly, Jesse W. Beams Lab. of Physics, Univ. of Virginia, McCormick Rd., Charlottesville, VA 22901
202. Dr. R. A. Kenefick, Dept. of Physics, Texas A&M Univ., College Station, TX 77843-4242
203. Dr. Y. K. Kim, Bldg. 221, Rm A-267, NIST, Gaithersburg, MD 20899
204. Dr. J. L. Kohl, MS 50, Center for Astrophysics, 60 Garden St., Cambridge, MA 02138
205. Dr. R. A. Krakowski, Los Alamos National Lab., P.O. Box 1663, MS F-647, Los Alamos, NM 87545
206. Dr. G. L. Kulcinski, Fusion Technology Inst., 1500 Johnson Dr., Univ. of Wisconsin, Madison, WI 53706
207. Dr. W. B. Kunkel, Lawrence Berkeley Lab., MS 4-230, Univ. of California, Berkeley, CA 94720
208. Dr. R. Kurucz, Smithsonian Astrophys. Obs., Harvard Univ., 60 Garden St., Cambridge, MA 02138
209. Dr. T. Kvale, Dept. of Physics, Univ. of Toledo, Toledo, OH 43606
210. Dr. N. F. Lane, Rice Univ., P.O. Box 1892, Houston, TX 77251-1892
211. Dr. W. Langer, Princeton Plasma Physics Lab., P.O. Box 451, Princeton, NJ 08543

212. Lawrence Livermore National Lab., Attn: Magnetic Fusion Energy Library, L-440, P.O. Box 808, Livermore, CA 94550
213. Lawrence Livermore National Lab., Attn: Tech. Infor. Dept. Library, L-53, P.O. Box 808, Livermore, CA 94550
214. Lawrence Livermore National Lab., Univ. of California, P.O. Box 808, MS L-298, Livermore, CA 94550
215. Dr. S. A. Lawton, Mail Code 111-1041, McDonnell-Douglas Research Labs., P.O. Box 516, St. Louis, MO 63166
216. Dr. N. H. Lazar, 01-1280, TRW Space & Technology Group, One Space Park, Redondo Beach, CA 90278
217. Dr. Y. Lee, LLNL, P.O. Box 808, MS L-298, Livermore, CA 94550
218. Dr. K. N. Leung, MFE Group, Bldg. 4, Lawrence Berkeley Lab., 1 Cyclotron Rd., Berkeley, CA 94720
219. Dr. F. Levinton, Princeton Univ., Princeton Plasma Physics Lab., P.O. Box 451, Princeton, NJ 08544
220. Dr. A. Lietzke, Lawrence Berkeley Lab., 1 Cyclotron Rd., Bldg. 4, Berkeley, CA 94720
221. Dr. C. D. Lin, Dept. of Physics, Cardwell Hall, Kansas State Univ., Manhattan, KS 66506
222. Dr. C. C. Lin, Dept. of Physics, Univ. of Wisconsin, Madison, MI 53706
223. Dr. J. L. Linsky, JILA, Campus Box 440, Univ. of Colorado, Boulder, CO 80309-0440
224. Dr. A. Li-Scholz, Dept. of Physics, State Univ. of New York, Albany, NY 12222
225. Dr. G. J. Lockwood, Div. 9312, Sandia National Labs., P.O. Box 5800, Albuquerque, NM 87185
226. Dr. D. C. Lorents, Molecular Physic Lab., SRI International, 333 Ravenwood Ave., Menlo Park, CA 94025
227. Los Alamos National Lab., Attention: Librarian, MS P362, P.O. Box 1663, Los Alamos, NM 87545
228. Ms. H. W. Lovitts, Sci. Information Service, Inc., 7 Woodland Ave., Larchmont, NY 10538
229. Dr. S. R. Lundeen, Univ. of Notre Dame, Dept. of Physics, College of Science, Notre Dame, IN 46556
230. Dr. K. MacAdam, Dept. of Physics & Astronomy, Univ. of Kentucky, Lexington, KY 40506
231. Dr. D. M. Manos, Plasma Physics Lab., Princeton Univ., P.O. Box 451, Princeton, NJ 08543
232. Dr. S. T. Manson, Dept. of Physics, Georgia State Univ., Atlanta, GA 30303
233. Dr. J. V. Martinez, ER-141, G-337 GTN, Division of Chemical Sciences, Off. of Basic Energy Sci, U.S. Dept. of Energy, Washington, DC 20545
234. Dr. E. D. Mays, Sandia National Lab., Technical Library, Org. 3144, P.O. Box 5800, Albuquerque, NM 87185
235. Dr. J. W. McConkey, Dept. of Physics, Univ. of Windsor, Windsor, Ontario, Canada N9B 3P4
236. Dr. D. McCune, Princeton Plasma Physics Lab., James Forrestal Campus, P.O. Box 451, Princeton, NJ 08544
237. Prof. E. W. McDaniel, School of Physics, Georgia Inst. of Technology, Atlanta, GA 30332
238. Dr. R. McFarland, 309 Christy Dr., Rolla, MO 65401
239. Dr. R. T. McGrath, Organization 1128, Sandia National Lab., P.O. Box 5800, Albuquerque, NM 87185
240. Dr. E. J. McGuire, 1271, Sandia National Labs., Albuquerque, NM 87185
241. Dr. J. H. McGuire, Dept. of Physics, Kansas State Univ., Manhattan, KS 66506
242. Mr. E. A. McLean, Code 4780.1 EM, Naval Research Lab., Washington, DC 20375-5000
243. Dr. T. A. Mehlhorn, Div. 1262, Sandia National Labs., P.O. Box 5800, Albuquerque, NM 87185
244. Dr. A. L. Merts, Los Alamos National Lab., T-4, MS B-212, P.O. Box 1663, Los Alamos, NM 87545
245. Dr. D. R. Mikkelsen, B-148, Plasma Physics Lab., Princeton Univ., P.O. Box 451, Princeton, NJ 08543
246. Dr. G. H. Miley, Univ. of Illinois, 214 Nucl. Eng. Lab., 103 S. Goodwin Ave., Urbana, IL 61801-2984
247. Mirror Fusion Experimental Group, Lawrence Livermore National Lab., P.O. Box 808, Livermore, CA 94550
248. Dr. J. Molitoris, Lawrence Livermore National Lab., P.O. Box 808, Livermore, CA 04550
249. Dr. A. A. Mondelli, SAIC, P.O. Box 1303, 1710 Goodridge Dr., McLean, VA 22102
250. Dr. H. W. Moos, Physics & Astronomy, Johns Hopkins Univ., Baltimore, MD
251. Dr. T. J. Morgan, Dept. of Physics, Wesleyan Univ., Middletown, CT 06457-6036
252. Dr. K. Moses, Mgr., JAYCOR, Plasma Technology Division, 3547 Voyager St., Suite B-104, Torrance, CA 90503-1667
253. Prof. J. R. Mowat, Dept. of Physics, North Carolina State Univ., P.O. Box 8202, Raleigh, NC 27695-8202
254. Dr. D. W. Norcross, JILA, Campus Box 440, Univ. of Colorado, Boulder, CO 80309-0440
255. Dr. K. W. Ogilvie, Code 692, Goddard Space Flight Center, Greenbelt, MD 20771
256. Dr. R. E. Olson, Dept. of Physics, Univ. of Missouri, Rolla, Rolla, MO 65401
257. Dr. J. E. Osher, Lawrence Livermore National Lab., Univ. of California, P.O. Box 808, L-637, Livermore, CA 94550

258. Dr. J. T. Park, Vice Chancellor, Physics Dept., 102 Physics, Univ. of Missouri - Rolla, Rolla, MO 65401-0249
259. Dr. J. M. Peek, Group T-4, MS B-212, Los Alamos National Lab., Los Alamos, NM 87545
260. Dr. P. J. Persiani, Argonne National Laboraory, 9700 South Cass Ave., Bldg. 208, Argonne, IL 60439
261. Dr. J. R. Peterson, SRI International, Molecular Physics Lab., Menlo Park, CA 94025
262. Dr. R. Petrasso, NW16-132, MIT, Plasma Fusion Center, 167 Albany St., Cambridge, MA 02139
263. Dr. A. V. Phelps, JILA, Univ. of Colorado, Campus Box 440, Boulder, CO 80309-0440
264. Physics-Astronomy Library, 290 Dennison Bldg., Univ. of Michigan, Ann Arbor, MI 48109-1090
265. Physics-Astronomy Library, FM-15, Ms. Martha Austin, Librarian, Univ. of Washington, Seattle, WA 98195
266. Physics/Optics/Astron. Library, 374 Bausch & Lomb Bldg., Univ. of Rochester, Rochester, NY 14627
267. Dr. F. M. Pipkin, Dept. of Physics, Harvard Univ., Cambridge, MA 02138
268. Plasma Fusion Center Library, Massachusetts Inst. of Technology, NW16-153, Cambridge, MA 02139
269. Dr. D. E. Post, Jr., Plasma Physics Lab., P.O. Box 451, Princeton, NJ 08544
270. Dr. R. S. Post, Massachusetts Inst. of Techn., Plasma Fusion Center, 190 Albany St., Cambridge, MA 02139
271. Dr. K. Prelec, Accelerator Dept., Building 911B, Brookhaven National Lab., Upton, NY 11973
272. Dr. R. E. Price, Office of Fusion Energy, Office of Energy Research, Mail Station G-256, U.S. Dept. of Energy, Washington, DC 20545
273. Princeton Univ., Library/James Forrestal Campus, Plasma Physics Lab., P.O. Box 451, Princeton, NJ 08543
274. Dr. R. V. Pyle, Bldg. 4-230, Lawrence Berkeley Lab., Berkeley, CA 94720
275. Dr. Niansheng Qi, Lab. of Plasma Studies, 369 Upson Hall, Cornell Univ., Ithaca, NY 14853
276. Dr. B. H. Quon, Applied Microwave Plasma Concepts, Inc., 2075-N Corte del Nogal, Carlsbad, CA 92009
277. Dr. D. J. Rej, Los Alamos National Lab., CTR-1, MS K-638, P.O. Box 1663, Los Alamos, NM 87545
278. Dr. T. Renk, Div. 1264, Sandia National Labs., Albuquerque, NM 87185
279. Dr. J. S. Risley, Dept. of Physics, North Carolina State Univ., Raleigh, NC 27695-8202
280. Dr. J. R. Roberts, A-167, Physics Bldg., NIST, Gaithersburg, MD 20899
281. Dr. E. C. Roelof, Johns Hopkins Univ., Applied Physics Lab., Johns Hopkins Rd., Laurel, MD 20723-6099
282. Dr. A. B. Ross, Radiation Chem. Data Center, Radiation Lab., Univ. of Notre Dame, Notre Dame, IN 46556
283. Prof. J. Reese Roth, Univ. of Tennessee, 409 Ferris Hall, Knoxville, TN 37996-2100
284. Dr. W. L. Rowan, Fusion Research Center, Univ. of Texas, Austin, TX 78712
285. Dr. M. E. Rudd, Dept. Physics and Astronomy, Univ. of Nabraska, Lincoln, NE 68588-0111
286. Prof. D. N. Ruzic, Univ. of Illinois, 214 Nuclear Eng. Lab., 103 S. Goodwin Ave., Urbana, IL 61801
287. Sandia National Lab., Attn: Techn. Library, P.O. Box 5800, Albuquerque, NM 87185-5800
288. Dr. G. Schilling, Plasma Physics Lab., Princeton Univ., P.O. Box 451, Princeton, NJ 08543
289. Dr. A. S. Schlachter, Lawrence Berkeley Lab., MS 46-161, Univ. of California, Berkeley, CA 94720
290. Dr. H. A. Schuessler, Dept. of Physics, Texas A&M Univ., College Station, TX 77843-4242
291. Science & Vie Magazine, P.O. Box 4373, Grand Central Station, New York, NY 10163-4373
292. Dr. M. Seidl, Stevens Inst. Techn., Dept. of Physics, Hoboken, NJ 07030
293. Dr. A. Sen, Univ. Toledo, 2801 W. Bancroft St., Phys. & Astron., Toledo, OH 43606
294. Dr. S. Shafroth, Dept. of Physics, Univ. of North Carolina, Chapel Hill, NC 27514
295. Dr. D. E. Shemansky, Lunar & Planetary Lab., West, Gould-Simpson Bldg., The Univ. of Arizona, Tucson, AZ 85621
296. Dr. J. Sherman, AT-10, MS 818, Los Alamos National Lab., Los Alamos, NM 87545
297. Prof. H. A. Shugart, Dept. of Physics, Univ. of California, Berkeley, CA 94720
298. Dr. C. H. Skinner, Plasma Physics Lab., Princeton Univ., P.O. Box 451, James Forrestal Campus, Princeton, NJ 08543
299. Dr. R. Slater, Avco-Everett Research Lab., 2385 Revere Beach Parkway, Everett, MA 02149
300. Dr. J. N. Smith, Mail Stop L-520, General Atomics, P.O. Box 85608, San Diego, CA 92138-5608
301. Dr. P. L. Smith, Center for Astrophysics-50, 60 Garden St., Cambridge, MA 02138
302. Dr. L. G. Sobotka, Dept. of Chem., Washington Univ., Campus Box 1134, St. Louis, MO 63130-4899
303. Prof. J. O. Sprott, Plasma Physics, Univ. of Wisconsin, 1150 Univ. Ave., Madison, WI 53706
304. Dr. S. K. Srivastava, Jet Propulsion Lab., MS 183-601, California Inst. Techn., 4800 Oak Grove Dr., Pasadena, CA 91109
305. Dr. K. R. Stalder, SRI International, PS-019, 333 Ravenswood Ave., Menlo Park, CA 94025
306. Dr. J. W. Stearns, MS 5-119, Lawrence Berkeley Lab., 1 Cyclotron Rd., Berkeley, CA 94720

307. Prof. R. F. Stebbings, Dept. Space Phys. and Astronomy, Rice Univ., P.O. Box 1892, Houston, TX 77251
308. Prof. R. A. Stern, Univ. of Colorado, Dept. Astrophysic Planetary & Atmospheric Sciences, Campus Box 391, Boulder, CO 80309
309. Dr. B. Stratton, Princeton Plasmas Physics Lab., P.O. Box 451, Princeton, NJ 08543
310. Dr. S. Suckewer, Plasma Physics Lab., Princeton Univ., James Forrestal Campus, P.O. Box 451, Princeton, NJ 08544
311. Dr. W. C. Stwalley, Dept. of Chem., Univ. of Iowa, Iowa City, IA 52242
312. Dr. H. Takahashi, Brookhaven National Lab., Bldg. 130, Upton, Long Island, NY 11973
313. Dr. J. A. Tanis, Western Michigan Univ., Dept. of Physics, Kalamazoo, MI 49008-5151
314. Technical Information Center, Stone & Webster Engineering Corp., 3 Executive Campus, P.O. Box 5200, Cherry Hill, NJ 08034
315. Dr. D. M. Thomas, 02/527, General Atomics, 3550 General Atomics Ct., San Diego, CA 92121-1194
316. Dr. E. W. Thomas, School of Physics, Georgia Inst. of Technology, Atlanta, GA 30332
317. Dr. L. H. Toburen, Battelle Northwest Labs, P.O. Box 999 (P8-47), Richland, WA 99352
318. Dr. B. Van Zyl, Dept. of Physics, Univ. of Denver, Denver, CO 80208
319. Dr. P. Vitello, Science Applications, Inc., 1710 Goodridge Dr., P.O. Box 1303, McLean, VA 22102
320. Prof. G. C. Vlases, Aerospace and Energetics Research Program, FL-10, Univ. of Washington, Seattle, WA 98195
321. Dr. S. E. Von Goeler, Plasma Physics Lab., Princeton Univ., P.O. Box 451, Princeton, NJ 08544
322. Dr. G. K. Walters, Dept. of Physics, Rice Univ., P.O. Box 1892, Houston, TX 77251
323. Dr. P. G. Weber, Group CTR 1, MS F-648, Los Alamos National Lab., Los Alamos, NM 87545
324. Dr. W. L. Wiese, NIST, Atomic & Plasma Radiation Div., Gaithersburg, MD 20899
325. Dr. T. G. Winter, Dept. of Physics, Pennsylvania State Univ., Wilkes-Barre Campus, Lehman, PA 18627-0217
326. John G. Wolbach Library, Attention: Librarian, Harvard College Observatory, 60 Garden St., Cambridge, MA 02138
327. Dr. J. T. Woo, Interscience, Inc., 105 Jordan Rd., Troy, NY 12180
328. Dr. T. T. Yang, Rocketdyne Div., D/631-FA38, Rockwell International Corp., 6633 Canoga Park, Canoga Park, CA 91303
329. Dr. C. E. Young, CHM 200, Argonne National Lab., 9700 South Cass Ave., Argonne, IL 60439
330. Dr. A. R. Zander, Dept. of Physics, East Texas State Univ., Commerce, TX 74528
331. Prof. R. N. Zare, Dept. of Chem., Stanford Univ., Stanford, CA 94305
332. Dr. B. Zygelman, Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138

EXTERNAL DISTRIBUTION - FOREIGN

333. Dr. V. A. Abramov, Inst. Atommoi Energii, I. V. Kurchatova, 46 Ulitsa Kurchatova, Moscow D-182, U.S.S.R.
334. V. V. Afrosimov, A. F. Ioffe Phys.-Techn. Inst., Leningrad, 194021, U.S.S.R.
335. Dr. I. Alvarez-Torre and Dr. C. Cisneros, Inst. de Fisica, Univ. Nacional Autonoma de Mexico, Apartado Postal 139-B, Cuernavaca, Morelos 62191, Mexico
336. Dr. N. O. Andersen, Inst. of Physics, Univ. of Aarhus, DK-8000 Aarhus C, Denmark
337. Dr. K. Ando, Atomic Process Lab., Rikagaku Kenkyusho, The Inst. of Phys. & Chem. Res., Wako-Shi, Saitama, 351-01, Japan
338. Prof. J. Andra, LAGRIPPA, CEA-CNRS, CEN-Grenoble, Ave. des Martyrs 85X, F-38041 Grenoble, France
339. Dr. R. J. Armstrong, Univ. of Tromso, Dept. of Physics, P.O. Box 953, N 9001 Tromso, Norway
340. Dr. S. Aynacioglu, Technische Univ. Berlin, Inst. fur Strahl. und Kernphys., Sekr. PN 3-2, Hardenbergstr. 36, D-1000 Berlin, Germany
341. Dr. M. Bacal, Laboratoire P.M.I., Ecole Polytechnique, 91128 Palaiseau Cedex, France
342. Dr. J. L. Bahr, Physics Dept., Univ. of Otago, Box 56 Dunedin, New Zealand
343. Dr. A. Barany, Manne Siegbahn Inst., Frescativagen 24, S-104 05 Stocholm, Sweden
344. Dr. M. Barat, LCAM Bat 351, Univ. Paris-Sud, 91405-Orsay Cedex, France
345. Dr. H. Behrens, Fachinformationszentrum, Karlsruhe mbH, FA III, 7514 Eggenstein-Leopoldshafen 2, West Germany

346. Dr. R. Behrisch, Max-Planck-Institut fur Plasmaphysik, Boltzmannstr. 2, D-8046 Garching, B. Munchen, West Germany
347. Dr. D. Berenyi, ATOMKI, Debrecen, P.F. 51. H-4001, Hungary
348. Biblioteca - Att. Clelia De Palo, ENEA-Centro Ricerche Energia Frascati, C.P. 65, 00044 Frascati, Roma, Italy
349. Dr. Xu Bing, Head, Information Center, Southwestern Inst. of Phys., P.O. Box 16, Leshan, Sichuan, People's Republic of China
350. Dr. S. L. Bliman, Centre d'Etudes Nucleaires de Grenoble, 85 X Ave. Des Martyrs, F-38041 Grenoble, Cedex, France
351. Dr. P. Bogen, Forschungszentrum Julich GMBH, Postfach 1913, D-5170 Julich 1, West Germany
352. Dr. J. S. Briggs, Fakultaet fuer Phys., Universitaet Freiburg, D-7800 Freiburg 1 BR., Germany
353. Dr. F. Brouillard, Dept. of Physics, Catholic Univesity of Louvain, Chemin du Cyclotron. 2, B 1348 Louvain-La-Neuve, Belgium
354. Fr. Weidemann's Buchhandlung, (H. Witt) Fachbuchandlung, Fur Technische Literatur, Georgstr. 11, Postfach 64 06, D-3000 Hannover 1, West Germany
355. Miss S. Burgess, Librarian, Physics Library, Blackett Lab., Prince Consort Rd., London, S.W. 7 2B2, United Kingdom
356. Dr. R. I. Campeanu, Atomic Physics Group, Dept. of Physics, Univ. of Cluj, Str. Kogalniceanu NR.1, 3400 Cluj-Napoca, Romania
357. Dr. X. Carlotti, CEA - Vautours, BP7, 77181 Courtry, France
358. Dr. Li Shi Chang, Inst. of Appl. Phys. and Computational Math., P.O. Box 8009, Beijing, The People's Republic of China
359. Dr. L.F.S. Coelho, Univ. Fed. Do Rio De Janeiro, Centro de Cienc. Matem. da Natur., Inst. de Fisica, Caixa Postal 68.528, 21945 Rio de Janeiro, RJ, Brazil
360. Dr. J.N.L. Connor, Dept. of Chem., Univ. of Manchester, Manchester M13 9PL, England
361. Dr. J. L. Cook, Aaec Research Establishment, Physics Division, AAEC - Lucas Heights, Sutherland 2232, N.S.W., Australia
362. Culham Lab., The Library, A.E.A. Technology, Abingdon, Oxfordshire OX14 3DB, England
363. Data & Planning Center, Attn: Librarian, National Inst. for Fusion Sci., Nagoya 464-01, Japan
364. Dr. F. J. de Heer, FOM-Instituut Voor Atoom - en Molecuul Fysica, Postbus 41883, 1009 DB Amsterdam, The Netherlands
365. Prof. J. L. Delcroix, Lab. de Physique des Gaz et des Plasmas, Batiment 212, Univ. de Paris-Sud, 91405 Orsay Cedex, France
366. Dr. W. Demtroeder, Fachbereich Physik, Universitaet Kaiserslautern, Postfach 3049, D-6750 Kaiserslautern, West Germany
367. Prof. K. T. Dolder, Dept. of Atomic Physics, School of Physics, Univ. of Newcastle Upon Tyne, Newcastle-on-Tyne NE1 7RU, United Kingdom
368. Dr. G. Drake, Dept. of Physics, Univ. of Windsor, Windsor, Ont. N9B 3P4, Canada
369. Dr. H. W. Drawin, Association Euratom-CEA, D.R.F.C., Centre d'Etudes Nucleaires de Cadarache, F-13108 St-Paul-lez-Durance, France
370. Dr. P. D. Dumont, Univ. De Liege, Inst. De Phys. Nucl. Exp. B-15, Sart Tilman B-4000 Liege 1, Belgium
371. Dr. K. F. Dunn, Dept. of Pure and Applied Physics, The Queen's Univ. of Belfast, Belfast BT 7 1NN, United Kingdom
372. Dr. W. Eckstein, Bereich Oberflachenphysik, Max-Planck-Inst. fur Plasmaphysik, Boltzmannstr. 2, D-8046 Garching Bei, Munchen, West Germany
373. Dr. M. Fabbriconi, Biblioteca - ENEA, C.P. 65, 00044 - Frascati (RM), Italy
374. Dr. P. Focke, Centro Atomico Bariloche, 8400 San Carlos de Bariloche, Argentina
375. Dr. A. Fukuda, Electrotechnical Lab., 1-1-4 Umezono, Tsukuba-shi, Ibaraki, 305, Japan
376. Prof. H. B. Gilbody, Dept. of Pure & Applied Phys., The Queen's Univ. of Belfast, Belfast BT 7, 1NN, Northern Ireland
377. Dr. I. N. Golovin, I. V. Kurchatov Inst., Atomic Energy, Ulitsa Kurchatove 46, Moscow 123182, U.S.S.R.
378. Dr. E. A. Gurtovenko, Main Astronomical Observatory of the Academy of Sci. Ukr. USSR, 252127 Kiev, U.S.S.R.
379. Dr. A. A. Haasz, Univ. of Toronto, Inst. for Aerospace Studies, 4925 Dufferin St., Downsview, Ontario, Canada M3H 5T6
380. Dr. M. Hack, Dept. of Astronomy, Via Tiepolo 11, I 34131 Trieste, Italy

381. Dr. L. Hage, Library Adm. Dept., Technische Hogeschool, Bibliotheek, 2120.41233, Postbus 513, 5600-MB Eindhoven, The Netherlands
382. Dr. K. I. Hamed, Univ. of Strathclyde, Dept. of Physics, John Anderson Bldg., Glasgow G4 ONG, Scotland, United Kingdom
383. Dr. J. E. Hansen, Zeeman Labs., PL Muidergracht 4, NL-1018TV Amsterdam, The Netherlands
384. Dr. M.F.A. Harrison, Culham Lab., UKAEA, Abingdon Oxfordshire, OX 14 3DB England
385. Dr. J. D. Hey, Dept. of Physics, Univ. of Cape Town, Rondebosch 7700, South Africa
386. Dr. R. Hoekstra, K. V. I., Univ. of Groningen, Zernikelaan 25, 9747 AA Groningen, The Netherlands
387. Dr. Keh-Ning Huang, Inst. of Atomic and Molecular Science, Academia Sinica, P.O. Box 23-166, Taipei, Taiwan 107
388. Prof. Dr. B. A. Huber, CEN-G SPhAt, Ave. des Martyrs, 38041 Grenoble Cedex, France
389. Dr. J. G. Hughes, Dept. of Computer Science, The Queen's Univ. of Bel, Belfast BT7 1NN, Northern Ireland
390. Dr. W. N. Hugrass, Dept. of Physics, Univ. of New England, Armidale NSW 2351, Australia
391. Dr. P. Hvelplund, Inst. of Physics, Univ. of Aarhus, DK 8000 Aarhus C, Denmark
392. Inst. for Plasma Research, Attn: Librarian, BHAT, Gandhinagar 382 424, India
393. Dr. Y. Itikawa, Inst. of Space, and Astronautical Science, 3-1-1 Yoshinodai, Sagamihara 229, Japan
394. Dr. R. K. Janev, Nuclear Data Section, International Atomic Energy Agency, Wagramerstrasse 5, P.O. Box 100, A-1400 Vienna, Austria
395. Dr. C. Joachain, Institut De Physique, Sciences 1, Universite De Louvain, B1348 Louvain-La-Neuve, Belgium
396. Dr. P. B. Johnson, Dept. of Physics, Victoria Univ. of Wellington, P.O. Box 600, Wellington, New Zealand
397. Dr. E. M. Jones, Jet Joint Undertaking, Codas Div. J2, Abingdon, Oxfordshire OX14 3EA, United Kingdom
398. Dr. T. Kato, Data & Planning Center, National Inst. for Fusion Science, Nagoya 464-01, Japan
399. Dr. K. Katsonis, Lab. Physique Des Plasmas, Faculte Des Sci. D'orsay, Batiment 212, F-91405 Orsay Cedex, France
400. Kernforschungszentrum Karlsruhe GMBH, Literaturabteilung, Postfach 3640, D-7500 Karlsruhe, Federal Republic of Germany
401. Prof. A. E. Kingston, Dept. Appl. Math. & Theor. Phys., The Queen's Univ., Belfast BT7 1NN, Northern Ireland
402. Dr. H. Knudsen, Inst. of Physics, Univ. of Aarhus, DK-8000, Aarhus C, Denmark
403. Dr. G. A. Kobzen, Inst. for High Temperatures, USSR Acad. of Science, 127912, Izhorskaya Str. 13/19, Moscow, USSR
404. Dr. K. Kondo, Plasma Physics Lab., Kyoto Univ., Gokasho Uji, Kyoto, Japan
405. Dr. N. Konjevic, Inst. of Physics, 11001 Beograd, P.O. Box 57, Yugoslavia
406. Dr. R. I. Kostik, Main Astronomical Observatory of the Academy of Sci. Ukr., 252127, Kiev, USSR
407. Dr. Ales Krejci, Czechoslovak Academy of Sciences, Inst. of Plasma Physics, Pod Vodarenskou vezi 4, 18211 Prague 8, Liben, Czechoslovakia
408. Dr. L. I. Krupnik, Inst. of Phys. & Techn., The Ukrainian Academy of Sci., Kharkov - 310108, U.S.S.R.
409. Dr. H. Kubo, Diagnostics Div., Dept. Large Tokamak Res., Naka Fusion Research Establishment, Naka-machi, Ibaraki 311-02, Japan
410. Prof. Dr. H.-J. Kunze, Inst. fuer Exp.-Physics V, Ruhr-Universitaet, Postfach 10 21 48, D-4630 Bochum 1, West Germany
411. Prof. B. Lehnert, The Royal Inst. of Techn., Div. of Plasma Phys. Fusion Res., Stockholm, 70, Sweden
412. Dr. W. N. Lennard, Solid State Science, Chem. & Mat. Div., Atomic Energy of Canada Limited, Nuclear Labs., Chalk River, Ontario, Canada K0J 1JO
413. Prof. F. Linder, Dept. of Physics, Univ. of Kaiserslautern, D-6750 Kaiserslautern, West Germany
414. Dr. A. Lorenz, Deputy Head, Nuclear Data Section, International Atomic Energy Agency, Wagramerstrasse 5, P.O. Box 100, A-1400 Vienna, Austria
415. Prof. M. Los, FOM-Inst. of Atomic and Molecular Physics, Kruislaan 407, Amsterdam/Watergraafsmeer, The Netherlands
416. Dr. M. W. Lucas, Univ. of Sussex, School of Math. and Physical Sciences, Falmer, Brighton BN1 9GH, United Kingdom
417. Dr. Y. Maron, Dept. Nuclear Physics, K. B. Weissman Inst. of Phys. Sci., The Weizmann Inst. of Sci., Rehovot 76 100, Israel

418. Dr. I. Martinson, Fysiska Institutionen, Lunds Universitet, Soelvagatan 14, 223 62 Lund, Sweden
419. Dr. D. Mathur, Tata Inst. of Fundamental Research, Homi Bhabha Rd., Bombay - 400 005, India
420. Dr. S. Matsuda, Fusion Research Center, Japan Atomic Energy Res. Inst., Mukaiyama, Naka-machi, Ibaraki 311-01, Japan
421. Dr. M. Mattioli, Assoc. Euratom-CEA sur la Fusion, Dept. de Recherches sur la Fusion Centrolle, Centre d'Etudes Nucl. de Cadarache, F-13108 St. Paul lez durance, Cedex, France
422. Max-Planck-Inst. F. Plasmaphys., E3, Mrs. A. Hohaus, Librarian, 8046 Garching Bei, Muenchen, West Germany
423. Dr. H. M. Mayer, Max-Planck-Inst. fuer Plasmaphys., E3, 8046 Garching, West Germany
424. Dr. J. W. McConkey, Dept. of Physics, Univ. of Windsor, Windsor, Ontario N9B 3P4, Canada
425. Dr. E. Minguez, Inst.o de Fusion Nuclear, c/Jose G. Abascal, 2, 28006 Madrid, Spain
426. Dr. S. V. Mirnov, I. V. Kurchatov Inst. of Atomic Energy, Ulitsa Kurchatova, 46, Moscow 123182, U.S.S.R.
427. Dr. J.B.A. Mitchell, Physics Dept., Univ. of Western Ontario, London, Ontario, Canada N6A 3K7
428. Dr. A. Miyahara, National Inst. for Fusion Science, Nagoya 464-01, Japan
429. Prof. R. Morgenstern, K.V.I., Univ. of Groningen, Zernikelaan 25, 9747 AA Groningen, The Netherlands
430. Dr. A. Muller, Strahlenzentrum, der Justus Liebig-Universitaet, Leihgesterner Weg 217, D-6300 Giessen, West Germany
431. Dr. Y. Nakai, Laser Atomic Separation Eng., Research Association of Japan, Tokai-mura Ibaraki 319-11, Japan
432. Dr. S. Nakazaki, Dept. of Applied Physics, Faculty of Eng., Miyazaki Univ., Miyazaki 889-21, Japan
433. Prof. H. Narumi, Fukuyama Univ., Fukuyama, 729-02, Japan
434. Dr. H. Nguyen, Lab. de Spectroscopie des Plasmas, Tour 22, Univ. Paris Vi, 4 Place Jussieu, 05 Cedex, 75252 Paris, France
435. Prof. Dr. H. Nussbaumer, Inst. of Astronomy, ETH Zentrum, CH-8092 Zurich, Switzerland
436. Dr. K. Ohno, National Center Sci. Information System, 3-29-1 Otsuka Bunkyo-ku, Tokyo 112, Japan
437. Dr. S. Ohtani, National Inst. for Fusion Science, Nagoya 464-01, Japan
438. Prof. A. Osman, Theoretical Nuclear Physics, Dept. of Physics, Faculty of Science, Cairo Univ., Cairo, Egypt
439. Dr. M. Otsuka, National Inst. for Fusion Science, Nagoya 646-01, Japan
440. Dr. W. Ott, Max-Planck-Inst. fuer Plasmaphysik, D-8046 Garching Bei, Muenchen, West Germany
441. Dr. E. S. Parilis, Inst. of Electronics, Tashkent, U.S.S.R.
442. Dr. O. S. Pavlichenko, Inst. of Phys. & Techn., The Ukrainian Academy of Sci., Kharkov - 310108, U.S.S.R.
443. Dr. N. J. Peacock, Culham Lab., UKAEA Research Group, Abingdon Oxfordshire OX 14 3DB, England
444. Dr. M. P. Petrov, A. F. Ioffe Phys.-Technical Inst., Leningrad K-21, U.S.S.R.
445. Physical Research Lab., Librarian, Plasma Physics Programme, Room No. 563, Navrangpura, Ahmedabad-380 009, India
446. Physics and Applied Maths Library, College Park, The Queen's Univ. of Belfast, Belfast BT 7 1NN, Northern Ireland
447. Dr. L. P. Presnyakov, P. N. Lebedev Physical Inst., USSR Academy of Science, Leninsky Prospect 53, 117924 Moscow, USSR
448. Dr. J. Radet (Dowmeutalist), Dept. de Recherches sur la Fusion Controlee BT 506, 13108 St. Paul lez, Durance Cedex, France
449. Research Information Center, National Inst. for Fusion Science, Nagoya 464-01, Japan
450. Dr. H. Ringler, 1 E 3, Max-Planck-Inst. fuer Plasmaphys., Boltzmannstr. 2, D-8046 Garching Bei, Munchen, West Germany
451. Dr. A. C. Riviere, A.E.A. Technology, Culham Lab., Abingdon, Oxfordshire 0X14 3DB, United Kingdom
452. Dr. H. Rohr, Max-Planck-Inst. fuer Plasmaphys., Boltzmann Str. 3, D-8046 Garching Bei, Munchen, West Germany
453. Prof. Dr. Z. Rudzikas, Inst. of Physics, K. Pozelos 54, 232600, Vilnius, Lithuanian SSR, USSR
454. Rutherford Appleton Lab., Attn: Librarian, Chilton, Didcot, Oxfordshire OX11 0QX, United Kingdom
455. Dr. U. I. Safronova, Inst. of Spectroscopy, USSR Academy of Science, Troitzk, Moscow Region, 142092 USSR
456. Dr. J. Sahade, Casilla de Correo 677, Observatorio Astronomico, 1900 La Plata, Argentina
457. Prof. E. Salzborn, Strahlenzentrum, der Justus Liebig-Universitaet, Leihgesterner Weg 217, D-6300 Giessen, West Germany
458. Dr. A. Sanchez, Calle Angostura Quinta-Carmen, El Marques, Caracas 1070, Venezuela

459. Prof. K. H. Schartner, I. Physikalisches Inst., Justus-Liebig Univ., Heinrich-Buff-Ring 16, D-6300 Giessen, West Germany
460. Dr. J. J. Schmidt, Chief, Nuclear Data Section, International Atomic Energy Agency, Wagramerstrasse 5, P.O. Box 100, A-1400 Vienna, Austria
461. Dr. V. S. Senashenko, Inst. of Nuclear Physics, M. V. Lomonosov Moscow State Univ., Moscow 119899, U.S.S.R.
462. Dr. A. Surjalal Sharma, Inst. for Plasma Research, BHAT, Gandhinagar 382 424, India
463. Dr. V. B. Sheorey, 461 Physical Research Lab., Navrangpura, Ahmedabad 380 009, India
464. Dr. I. Shimamura, Riken, Hirosawa, Wako-Shi, Saitama 351-01, Japan
465. Dr. T. Shirai, Dept. of Physics, Japan Atomic Energy Research Inst., Tokai-Mura, Ibaraki 319-11, Japan
466. Dr. R. Shuker, Dept. of Physics, Ben-Gurion Univ. of The Negev, P.O. Box 653, Beer Sheva 84 105, Israel
467. Prof. B. M. Smirnov, Inst. of Thermophysics, Siberian Branch of The USSR, Academy of Sciences, PR. AC. Lavrentyeva, 1, 630090 Novosibirsk - 90, U.S.S.R.
468. Prof. F. J. Smith, Dept. Computer Science, The Queen's Univ., Belfast BT7 1NN, Northern Ireland
469. Dr. J. J. Smith, Nuclear Data Section, International Atomic Energy Agency, Wagramerstrasse 5, P.O. Box 100, A-1400 Vienna, Austria
470. Prof. B. L. Stansfield, Univ. du Quebec, Inst. National de la Recherche Scientifique, Case Postale 1020, Varennes, Quebec, Canada J3X 1S2
471. Dr. H. P. Summers, Jet Joint Undertaking, Abingdon, Oxon, OX14 3EA, England
472. Dr. H. Suzuki, Dept. of Physics, Sophia Univ., Kioicho 7-1, Chiyoda-ku, Tokyo 102, Japan
473. Dr. D. R. Sweetman, United KingdomA.E.A., Culham Lab., Bldg. E-5, Abingdon, Oxfordshire OX14 3DB, United Kingdom
474. Prof. K. Takayanagi, Oji-Honcho 1-3-10, Kita-ku, Tokyo 114, Japan
475. Prof. H. Tawara, Data & Planning Center, National Inst. for Fusion Science, Nagoya 464-01, Japan
476. Dr. V. I. Tereshin, Inst. of Phys. & Techn., The Ukrainian Academy of Sci., Kharkov - 310108, U.S.S.R.
477. Univ. de Bordeaux I, Lab. des Collisions Atomiques, 351 Cours de la Liberation, 33405 Talence Cedex, France
478. Universitaetsbibliothek and Techn., Informationsbibliothek, Welfengarten 1 B, D-3000 Hannover 1, West Germany
479. Dr. E. Veje, Physics Lab., Univ. of Copenhagen, H. C. Orsted Inst., Universitetsparken 5, DK 2100 Copenhagen, Denmark
480. Dr. H. Verbeek, Max-Planck-Institut fuer Plasmaphys., 8046 Garching Bei, Munchen, West Germany
481. Dr. V. S. Voitsenya, Inst. of Phys. & Techn., The Ukrainian Academy of Sci., Kharkov 310108, U.S.S.R.
482. Prof. H. Walther, Sektioa Physik der Ludwig, Maximilians-Univ. Muenchen, Am. Coulombwall 1, D-8046 Garching, West Germany
483. Dr. Hongzhong Wang, Dept. of Physics, Huaihai Univ., Lianyungang, Jiangsu Province, The People's Republic of China
484. Prof. K. Wiesemann, Ruhr-Univ. Bochum, Postfach 102148, D-4630 Bochum 1, West Germany
485. Dr. J. F. Williams, Dept. of Physics, The Univ. of Western Australia, Nedlands Perth 6009, Australia
486. Prof. Dr. H. Winter, Inst. fur Allgemeine Physik, Der Technischen Univ. Wien, A-1040 Wien, Wiedner Hauptstr. 8-10, Austria
487. Dr. Z. Xu, Shanghai Inst. of Optics & Fine Mechanics, Academia Sinica, P.O. Box 800-211, Shanghai, 201800, The People's Republic of China
488. Prof. F. Yang, Inst. of Modern Physics, Fudan Univ., Shanghai, The People's Republic of China
489. Dr. Jin-Zhang Yao, Chinese Nuclear Data Center/A+M, Inst. of Atomic Energy, P.O. Box 275 (41), Beijing, The People's Republic of China
490. Mahroos Yaseen, Dept. of Physics, Univ. of Strathclyde, John Anderson Bldg., 107 Rottenrow, Glasgow G4 ONG, Scotland, U. K.
491. Dr. E. C. Zhou, Chinese Nuclear Data Center, Inst. of Atomic Energy, P.O. Box 275 (41), Beijing, The People's Republic of China
492. Dr. V. Zoita (IFTAR-L22), Central Inst. of Physics, R-76900, Magurele Bucharest, P.O. Box MG-7, Romania

493-
502. Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831

503-
552. Controlled Fusion Atomic Data Center