

ornl

OAK RIDGE
NATIONAL
LABORATORY

MARTIN MARIETTA


ORNL/TM-11559



3 4456 0349391 2

ASYMPTOTIC BEHAVIOR OF SOLUTIONS OF DIFFUSION-LIKE PARTIAL DIFFERENTIAL EQUATIONS INVARIANT TO A FAMILY OF AFFINE GROUPS

Lawrence Dresner



OPERATED BY
MARTIN MARIETTA ENERGY SYSTEMS, INC.
FOR THE UNITED STATES
DEPARTMENT OF ENERGY

This report has been reproduced directly from the best available copy.

Available to DCE and DCE contractors from the Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831; prices available from (615) 576-8401, FTS 623-8401.

Available to the public from the National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Rd., Springfield, VA 22161.
NTIS price codes--Printed Copy: A03 Microfiche A01

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

ORNL/TM-11559
Dist. Category UC-420

Fusion Energy Division


**ASYMPTOTIC BEHAVIOR OF SOLUTIONS OF
DIFFUSION-LIKE PARTIAL DIFFERENTIAL
EQUATIONS INVARIANT TO A FAMILY
OF AFFINE GROUPS**

Lawrence Dresner

Date published: July 1990

Prepared for the
Office of Fusion Energy
Budget Activity No. 6130 66 90 1

Prepared by the
OAK RIDGE NATIONAL LABORATORY
Oak Ridge, Tennessee 37831
operated by
MARTIN MARIETTA ENERGY SYSTEMS, INC.
for the
U.S. DEPARTMENT OF ENERGY
under contract DE-AC05-84OR21400



3 4456 0349391 2

CONTENTS

ABSTRACT	v
INTRODUCTION	1
FORM OF THE PARTIAL DIFFERENTIAL EQUATION	2
ORDERED SOLUTIONS	2
THE SUPERFLUID DIFFUSION EQUATION	3
THE THERMAL EXPULSION EQUATION	5
THE POROUS MEDIUM EQUATION	6
DISCUSSION	6
REFERENCES	7
Appendix A. MOST GENERAL FORM OF THE PARTIAL DIFFERENTIAL EQUATION	9
Appendix B. INVARIANCE OF THE PRINCIPAL ORDINARY DIFFERENTIAL EQUATION	11

ABSTRACT

This report deals with the asymptotic behavior of certain solutions of partial differential equations in one dependent and two independent variables (call them c , z , and t , respectively). The partial differential equations are invariant to one-parameter families of one-parameter affine groups of the form

$$c' = \lambda^\alpha c ,$$

$$t' = \lambda^\beta t ,$$

$$z' = \lambda z ,$$

where λ is the group parameter that labels the individual transformations and α and β are parameters that label groups of the family. The parameters α and β are connected by a linear relation,

$$M\alpha + N\beta = L ,$$

where M , N , and L are numbers determined by the structure of the partial differential equation.

It is shown that when L/M and N/M are <0 , certain solutions become asymptotic to $z^{L/M}t^{-N/M}$ for large z or small t . Some practical applications of this result are discussed.

INTRODUCTION

In two earlier publications^{1,2} I studied the properties of certain partial differential equations with one dependent and two independent variables (call them c , z , and t , respectively). These partial differential equations are invariant to one-parameter families of one-parameter affine groups of the form

$$\left. \begin{aligned} c' &= \lambda^\alpha c \\ t' &= \lambda^\beta t \\ z' &= \lambda z \end{aligned} \right\} \quad 0 < \lambda < \infty, \quad (1)$$

where λ is the group parameter that labels the individual transformations and α and β are parameters that label groups of the family. The parameters α and β are connected by a linear relation,

$$M\alpha + N\beta = L, \quad (2)$$

where M , N , and L are numbers determined by the structure of the partial differential equation. Because of relation (2), only one of the two parameters α and β may be chosen freely.

Similarity solutions are solutions of the partial differential equation that are invariant to one group of the family, say, that for which $\alpha = \alpha_0$ and $\beta = \beta_0$. Such solutions most generally have the form

$$c = t^{\alpha_0/\beta_0} y(z/t^{1/\beta_0}), \quad (3)$$

where y is a function of the single variable $x = z/t^{1/\beta_0}$. When substituted into the partial differential equation, Eq. (3) yields an ordinary differential equation for y called the principal ordinary differential equation. The form of the principal ordinary differential equation depends on both the form of the partial differential equation and the values of α_0 and β_0 .

Much attention has been paid in refs. 1 and 2 to diffusion-like partial differential equations such as $c_t = (c^m c_z)_z$ (the so-called porous medium equation), $cc_t = c_{zz}$ (which describes thermal expulsion of a compressible liquid from a long, slender, heated tube), and $c_t = (c_z^{1/3})_z$ [which describes heat transport in turbulent superfluid helium (He-II)]. Among the interesting solutions of these equations are those that obey the boundary and initial conditions

$$c(\infty, t) = 0, \quad (4a)$$

$$c(z, 0) = 0. \quad (4b)$$

To define a solution completely, an additional boundary condition is necessary. If it takes the form

$$c(0, t) = At^n, \quad (5)$$

where A and n are constants, then the solution is a similarity solution of the form (3). Equations (4a) and (4b) then collapse to the single condition $y(\infty) = 0$.

Many detailed calculations of similarity solutions of the three partial differential equations mentioned above for various α_0 and β_0 showed that quite often, when $L/M < 0$, the function y approaches zero as x approaches ∞ asymptotically to $ux^{L/M}$, where u is a constant. When substituted into Eq. (3), this gives the asymptotic form $uz^{L/M}t^{-N/M}$ for c , which would obey the conditions (4a) and (4b) if L/M and N/M were both < 0 . Demonstrating that c has the asymptotic form $uz^{L/M}t^{-N/M}$ in the special cases studied was quite laborious, and it is the purpose of this report to outline broad conditions under which this asymptotic form can be verified if not at a glance, then at least with a minimum of computational effort.

FORM OF THE PARTIAL DIFFERENTIAL EQUATION

The partial derivative c_t transforms under transformation (1) according to $c'_t = \lambda^{\alpha-\beta}c_t$; similarly, all other partial derivatives transform by multiplication by some power of λ . If a partial differential equation involving $z, t, c, c_t, c_z, c_{zz}, \dots$ is to be invariant to all groups of the family (1), then it can only contain λ -less combinations of $z, t, c, c_t, c_z, c_{zz}$, etc. Here the term “ λ -less” is to be understood as the term “dimensionless” is understood in ordinary dimensional analysis. Thus the partial differential equation must have the form

$$F\left(\frac{c}{z^{L/M}t^{-N/M}}, \frac{tc_t}{c}, \frac{zc_z}{c}, \frac{t^2c_{tt}}{c}, \frac{z^2c_{zz}}{c}, \dots\right) = 0, \quad (6)$$

where F can be any function.

It is easy enough to see that Eq. (6) is invariant to Eq. (1): if we imagine Eq. (6) to be written in terms of the primed variables and substitute for them from Eq. (1), we obtain Eq. (6) again in the unprimed variables. It is proved in Appendix A that only forms composed of λ -less terms have this property.

Can this partial differential equation have solutions of the form

$$c = uz^at^b? \quad (7)$$

Direct substitution into Eq. (6) shows that Eq. (7) can only be a solution when $a = L/M$, $b = -N/M$, and u satisfies the equation

$$F\left[u, -\frac{N}{M}, \frac{L}{M}, \frac{N}{M}\left(\frac{N}{M} + 1\right), \frac{L}{M}\left(\frac{L}{M} - 1\right), \dots\right] = 0. \quad (8)$$

If Eq. (8) has real solutions for u , then the partial differential equation has real solutions of the form Eq. (7); if not, it has none. If $L/M < 0$ and $N/M < 0$, then Eq. (7) is capable of representing the asymptotic limit of solutions obeying the boundary and initial conditions (4a) and (4b). But under what conditions *must* Eq. (7) represent this limit?

ORDERED SOLUTIONS

For the superfluid diffusion equation and the thermal expulsion equation, it can be shown that for solutions $c_1(z, t)$ and $c_2(z, t)$ that obey the boundary and initial

conditions (4a) and (4b), if $c_1(0, t) > c_2(0, t)$, then $c_1(z, t) \geq c_2(z, t)$ for all z . Such solutions are thus ordered according to their values at $z = 0$.

One consequence of such ordering is that for any similarity solutions, the function y obeys the inequalities

$$0 \leq y(x) \leq ux^{L/M} \quad (9)$$

as long as $y(0)$ is finite. Here u is the smallest real solution of Eq. (8). Let us now consider the functions $y(x)$ belonging to the values α_0 and β_0 of α and β . These functions are a one-parameter family ordered according to their intercepts $y(0)$ on the y -axis. In refs. 1 and 2, it is shown that the principal ordinary differential equation for the functions $y(x)$ is invariant to the affine group*

$$\left. \begin{aligned} y' &= \mu^{L/M} y \\ x' &= \mu x \end{aligned} \right\} 0 < \mu < \infty . \quad (10)$$

So each of the curves $y(x)$ is the image of any other because $y'(0) = \mu^{L/M} y(0)$ can be given any value by the appropriate choice of μ while $y'(\infty) = \mu^{L/M} y(\infty)$ remains zero. Thus, the entire family of these similarity solutions $y(x)$ is transformed into itself.

Because of Eq. (10),

$$\frac{y'(x')}{ux'^{L/M}} = \frac{\mu^{L/M} y(x)}{u(\mu x)^{L/M}} = \frac{y(x)}{ux^{L/M}} , \quad (11)$$

so that the limits as x and x' approach infinity of the left- and right-hand sides are the same. Suppose this limit < 1 . The infinitude of curves $y(x)$, being bounded from above, has an upper limit $y_\infty(x)$. This limit, too, is a solution of the principal ordinary differential equation. Furthermore, because the entire family is invariant to Eq. (10), its upper limit $y_\infty(x)$ must also be invariant to Eq. (10). Curves invariant to Eq. (10) must have the form $y = vx^{L/M}$, where v is a constant. Since this invariant curve also satisfies the principal ordinary differential equation, it must correspond to a solution $c = vz^{L/M}t^{-N/M}$ of the partial differential equation. But then v must be a root of Eq. (8). Since $v < u$, and u is the smallest root of Eq. (8), we have a contradiction. Therefore, the limit of both sides of Eq. (11) as x and x' approach infinity must be 1. This means that all the similarity solutions have the asymptotic form $uz^{L/M}t^{-N/M}$.

Because of the ordering of the solutions of the partial differential equations, this conclusion holds as well for any solution $c(z, t)$ that obeys the boundary and initial conditions (4a) and (4b) and whose value at the origin $c(0, t)$ is bounded above and below by powers of t .

THE SUPERFLUID DIFFUSION EQUATION

We now investigate the ordering of the solutions of the superfluid diffusion equation,

$$c_t = (c_z^{1/3})_z . \quad (12)$$

*A different proof from that given in refs. 1 and 2 can be found in Appendix B.

To do so, we consider the infinitesimal difference δc between two neighboring solutions. It obeys the linear partial differential equation obtained by taking the first variation of Eq. (12), namely,

$$(\delta c)_t = \left[\frac{1}{3} c_z^{-2/3} (\delta c)_z \right]_z, \quad (13a)$$

which can be rearranged as

$$3c_z^{2/3} u_t = -\frac{2}{3} \frac{c_{zz}}{c_z} u_z + u_{zz} \quad (13b)$$

after setting $\delta c = u$. The difference u obeys the boundary and initial conditions

$$u(0, t) > 0, \quad (14a)$$

$$u(\infty, t) = 0, \quad (14b)$$

$$u(z, 0) = 0. \quad (14c)$$

The conditions (14b) and (14c) follow from the fact that both neighboring solutions obey the boundary and initial conditions (4a) and (4b). To prove the ordering of the two solutions, we must prove that $u(z, t) \geq 0$ for all z and t .

To avoid difficulties created by semi-infinite domains, let us begin by replacing Eq. (14b) by

$$u(L, t) = 0 \quad (14d)$$

and restricting ourselves to the rectangle R of length L along the z -axis and length T along the t -axis. We propose to prove that $u \geq 0$ in R by proving that the smallest value of u must lie on one of the sides S_1 : ($z = 0, 0 \leq t \leq T$), S_2 : ($t = 0, 0 \leq z \leq L$), or S_3 : ($z = L, 0 \leq t \leq T$). On S_1, S_2 , and S_3 , the smallest value of u is zero.

If Eq. (13b) were replaced by the strict differential inequality

$$3c_z^{2/3} u_t > -\frac{2}{3} \frac{c_{zz}}{c_z} u_z + u_{zz}, \quad (15)$$

we could easily prove that the smallest value of u could not be attained either in the interior of R or at an interior point of side S_4 : ($t = T, 0 \leq z \leq L$). Suppose, for example, that the smallest value of u were attained at a point P in the interior of R . The point P would then be a relative minimum at which $u_z(P) = u_t(P) = 0$ and $u_{zz}(P) \geq 0$. But these stipulations contradict Eq. (15). Suppose, instead, that the smallest value of u were attained at an interior point P of S_4 . Then the point P would be a relative minimum along S_4 so that at P , $u_z(P) = 0$ and $u_{zz}(P) \geq 0$. Since $c_z^{2/3}$ is always positive no matter what the sign of c_z , Eq. (15) then implies that $u_t > 0$; this means that yet smaller values of u lie inside R directly under point P , again a contradiction. Since the smallest value of u cannot be attained either in R or on S_4 , it must lie on S_1, S_2 , or S_3 .

We can convert Eq. (13b) into the strict inequality (15) by adding a small, positive source term ϵ to the right-hand side of Eq. (13b). If we assume that δc is

a continuous function of ϵ , then $\lim_{\epsilon \rightarrow 0} \delta c \geq 0$ in R . It is worth noting that this limiting process dilutes the strict inequality $\delta c > 0$ in R , which is what we have actually proved when $\epsilon > 0$, to the weaker inequality $\delta c \geq 0$ in R , but the latter is sufficient for our purposes.

Finally, we let L approach infinity and so return from the boundary condition (14d) to the boundary condition (14b).

THE THERMAL EXPULSION EQUATION

The thermal expulsion equation

$$cc_t = c_{zz} \tag{16}$$

has as its first variation the following equation for $u = \delta c$:

$$uc_t + cu_t = u_{zz} , \tag{17}$$

which we again consider in the rectangle R with the boundary and initial conditions (14a), (14c), and (14d). As before, we can convert Eq. (17) to a strict inequality by adding an infinitesimal positive source term ϵ to its right-hand side. We cannot be sure of the sign of c_t , and the standard trick for dealing with such uncertainty is to set

$$u = ve^{\lambda t} , \tag{18}$$

in which case Eq. (17) becomes

$$(c_t + \lambda c)v + cv_t = v_{zz} + \epsilon e^{-\lambda t} , \tag{19}$$

where v , too, obeys the boundary and initial conditions (14a), (14c), and (14d). If $c \geq \delta > 0$ in R , we can choose λ large enough so that $c_t + \lambda c > 0$ in R . Then if the minimum of v were attained at a point P in R , P would be a relative minimum at which $v(P) \leq 0$, $v_t(P) = 0$, and $v_{zz}(P) \geq 0$. But these stipulations contradict Eq. (19). If the minimum of v were attained at an interior point P of S_4 , then $v_{zz}(P) \geq 0$, $v(P) \leq 0$, so that from Eq. (19) we find that $v_t(P) > 0$. Thus, yet smaller values of v lie inside R , again a contradiction. Since the smallest value of v must then lie on S_1 , S_2 , or S_3 , $v > 0$ in R . In the limit as ϵ approaches zero, this strict inequality weakens to $v \geq 0$ in R . In view of Eq. (18), this is equivalent to $u \geq 0$ in R , which was to be proved.

All of this depends on showing that $c \geq \delta$ in the interior of R and on S_4 . This will be so if $c(0, t) \geq \delta$. As before, we add a small, positive source term ϵ to Eq. (16) and also replace the right-hand sides of Eqs. (14a), (14c), and (14d) written for c with a small, positive quantity δ . Later, we shall let ϵ and δ approach 0.

The smallest value of c cannot occur in the interior of R . If it did, say at a point P , then P would be a relative minimum and $c_t(P) = 0$ and $c_{zz}(P) \geq 0$. These two requirements contradict Eq. (16) augmented by the source term ϵ .

The smallest value of c cannot occur at an interior point of S_4 either. But now, owing to the factor c (of uncertain sign on S_4) on the left-hand side of Eq. (16), we cannot prove this with the argument of the previous section. But let us consider

the rectangle R' : ($0 \leq z \leq L$, $0 \leq t \leq T'$), where $T' \gg T$, and let $c(0, t)$ take the fixed value $c(0, T)$ for $T \leq t \leq T'$. If T' is large enough, then on S'_4

$$c(z, T') = \frac{z}{L} \delta + \frac{L-z}{L} c(0, T) , \quad (20)$$

which is the steady solution that $c(z, t)$ approaches when $c(0, t)$ is constant. The smallest value of c cannot occur in R' , as just shown, and surely does not occur along the interior of S'_4 , where c is given by Eq. (20). Hence, it must occur on S'_1 , S'_2 , or S'_3 , which means that $c > \delta$ in R' and thus in R . In the limit as ϵ approaches zero, we find $c \geq \delta$ in R .

By the argument previously given, $v \geq 0$ in R for any value of $\delta > 0$. In the limit as δ approaches zero, $v \geq 0$ in R . Thus $u \geq 0$ in R , which proves the ordering of the solutions. Finally, as before, we let L approach infinity.

THE POROUS MEDIUM EQUATION

Both of the two partial differential equations just discussed have L/M and $N/M < 0$ as required for the solution (7) to fulfill the boundary and initial conditions (4a) and (4b). For the porous medium equation

$$c_t = (c^m c_z)_z , \quad (21)$$

$L/M = 2/m$ and $N/M = 1/m$, so that these two ratios can only be negative if $m < 0$. In many applications, $m > 0$, so that solutions obeying the boundary and initial conditions (4a) and (4b) cannot have asymptotic limits of the form (7). At least some of the similarity solutions of Eq. (21) for $m > 0$ are known to vanish at and beyond certain finite, time-dependent values of z (Refs. 3-5).

DISCUSSION

The limiting processes used in demonstrating the ordering of the solutions of the superfluid diffusion equation and the thermal expulsion equation are based on unproven assumptions of continuity. For certain linear partial differential equations related to the ordinary heat diffusion equation, rigorous proofs exist that do not depend on such assumptions.⁶ These proofs are quite lengthy and involved; what is more, they are not always easy to generalize for use with nonlinear partial differential equations. Therefore, I prefer the heuristic approach given here, even though it does not conform to a high standard of rigor.

The importance of the results proved here rests on the fact that the asymptotic form $c \sim uz^{L/M}t^{-N/M}$ is simple and independent of the boundary value $c(0, t)$. This form can therefore be used without paying an exorbitant cost in computation; furthermore, it can be relied on even when the boundary value $c(0, t)$ is uncertain. An excellent example of such an application is the protection of superconducting magnets wound with cable-in-conduit conductors.⁷ A local normal zone in such a conductor induces flow in the helium, the velocity of which can be modeled by the thermal expulsion equation.⁸ Since such conductors are typically very long

compared with their hydraulic diameters ($L/D \sim 10^5$), the induced velocity in the helium at the end of a hydraulic path can be estimated using the asymptotic form given above. Two data of great practical utility that can be obtained in this way are the velocity of expulsion from the ends of the tube and the time at which thermal hydraulic quenchback⁹ is complete. The expulsion velocity is large enough in many practical situations to be used as a nonelectrical means of quench detection.¹⁰ The phenomenon of thermal hydraulic quenchback strongly affects the maximum quench pressure in the conductor as well as the hot-spot temperature. Both thermal expulsion and thermal hydraulic quenchback have been studied by means of detailed calculations based on simple models that give plausible values of $c(0, t)$ (refs. 10 and 11). From the conclusions reached here, we can see that those results of the detailed calculations that are founded on the asymptotic form $uz^{L/M}t^{-N/M}$ are model independent and could have been obtained after only a few lines of calculation.

A similar conclusion applies to the experimental temperature distributions in superfluid He-II measured by van Sciver¹² and by van Sciver and Lottin.¹³ Although these two sets of measurements differ markedly in their boundary and initial conditions (van Sciver introduced a constant heat flux into a half-space; van Sciver and Lottin delivered an instantaneous heat pulse to an infinite medium), in both cases the temperature at short times and large distances is described by the asymptotic form $uz^{L/M}t^{-N/M}$, as detailed calculations show.^{14,15} With the theorems of this paper, a partial but convincing comparison of theory and experiment can be carried out with a minimum of computation.

REFERENCES

1. Lawrence Dresner, *Similarity Solutions of Nonlinear Partial Differential Equations*, Pitman Publishing, Inc., Marshfield, Mass., 1983.
2. Lawrence Dresner, "Similarity Solutions of Nonlinear Partial Differential Equations Invariant to a Family of Affine Groups," *Math. Comput. Model.* **11**, 531-534 (1988).
3. R. E. Pattle, "Diffusion from an Instantaneous Point Source with a Concentration-Dependent Coefficient," *Q. J. Mech. Appl. Math.* **12** (4), 407-409 (1959).
4. B. H. Gilding and L. A. Peletier, "On a Class of Similarity Solutions of the Porous Media Equation," *J. Math. Anal. Appl.* **55**, 351-364 (1976).
5. B. H. Gilding, "Similarity Solutions of the Porous Media Equation," *J. Hydrol.* **56**, 251-63 (1982).
6. M. H. Protter and H. F. Weinberger, *Maximum Principles in Differential Equations*, Prentice-Hall, Inc., Englewood Cliffs, N.J., 1967.
7. L. Dresner, "Superconductor Stability, 1983: A Review," *Cryogenics* **24**, 283-92 (1984).
8. Lawrence Dresner, "Thermal Expulsion of Helium from a Quenching Cable-in-Conduit Conductor," pp. 618-21 in *Proceedings of the Ninth Symposium on the Engineering Problems of Fusion Research, Chicago, IL, October 26-29, 1981*, IEEE, New York, 1982.

9. C. A. Luongo, R. J. Loyd, F. K. Chen, and S. D. Peck, "Thermal-Hydraulic Simulation of Helium Expulsion from a Cable-in-Conduit Conductor," *IEEE Trans. Magn.* **MAG-25**, 1589–95 (1989).
10. Lawrence Dresner, "Quench Detection by Fluid Dynamic Means in Cable-in-Conduit Superconductors," *Adv. Cryog. Eng.* **33**, 167–174 (1988).
11. Lawrence Dresner, "Thermal Hydraulic Quenchback in Cable-in-Conduit Superconductors," to be published.
12. S. W. van Sciver, "Transient Heat Transport in He-II," *Cryogenics* **19**, 385 (1979).
13. J. C. Lottin and S. W. van Sciver, "Heat Transport Mechanisms in a 2.3-meter-Long Cooling Loop Containing He-II," p. 269 in *Proceedings of the Ninth International Cryogenic Engineering Conference, Kobe, Japan, May 11–14, 1982*, Butterworths, Guildford, Surrey, U.K., 1982.
14. L. Dresner, "Transient Heat Transfer in Superfluid Helium," *Adv. Cryog. Eng.* **27**, 411–19 (1982).
15. Lawrence Dresner, "Transient Heat Transfer in Superfluid Helium—Part II," *Adv. Cryog. Eng.* **29**, 323–33 (1984).

APPENDIX A
MOST GENERAL FORM OF THE PARTIAL
DIFFERENTIAL EQUATION

As we have seen in the main text, the partial derivative c_t transforms under the family of groups (1) by multiplication by $\lambda^{\alpha-\beta}$: $c'_t = \lambda^{\alpha-\beta} c_t$. In view of the linear constraint (2), we can eliminate β from the exponent in favor of α ; thus $\alpha - \beta = (M/N + 1)\alpha - L/N$. So c_t transforms by multiplication by a power of λ , the power being a linear function of α . The same is true of all other partial derivatives of c . Thus, in finding the most general form of a partial differential equation involving z, t, c , and its derivatives, we are led to consider functions F of N arguments x_i that transform according to the one-parameter family of groups

$$x'_i = \lambda^{a_i\alpha + b_i} x_i, \quad 0 < \lambda < \infty. \quad (\text{A1})$$

Here a_i and b_i are constants [$(M/N + 1)$ and $-L/N$ in the case of c_t].

If the function F is invariant to all the groups of the family (A1), then

$$F(x_1, x_2, \dots, x_N) = F(x'_1, x'_2, \dots, x'_N) \quad (\text{A2})$$

or

$$F(x_1, x_2, \dots, x_N) = F(\lambda^{a_1\alpha + b_1} x_1, \lambda^{a_2\alpha + b_2} x_2, \dots, \lambda^{a_N\alpha + b_N} x_N). \quad (\text{A3})$$

Equation (A3) is an identity true for all values of α and λ . If we differentiate with respect to λ and then set $\lambda = 1$, we find

$$\sum_{i=1}^N (a_i\alpha + b_i) x_i \frac{\partial F}{\partial x_i} = 0. \quad (\text{A4})$$

Since Eq. (A4) is also an identity true for all values of α , it is equivalent to the pair of first-order, linear partial differential equations

$$\sum_{i=1}^N a_i x_i \frac{\partial F}{\partial x_i} = 0, \quad (\text{A5a})$$

$$\sum_{i=1}^N b_i x_i \frac{\partial F}{\partial x_i} = 0. \quad (\text{A5b})$$

The characteristic equations of these two partial differential equations are

$$\frac{dx_1}{a_1 x_1} = \frac{dx_2}{a_2 x_2} = \dots = \frac{dx_N}{a_N x_N}, \quad (\text{A6a})$$

$$\frac{dx_1}{b_1 x_1} = \frac{dx_2}{b_2 x_2} = \dots = \frac{dx_N}{b_N x_N}. \quad (\text{A6b})$$

The most general solution of Eq. (A5a) is an arbitrary function of the $N - 1$ integrals of Eq. (A6a), and similarly the most general solution of Eq. (A5b) is

an arbitrary function of the $N - 1$ integrals of Eq. (A6b). The integrals of either set of characteristic equations are products of powers of the x_i , that is, functions $u(x_1, \dots, x_N)$ of the form

$$u = \prod_{i=1}^N x_i^{c_i} . \quad (\text{A7})$$

Since $du = 0$ in the direction given by Eq. (A6a), if we take the logarithm of Eq. (A7) and then differentiate in that direction we find

$$0 = \frac{dx_1}{a_1 x_1} \left(\sum_{i=1}^N c_i a_i \right) , \quad (\text{A8})$$

so that the constants c_i must obey the constraint

$$\sum_{i=1}^N c_i a_i = 0 . \quad (\text{A8a})$$

Similarly, the powers c_i in the integrals of Eq. (A6b) must obey the constraint

$$\sum_{i=1}^N c_i b_i = 0 . \quad (\text{A8b})$$

If we look at Eqs. (A8a) and (A8b) as expressing the orthogonality of an N -dimensional vector c to two other vectors a and b , then we see at once that the admissible vectors span a subspace of dimension $N - 2$. Hence there are $N - 2$ independent mutual integrals of Eqs. (A6a) and (A6b). The most general simultaneous solution of Eqs. (A5a) and (A5b) is an arbitrary function of these $N - 2$ independent mutual integrals.

Now let us consider invariants of the entire family of groups of the form

$$v = \prod_{i=1}^N x_i^{c_i} . \quad (\text{A9})$$

Invariance means that

$$\prod_{i=1}^N x_i^{c_i} = \prod_{i=1}^N x_i'^{c_i} = \prod_{i=1}^N x_i^{c_i} \lambda^{(a_i \alpha + b_i) c_i} \quad (\text{A10})$$

for any choice of x_i , α , and λ . Taking logarithms, we find that

$$\ln \lambda \sum_{i=1}^N (a_i \alpha + b_i) c_i = 0 , \quad (\text{A11})$$

which leads as before to the constraints (A8a) and (A8b) on the c_i . Thus there are $N - 2$ independent invariants v of the entire family of groups, and their exponent vectors c span the same subspace as the exponent vectors of the mutual integrals of Eqs. (A6a) and (A6b). So the most general function F is then an arbitrary function of the $N - 2$ invariants v . Because of the first equality in Eq. (A10), these invariants are “ λ -less” combinations of the x_i .

APPENDIX B
INVARIANCE OF THE PRINCIPAL ORDINARY
DIFFERENTIAL EQUATION

We start by writing the “ λ -less” equation (6) in the more convenient form

$$G \left[\frac{c}{z^{L/M} t^{-N/M}}, \frac{tc_t}{c}, \frac{zc_z}{c}, \frac{t(tc_t)_t}{c}, \frac{z(zc_z)_z}{c}, \dots \right] = 0. \quad (\text{B1})$$

In view of the definition $x = zt^{-1/\beta}$, we can write the operator equations

$$t \frac{\partial}{\partial t} = -\frac{1}{\beta} x \frac{\partial}{\partial x} \quad (z \text{ held fixed}), \quad (\text{B2a})$$

$$z \frac{\partial}{\partial z} = x \frac{\partial}{\partial x} \quad (t \text{ held fixed}). \quad (\text{B2b})$$

Furthermore, we can write Eq. (3) in either of the following alternate forms:

$$c = t^{\alpha/\beta} y(x) = z^\alpha x^{-\alpha} y(x). \quad (\text{B3})$$

If we substitute Eq. (B3) into Eq. (B1), we find with the help of Eq. (B2) that Eq. (B1) takes the form

$$G \left\{ \frac{y(x)}{x^{L/M}}, -\frac{1}{\beta} \frac{x \frac{d}{dx} (x^{-\alpha} y)}{x^{-\alpha} y}, \frac{x dy}{y dx}, \frac{1}{\beta^2} \frac{x \frac{d}{dx} [x \frac{d}{dx} (x^{-\alpha} y)]}{x^{-\alpha} y}, \right. \\ \left. \frac{x \frac{d}{dx} \left(x \frac{dy}{dx} \right)}{y}, \dots \right\} = 0, \quad (\text{B4})$$

which one can immediately see is invariant to the group (10).

INTERNAL DISTRIBUTION

- | | |
|--------------------------------------|--|
| 1. D. B. Batchelor | 25. Laboratory Records, ORNL-RC |
| 2. J. B. Drake | 26. Document Reference Section |
| 3-17. L. Dresner | 27. Central Research Library |
| 18. M. S. Lubell | 28. Fusion Energy Division Library |
| 19. C. W. Nestor, Jr. | 29-30. Engineering Technology/Fusion
Energy Division Publications
Office |
| 20. M. W. Rosenthal | 31. ORNL Patent Office |
| 21. J. Sheffield | |
| 22. G. V. Wilson | |
| 23-24. Laboratory Records Department | |

EXTERNAL DISTRIBUTION

32. Office of the Assistant Manager for Energy Research and Development, U.S. Department of Energy, Oak Ridge Operations Office, Oak Ridge, TN 37831
33. J. D. Callen, Department of Nuclear Engineering, University of Wisconsin, Madison, WI 53706-1687
34. R. W. Conn, Department of Chemical, Nuclear, and Thermal Engineering, University of California, Los Angeles, CA 90024
35. N. A. Davies, Acting Director, Office of Fusion Energy, Office of Energy Research, ER-50 Germantown, U.S. Department of Energy, Washington, DC 20545
36. S. O. Dean, Fusion Power Associates, Inc., 2 Professional Drive, Suite 248, Gaithersburg, MD 20879
37. H. K. Forsen, Bechtel Group, Inc., Research Engineering, P. O. Box 3965, San Francisco, CA 94105
38. J. R. Gilleland, L-644, Lawrence Livermore National Laboratory, P.O. Box 5511, Livermore, CA 94550
39. R. W. Gould, Department of Applied Physics, California Institute of Technology, Pasadena, CA 91125
40. R. A. Gross, Plasma Research Laboratory, Columbia University, New York, NY 10027
41. D. M. Meade, Princeton Plasma Physics Laboratory, P.O. Box 451, Princeton, NJ 08543
42. M. Roberts, International Programs, Office of Fusion Energy, Office of Energy Research, ER-52 Germantown, U.S. Department of Energy, Washington, DC 20545

43. W. M. Stacey, School of Nuclear Engineering and Health Physics, Georgia Institute of Technology, Atlanta, GA 30332
44. D. Steiner, Nuclear Engineering Department, NES Building, Tibbetts Avenue, Rensselaer Polytechnic Institute, Troy, NY 12181
45. R. Varma, Physical Research Laboratory, Navrangpura, Ahmedabad 380009, India
46. Bibliothek, Max-Planck Institut für Plasmaphysik, Boltzmannstrasse 2, D-8046 Garching, Federal Republic of Germany
47. Bibliothek, Institut für Plasmaphysik, KFA Jülich GmbH, Postfach 1913, D-5170 Jülich, Federal Republic of Germany
48. Bibliothek, KfK Karlsruhe GmbH, Postfach 3640, D-7548 Karlsruhe 1, Federal Republic of Germany
49. Bibliotheque, Centre de Recherches en Physique des Plasmas, Ecole Polytechnique Federale de Lausanne, 21 Avenue des Bains, CH-1007 Lausanne, Switzerland
50. R. Aymar, CEN/Cadarache, Departement de Recherches sur la Fusion Contrôlée, F-13108 Saint-Paul-lez-Durance Cedex, France
51. Bibliothèque, CEN/Cadarache, F-13108 Saint-Paul-lez-Durance Cedex, France
52. Documentation S.I.G.N., Departement de la Physique du Plasma et de la Fusion Contrôlée, Association EURATOM-CEA, Centre d'Etudes Nucléaires, B.P. 85, Centre du Tri, F-38041 Grenoble, France
53. Library, Culham Laboratory, UKAEA, Abingdon, Oxfordshire, OX14 3DB, England
54. Library, JET Joint Undertaking, Abingdon, Oxfordshire OX14 3EA, England
55. Library, FOM-Instituut voor Plasmafysica, Rijnhuizen, Edisonbaan 14, 3439 MN Nieuwegein, The Netherlands
56. Library, National Institute for Fusion Science, Chikusa-ku, Nagoya 464-01, Japan
57. Library, International Centre for Theoretical Physics, P.O. Box 586, I-34100 Trieste, Italy
58. Library, Centro Ricerca Energia Frascati, C.P. 65, I-00044 Frascati (Roma), Italy
59. Library, Plasma Physics Laboratory, Kyoto University, Gokasho, Uji, Kyoto, Japan
60. Plasma Research Laboratory, Australian National University, P.O. Box 4, Canberra, A.C.T. 2601, Australia
61. Library, Japan Atomic Energy Research Institute, Naka Fusion Research Establishment, 801-1 Mukoyama, Naka-machi, Naka-gun, Ibaraki-ken, Japan
62. Library, Japan Atomic Energy Research Institute, Naka Research Establishment, Naka-machi, Naka-gun, Ibaraki-ken, Japan
63. P. Komarek, KfK Karlsruhe GmbH, Postfach 3640, D-7548 Karlsruhe 1, Federal Republic of Germany

64. S. Shimamoto, Japan Atomic Energy Research Establishment, Tokai Research Establishment, Tokai, Naka-gun, Ibaraki-ken 311-02, Japan

Department of Mathematics, The University of Tennessee,
Knoxville, TN 37916

65. V. Alexiades
66. J. S. Bradley
67. S. Lenhart

Department of Nuclear Engineering and Engineering Physics,
University of Wisconsin, Madison, WI 53706

68. M. M. Abdelsalam
69. R. Boom
70. M. Carbon
71. M. L. Corradini
72. Y. M. Eyssa
73. G. L. Kulcinski
74. D. C. Larbalestier
75. G. A. Moses
76. I. N. Sviatoslavsky
77. S. W. van Sciver

Swiss Institute for Nuclear Research, CH-5234 Villigen, Switzerland

78. G. Vécsey
79. J. A. Zichy

- 80–120. Given distribution as shown in DOE/OSTI-4500, Magnetic Fusion Energy
(Dist. Category UC-420)